

EPA REGISTRATION
NO.86004-4
Vol.1

DATA PACKAGE BEAN SHEET

Date: 01-Apr-2009

Page 1 of 2

Decision #: 406734

DP #: (363697)

PRIA

Parent DP #:

Submission #: 845401

*** Registration Information ***

Registration: 86004-U - GLYPHOSATE 41% SL

Company: 86004 - REPAR-GLYPHO, LLC

Risk Manager: RM 25 - James Tompkins - (703) 305-5697 Room# PY1 S-7337

Risk Manager Reviewer: Vickie Walters VWALTERS

Sent Date: _____

Calculated Due Date: 17-Sep-2009

Edited Due Date: _____

Type of Registration: Product Registration - Section 3

Action Desc: (R310.1) NEW PRODUCT;NON-FAST TRACK (INCLUDES REVIEWS OF PRODUCT CHEM)

Ingredients: _____

*** Data Package Information ***

Expedite: ☐ Yes ☒ No

Date Sent: 01-Apr-2009

Due Back: _____

DP Ingredient: _____

DP Title: Glyphosate 41% SL

CSF Included: ☒ Yes ☐ No

Label Included: ☒ Yes ☐ No

Parent DP #: _____

Assigned To

Date In

Date Out

Organization: RD / TRB

Last Possible Science Due Date: 18-Aug-2009

Team Name: CHEM

Science Due Date: _____

Reviewer Name: _____

Sub Data Package Due Date: _____

Contractor Name: _____

*** Studies Sent for Review ***

Printed on Page 2

*** Additional Data Package for this Decision ***

No Additional Data Packages

*** Data Package Instructions ***

Please review enclosed information to determine if proposed product is similar enough to 524-445 for submitted and cited data to be used for registration.

Please note source of ai is currently in TRB/Chem under DP 363572

Enclosed are comp appl, letter, data matrix, proposed and cited CSFs, proposed label, pages from cited label and MRID 47685101

MRID	MRID Status	Citation Reference	Guideline
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1550/Product Identity and composition
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1600/Description of materials used to produce the product
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1620/Description of production process
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1650/Description of formulation process
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1670/Discussion of formation of impurities
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1700/Preliminary analysis
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1750/Certified limits
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1800/Enforcement analytical method
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6302/Color
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6303/Physical state
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6304/Odor
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6313/Stability to sunlight, normal and elevated temperatures, metals, and metal ions
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6315/Flammability
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6319/Miscibility
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7000/pH of water solutions or suspensions
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7100/Viscosity
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7300/Density/relative density

Fee for Service

JK
{845401p~

This package includes the following

- ☒ New Registration
- ☐ Amendment

✓ Studies? Fee Waiver?
✓ volpay % Reduction: _____

for Division

- ☐ AD
- ☐ BPPD
- ☒ RD

Risk Mgr. 25

Receipt No.

S- 845401

EPA File Symbol/Reg. No.

86004-U

Pin-Punch Date:

2/23/2009

☐ This item is NOT subject to FFS action.

Action Code:

Requested: —

Granted: R31D.1

Amount Due: \$ ~~136.50~~ 136.5

Parent/Child Decisions:

86004-R / 8045378

per Linda Q
2/26/09

☒ Inert Cleared for Intended Use

☐ Uncleared Inert in Product

Reviewer: Venus Quejle

Date: 2/25/09

Remarks:

DATA PACKAGE BEAN SHEET

Date: 01-Apr-2009

Page 1 of 1

Decision #: 406734

DP #: (363699)

PRIA

Parent DP #:

Submission #: 845401

*** Registration Information ***

Registration: 86004-U - GLYPHOSATE 41% SL

Company: 86004 - REPAR-GLYPHO, LLC

Risk Manager: RM 25 - James Tompkins - (703) 305-5697 Room# PY1 S-7337

Risk Manager Reviewer: ~~Vickie Walters~~ VWALTERS

Sent Date: _____

Calculated Due Date: 17-S

Type of Registration: Product Registration - Section 3

Action Desc: (R310.1) NEW PRODUCT;NON-FAST TRACK (INCLUDES

Ingredients:

*** Data Package Information ***

Expedite: ☐ Yes ☒ No

Date Sent: 01-Apr

DP Ingredient: _____

DP Title: Glyphosate 41% SL

CSF Included: ☒ Yes ☐ No

Label Included: ☒ Yes ☐ No

Assigned To

Date In

Date Out

Organization: RD / TRB

Last Possible Science Due Date: 18 Aug 2009

Team Name: TOX

Science Due Date: _____

Reviewer Name: _____

Sub Data Package Due Date: _____

Contractor Name: _____

*** Studies Sent for Review ***

No Studies

*** Additional Data Package for this Decision ***

Can be printed on its own page

*** Data Package Instructions ***

Please review enclosed information to determine if proposed product is similar enough to 524-445 for cited and submitted data to support registration.

Please note, source of ai is currently in TRB/Tox under 363573.

Enclosed are comp appl, letter, data matrix, proposed and cited CSFs, proposed label and pages from cited labels

No tox data

Score 2/

IG DP 363655

Proposed: 86004-U

Cited: 524-445

BoB CSF: (02-09-09)
A/c ep. label

Similarity: Similar
to cited
product

17-06

Signature

Receipt for Section 3

S: 845401 Resubmission: ☐ Yes ☒ No

Regulatory Type: Product Registration - Section 3 Fee For Service: ☒ Yes ☐ No

Application Type: New Registration Billable: ☒ Yes ☐ No

Company: 86004 REPAR-GLYPHO, LLC V

Risk Manager: Registration Division, Risk Management Team 25

Product #: 86004-U Product Name: GLYPHOSATE 41% SL

Override: _____

Me Too Section 3: _____ Me Too Product Name: _____

Application Date: 17-Feb-2009 OPP Rec'd Date: 23-Feb-2009

Front End Date: 25-Feb-2009 Risk Manager Send Date: _____

FFS Due Date: _____ Negotiated Due Date: _____

OPP Target Date: _____

Fast Track: ☐ New Ingredient: ☐

Receipt Description:

Application for registration	524-445	New Ingredient	_____
Glyphosate 41% SL	42750-60-Active	Request Date:	_____
D 406 734	R 310.1	New Ingredient	_____
		Received Date:	_____

Form A: ☐ Signature Date: _____ Form B: ☐ Signature Date: _____

Receipt Content

Study: _____

CSF: _____

< >

View/Edit

Print Letter

Enter More Information

Tracking

para 6111.6 9/11/09

due: 9/17/09

11/11/09 363697

Please review enclosed information to determine if proposed product is similar enough to 524-445 as that it is added or deleted from support registration. Please note, source of data is currently under review. T.E. 11/11/09 under DP 363572.

Enclosed are company application, letters, data matrix, proposed and actual CSFs, proposed label, pages from actual label and 42750-60-Active.

TRB/TM DP 363699

Please review enclosed information to determine if proposed product is similar enough to 42750-60 for its active to support registration. T.E. 11/11/09 under DP 363573.

Enclosed are company app, letters, data matrix, proposed and actual CSFs, proposed label and pages from actual label.

DATA PACKAGE BEAN SHEET

Date: 01-Apr-2009

Page 1 of 2

Decision #: 406734

DP #: (363697)

PRIA

Parent DP #:

Submission #: 845401

*** Registration Information ***

Registration: 86004-U - GLYPHOSATE 41% SL

Company: 86004 - REPAR-GLYPHO, LLC

Risk Manager: RM 25 - James Tompkins - (703) 305-5697 Room# PY1 S-7337

Risk Manager Reviewer: Vickie Walters VWALTERS

Sent Date: _____

Calculated Due Date: 17-Sep-2009

Edited Due Date: _____

Type of Registration: Product Registration - Section 3

Action Desc: (R310.1) NEW PRODUCT;NON-FAST TRACK (INCLUDES REVIEWS OF PRODUCT CHEMIST

Ingredients: _____

*** Data Package Information ***

Expedite: ☐ Yes ☒ No

Date Sent: 01-Apr-2009

Due Back: _____

DP Ingredient: _____

DP Title: Glyphosate 41% SL

CSF Included: ☒ Yes ☐ No

Label Included: ☒ Yes ☐ No

Parent DP #: _____

Assigned To

Date In

Date Out

Organization: RD / TRB

Last Possible Science Due Date: 18-Aug-2009

Team Name: CHEM

Science Due Date: _____

Reviewer Name: _____

Sub Data Package Due Date: _____

Contractor Name: _____

*** Studies Sent for Review ***

Printed on Page 2

*** Additional Data Package for this Decision ***

No Additional Data Packages

*** Data Package Instructions ***

Please review enclosed information to determine if proposed product is similar enough to 524-445 for submitted and cited data to be used for registration.

Please note source of ai is currently in TRB/Chem under DP 363572

Enclosed are comp appl, letter, data matrix, proposed and cited CSFs, proposed label, pages from cited label and MRID 47685101

MRID	MRID Status	Citation Reference	Guideline
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1550/Product Identity and composition
4	101	Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1600/Description of materials used to produce the product
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1620/Description of production process
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1650/Description of formulation process
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1670/Discussion of formation of impurities
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1700/Preliminary analysis
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1750/Certified limits
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.1800/Enforcement analytical method
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6302/Color
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6303/Physical state
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6304/Odor
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6313/Stability to sunlight, normal and elevated temperatures, metals, and metal ions
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6315/Flammability
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.6319/Miscibility
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7000/pH of water solutions or suspensions
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7100/Viscosity
47685101		Mandava, N. (2009) Glyphosate 415 SL: Product Properties Data in Support of FIFRA Registration. Project Number: REPAR/GLYPHO41SL/5. Unpublished study prepared by Repar-Glypho, LLC. 65 p.	830.7300/Density/relative density

DATA PACKAGE BEAN SHEET

Date: 01-Apr-2009

Page 1 of 1

Decision #: 406734

DP #: (363699)

PRIA

Parent DP #:

Submission #: 845401

*** Registration Information ***

Registration: **86004-U - GLYPHOSATE 41% SL**

Company: 86004 - REPAR-GLYPHO, LLC

Risk Manager: RM 25 - James Tompkins - (703) 305-5697 Room# PY1 S-7337

Risk Manager Reviewer: Vickie Walters VWALTERS

Sent Date: _____

Calculated Due Date: 17-Sep-2009

Edited Due Date: _____

Type of Registration: Product Registration - Section 3

Action Desc: (R310.1) NEW PRODUCT;NON-FAST TRACK (INCLUDES REVIEWS OF PRODUCT CHEMIS

Ingredients: _____

*** Data Package Information ***

Expedite: ☐ Yes ☒ No

Date Sent: 01-Apr-2009

Due Back: _____

DP Ingredient: _____

DP Title: Glyphosate 41% SL

CSF Included: ☒ Yes ☐ No

Label Included: ☒ Yes ☐ No

Parent DP #: _____

Assigned To

Date In

Date Out

Organization: RD / TRB

Last Possible Science Due Date: 18-Aug-2009

Team Name: TOX

Science Due Date: _____

Reviewer Name: _____

Sub Data Package Due Date: _____

Contractor Name: _____

*** Studies Sent for Review ***

No Studies

*** Additional Data Package for this Decision ***

Can be printed on its own page

*** Data Package Instructions ***

Please review enclosed information to determine if proposed product is similar enough to 524-445 for cited and submitted data to support registration.

Please note, source of ai is currently in TRB/Tox under 363573.

Enclosed are comp appl, letter, data matrix, proposed and cited CSFs, proposed label and pages from cited labels



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 86004-U	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Glyphosate 41% SL	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Repar-Glypho, LLC 8070 Georgia Avenue, Suite 209 Silver Spring, MD. 20910 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 524-445 Product Name Roundup Original Herbicide	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

- 1) Submission of revised draft label with corrections in support of registration for Glyphosate 41% SL (EPA Reg. No.: 86004-U)
- 2) Please direct all correspondence to: Mandava Associates, LLC; 1050 Connecticut Avenue, NW / Suite 1000; Washington, D.C. 20036
- 3) The contact Person fax number is (202) 223 - 0141 and e-mail address is: mandava@compuserve.com.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	
				<input checked="" type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 20 Kg, 50 Kg		5. Location of Label Directions <input checked="" type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name N. Bhushan Mandava		Title Agent for Repar-Glypho, LLC		Telephone No. (Include Area Code) (202) 223-1424	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature <i>N. Bhushan Mandava</i>		3. Title Agent for Repar-Glypho, LLC			
4. Typed Name N. Bhushan Mandava		5. Date 09/16/09			

1 Reg.

P. 43 T.

46 "

106 ~~Test~~ purpose

MASTER LABEL FOR EPA REG. NO. 86004-U

Registered Brand Names:
GLYPHOSATE 41% SL

Table of Contents for Master Label

12wd -

09-17-08

E. Wilson

	Main Label for Food Crop Uses	Page 2
	Main Label for Industrial, Turf, & Ornamental Uses	Page 70
	Supplemental Labeling	Page 100

See each label part for more detailed table of contents

I. MAIN LABEL FOR FOOD CROP USES

GLYPHOSATE 41% SL

Herbicide

Complete Directions for Use EPA Reg. No. 86004-U

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Herbicide for Roundup Ready® Crops.

Selective broad-spectrum weed control in Roundup Ready crops. Non-selective, broad-spectrum weed control for many agricultural systems and farmsteads.

Not all products on this label are registered for use in California. Check the registration status of each product in California before using.

Read the entire label before using this product.

Use only according to label instructions.

[Container Label Statement (5 gallons or smaller):]

Do not use this container for residential treatments.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC REPACKAGING OR TOLL REPACKAGING AGREEMENT.

Non-refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

CONTENTS

1.	1.0	INGREDIENTS
2.	2.0	IMPORTANT PHONE NUMBERS
3.	3.0	PRECAUTIONARY STATEMENTS
	3.1	Hazards to Humans and Domestic Animals
	3.2	Environmental Hazards
	3.3	Physical or Chemical Hazards
4	4.0	STORAGE AND DISPOSAL
5	5.0	GENERAL INFORMATION (How This Product Works)
6	6.0	Weed Resistance Management
	6.1	General Weed Management Recommendations
	6.2	Management Recommendations for Glyphosate Resistant Biotype
7	7.0	MIXING
	7.1	Mixing with Water
	7.2	Tank Mixtures
	7.3	Tank Mixing Procedures
	7.4	Mixing for Hand-Held Sprayers
	7.5	Surfactants
	7.6	Ammonium Sulfate
	7.7	Colorants or Dyes
	7.8	Drift Reduction Additives
8	8.0	APPLICATION EQUIPMENT AND TECHNIQUES
	8.1	Aerial Equipment
	8.2	Ground Broadcast Equipment
	8.3	Hand-Held or Backpack Equipment
	8.4	Selective Equipment
	8.5	Injection Systems 8.6 CDA Equipment
9	9.0	ANNUAL AND PERENNIAL CROPS (Alphabetical)
	9.1	Cereal and Grain Crops
	9.2	Corn
	9.3	Cotton
	9.4	Fallow Systems
	9.5	Grain Sorghum (Milo)
	9.6	Herbs and Spices
	9.7	Oil Seed Crops
	9.8	Soybeans
	9.9	Sugarcane
	9.10	Vegetable Crops 9.11 Miscellaneous Crops
10	10.0	TREE, VINE, AND SHRUB CROPS (Alphabetical)
	10.1	Berry Crops
	10.2	Citrus
	10.3	Miscellaneous Tree Food Crops
	10.4	Non-Food Tree Crops
	10.5	Pome Fruit

- 10.6 Stone Fruit
- 10.7 Tree Nuts
- 10.8 Tropical and Subtropical Trees and Fruits
- 10.9 Vine Crops
- 11 11.0 **PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS**
 - 11.1 Alfalfa, Clover, and Other Forage Legumes
 - 11.2 Conservation Reserve Programs (CRP)
 - 11.3 Grass Seed or Sod Production
 - 11.4 Pastures
 - 11.5 Rangelands
- 12 12.0 **ROUNDUP READY® CROPS**
 - 12.1 Roundup Ready® Alfalfa
 - 12.2 Roundup Ready® Canola (Spring Varieties)
 - 12.3 Roundup Ready® Canola (Winter Varieties)
 - 12.4 Roundup Ready® Corn 2 (Supplemental Labeling for Corn 2)
 - 12.5 Roundup Ready® Gene Cotton
 - 12.6 Roundup Ready® Flex Cotton
 - 12.7 Roundup Ready® Gene Soybeans
 - 12.8 Roundup Ready® Gene Sugar Beets
- 13 13.0 **NON-CROP USES AROUND THE FARMSTEAD**
 - 13.1 General Weed Control and Trim-And-Edge
 - 13.2 Greenhouse/Shade house
 - 13.3 Chemical Mowing
 - 13.4 Cut Stumps
 - 13.5 Habitat Management
- 14 14.0 **ANNUAL WEEDS RATE SECTION**
 - 14.1 Annual Weeds - Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K
 - 14.2 Annual Weeds - Hand-Held or Backpack Equipment
 - 14.3 Annual Weeds - Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems
- 15 15.0 **PERENNIAL WEEDS RATE SECTION**
- 16 16.0 **WOODY BRUSH AND TREES RATE SECTION**
- 17 17.0 **LIMIT OF WARRANTY AND LIABILITY**

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....41.0%

OTHER INGREDIENTS:.....59.0%
100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

2.0 IMPORTANT PHONE NUMBERS

(To be completed later)

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Precautions:

3.2 Environmental Hazards

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and transported in stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR TRANSPORT THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Supplemental labeling may be found on the cdms.net or greenbook.net websites or obtained by contacting your authorized Repar-Glypho, LLC, retailer or Repar-Glypho, LLC, Company representative.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

[Container Label Storage and Disposal Statements]:

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact Repar-Glypho, LLC. Company at 1-301-562-7330.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

Refiners: Do not reuse this container except for refill in accordance with a valid Repar-Glypho, LLC. Agreement for Repackaging or Toll Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Repar-Glypho, LLC. container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

Product Description: This product is a postemergence, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds effects may not be visible for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" for specific weed recommendations.

Always use the higher product application rate within the range when weed growth is heavy or dense or when weeds are growing in an undisturbed (non-cultivated) area.

Reduced weed control may result from treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by this herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates: The maximum application or use rates stated throughout this product's labeling are given in units of volume (fluid ounces, pints, or quarts) of this product per acre. However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as text mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of this product (6 pounds of glyphosate acid) per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product (8 pounds of glyphosate acid) per acre per year.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

6.0 WEED RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE
-------	---	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using cultural and mechanical practices.

6.1 General Weed Management Recommendations

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management recommendations:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready crops without rotation of crops is to rotate to other Roundup Ready crops to allow use of alternate tank mixtures.
- Utilize the label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Repar-Glypho, LLC. representative, local retailer, or county extension agent.

6.2 Management For Glyphosate Resistant Biotypes

Note: Contact your Repar-Glypho, LLC. representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

The following agronomic practices may be used to reduce the spread of confirmed or unconfirmed glyphosate resistant weed biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops that allows use of herbicides that controls the resistant biotypes.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes:

7.0 MIXING

Spray solutions of this product should be mixed, stored and applied using only clean stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: PRODUCT PERFORMANCE MAY BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

7.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Begin filling the mixing tank or spray tank with clean water. Add the amount of this product near the end of the filling process and mix gently. Use caution to avoid siphoning back into the carrier source. During mixing, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

7.2 Tank Mixtures This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and follow all label directions of all products in the tank mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not in this labeling, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC. for this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT for applications of this product unless otherwise noted in this product label, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC..

When a tank mixture with a generic active ingredient, such as diuron, atrazine, 2,4-D or dicamba, as in this label, the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank mixture.

Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture, and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Use according to the most restrictive label requirements.

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance. For best results, tank mixtures with this product be applied at a minimum spray volume rate of 10 gallons per acre.

7.3 Tank Mixing Procedure

Prepare tank mixtures of this product as follows:

1. Place a 20- to 35-mesh screen or wetting basket over the filling port of the tank.
2. Through the screen, fill the tank one-half full with water and start gentle agitation.
3. If ammonium sulfate is to be used, add it slowly through the screen into the tank, and continue adding water into the tank through the screen. If dry ammonium sulfate is being used, ensure that it is completely dissolved in the tank before adding other products.
4. If a wettable powder is used, make a slurry with the water, and add it SLOWLY through the screen into the tank while continuing gentle agitation.
5. If a flowable formulation is used, premix one part flowable with one part water, and add the diluted mixture SLOWLY through the screen into the tank while continuing gentle agitation.
6. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water, and add the diluted mixture slowly through the screen into the tank while continuing gentle agitation.
7. Continue filling the tank with water through the screen and add the required amount of this product near the end of the filling process.
8. If a nonionic surfactant is used, add it to the tank before completing the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquids (this product) followed by surfactant.

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying.

Keep by-pass and return lines on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

7.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product as indicated in the following table in water.

Spray Solution**Amount of GLYPHOSATE 41% SL**

Desired Volume	0.5%	1%	1.5%	2%	5%	10%
1 gal	0.7 oz.	1.3 oz.	2 oz.	2.7 oz.	6.5 oz	13 oz.
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

2 tablespoons – 1 fluid ounce

For use in backpack sprayers, the appropriate amount of this product be mixed with water in a larger container and then fill the sprayer with the mixed solution.

7.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

7.6 Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates in this label. Lower rates will result in reduced performance.

7.7 Colorants or Dyes

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce performance. Use colorants or dyes according to the manufacturer's recommendations.

7.8 Drift Reduction Additives

Drift reduction additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet Applicator (CDA) equipment. When a drift reduction additive is used, read and carefully observe the precautions, limitations, and all other information appearing on the additive label. Use of drift reduction additives can affect spray coverage, which can reduce product performance.

8.0 APPLICATION EQUIPMENT AND TECHNIQUES Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment: Aerial –Fixed wing and helicopter.

Ground Broadcast Spray--Boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held or Backpack Spray Equipment--Backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment-- Shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems--Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA)---Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING THE DESIRED VOLUMES.

8.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED IN THIS LABEL.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling. Use the rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 2 quart per acre. Refer to the individual use area sections of this label for further instructions.

FOR AERIAL APPLICATION IN ARKANSAS OR CALIFORNIA, OR SPECIFIC COUNTIES THEREIN, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE OR COUNTY FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

This product, when tank mixed with dicamba, may not be applied by air in California.

Ensure uniform application - to avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target movement during aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Comply with all State regulations where applicable.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (read the following sections on "Wind", "Temperature and Humidity", and "Temperature Inversions").

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures rates for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing spray pressure.
- **Number of nozzles:** Use the minimum number of nozzles that will provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind speed, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke produced by a ground source or an aircraft smoke generator can also identify temperature inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water. **Aircraft Maintenance**

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may help prevent corrosion.

8.2 Ground Broadcast Equipment

Apply the product rates in this label at 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the range to ensure complete coverage. Carefully select proper nozzles to avoid generating a fine mist. For best results with ground application equipment, use flat spray nozzles. Check spray pattern for uniform distribution.

8.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled on a spray-to-wet basis; do not spray to the point of run-off. Spray coverage should be uniform and complete. Use coarse sprays only. For rates and timing, refer to the "ANNUAL WEEDS HAND-HELD OR BACKPACK EQUIPMENT" section of this product label.

8.4 Selective Equipment

This product may be diluted in water and applied through shielded sprayers, hooded sprayers, wiper applicators or sponge bars to weeds growing in any non-crop site specified on this label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants), and wipers may be used over-the-top of crops only when specifically in this product's labeling. Such equipment must be capable of preventing all crop contact with the herbicide solutions and operated without leakage of spray mists or dripping onto crop.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of this product with desirable vegetation may result in unwanted plant damage or destruction. Crop injury may occur when the foliage of treated weeds comes into contact with leaves of the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Sprayers

This product, when applied at rates under the conditions described in the following paragraphs for shielded and hooded sprayers, will control those weeds listed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Adjust the shields on these sprayers to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is capable of completely enclosing the spray pattern. If necessary, extend the front and rear flaps of the hooded applicator downward to reach the ground in deep furrows. **EXTREME CARE MUST BE TAKEN TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be configured and operated in a manner that minimizes bouncing and avoids raising the hoods up off the ground at any time. If the hood is raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operating this equipment on rough or sloping terrain where the spray hoods might be raised up off the ground surface.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure low-drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is used. Spray volume should be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- Spray hoods must be operated on the ground or skimming across the ground surface.

- Leave at least an 8-inch untreated strip over the drill row. (For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.)
- Operate at ground speeds of no greater than 5 miles per hour to avoid bouncing of the spray hoods.
- Apply when wind speeds are 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Wiper Applicators

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators: Use solutions ranging from 33 to 75 percent of this product in water. **For**

Panel Applicators: Use solutions ranging from 33 to 100 percent of this product in water.

8.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products for use in injection systems.

8.6 CDA Equipment

The rate of this product applied per acre by vehicle-mounted controlled droplet applicator (CDA) equipment must not be less than the amount used in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply in 2 to 15 gallons of water per acre.

For control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet applicators produce a spray pattern that is not easily visible. Extreme care must be taken to avoid spray or drift onto the foliage or any other green tissue of desirable vegetation, as damage or destruction of the plant may result.

9.0 ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW, SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the "ROUNDUP READY CROPS" section of this label or separately published Repar-Glypho, LLC. Supplemental Labeling for this product for instructions on treating Roundup Ready crops.

TYPES OF APPLICATIONS:

Chemical Fallow, Preplant Fallow Beds, Preplant, At-Planting, Preemergence, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest.

GENERAL USE INSTRUCTIONS:

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", AND WOODY BRUSH AND TREES RATE SECTIONS" in this label. Use rates of this product that are emphasized in this product's labeling to control tough weeds take precedence over the general rates in the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS". Repeat applications may be made up to a maximum of 8 quarts of this product per acre per year.

Post-directed hooded sprayers and wiper applicators capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted in the individual crop sections that follow, wipers may also be used above certain crops to control tall weeds. Refer to the "SELECTIVE EQUIPMENT" section of this label for essential precautions regarding crop injury. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified in this product's labeling. Refer to the "AERIAL EQUIPMENT" section of this label for additional information.

GENERAL PRECAUTIONS, RESTRICTIONS:

Do not allow this herbicide to contact foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of. When making at-planting and preemergence applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death of emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any crops not listed on this label. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction-

For broadcast postemergence treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

9.1 Cereal and Grain Crops

Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Rye, Quinoa, Teff, Teosinte, Triticale, Wheat (all types), Wild rice.

RESTRICTIONS: Do not treat rice fields or levees when flooded.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Red Rice Control Prior to Planting Rice, Spot Treatment (Except Rice), Over-The-Top Wiper Application (Feed Barley and Wheat only), Preharvest (Wheat And Feed Barley Only).

Preplant, Preemergence, At-Planting

Apply before, during or after planting of cereal crops. Applications must be made prior to crop emergence.

TANK MIXTURES: In wheat, a tank mix with Aim may be used.

Red Rice Control Prior to Planting Rice

Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are at the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

RESTRICTIONS: Do not spray during conditions of low humidity, as reduced control may result. Do not treat rice fields or levees when the fields contain floodwater. Do not flood treated fields for 8 days following application.

Spot Treatment (Except Rice)

Apply as a spot treatment in cereal crops. Apply this product before heading in small grains.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Over-the-Top Wiper Application (Feed Barley and Wheat only)

Use Wiper applications in wheat and feed barley. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth.

PRECAUTIONS, RESTRICTIONS: Allow at least 35 days between application and harvest. Do not use roller applicators. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation will result in discoloration, stunting or destruction.

Preharvest (Feed Barley And Wheat Only)

This product provides weed control when applied prior to harvest of feed barley or wheat. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard-dough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest.

For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1 quart of this product per acre. Allow 7 days between application and harvest or grazing. Preharvest application is not used for barley or wheat grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

Apply after harvest of cereal crops. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.2 Corn

TYPES OF CORN: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest.

Use directions for corn containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply alone or in a tank-mixture before, during or after planting corn. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with the following products to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Ensure that the specific product is registered for application prior to planting corn. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

2,4-D	Distinct™	Leadoff™
Aim [®]	Dual Magnum™	Alachlor
Atrazine	Dual II Magnum™	Linex™/Lorox™
Axiom™	Epic™	Keystone LA
Balance™ / Balance PRO	Frontier™/Outlook™	Marksman™
Banvel™/Clarity™	Fultime™	Micro-Tech®
Bicep Magnum™	Guardsman™/Guardsman MAX	Prowl™
Bicep II Magnum™	Harness®	Python™
Bullet [®]	Harness Xtra	Resource®
Degree®	Harness Xtra 5.6 l	Simazine
Degree Xtra®	Hornet	Surpass
	Lariat®	Topnotch™

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

RESTRICTIONS: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. The area covered by this restriction includes Illinois and Indiana south of Route 50, and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

Hooded Sprayers

Apply with hooded sprayers for weed control between rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instructions for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Corn must be at least 12 inches tall, measured without extending leaves. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Do not apply more than 1 quart of this product per acre for each application and no more than 3 quarts per acre per year for hooded sprayer applications.

Spot Treatment

This product may be applied as a spot treatment prior to silking of corn.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

Apply at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the ear is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 2 quarts of this product per acre.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Preharvest application is not used for corn grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required to control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.3 Cotton

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Selective equipment, Spot treatment, Preharvest.

Use directions for cotton containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton.

TANK MIXTURES: This product may be tank-mixed with the following products provided that the specific product being used is registered for application prior to planting cotton. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Caparol®	Direx	Prowl
Clarity™	Dual Magnum™	Prowl H2O
Command	Dual II Magnum™	Staple®
Cotoran®	Karmex	Stalwart
Cotton PRO®	Meturon®	Zorbal®

PRECAUTIONS, RESTRICTIONS: Applications must be made prior to crop emergence. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Selective Equipment

Apply through hooded sprayers, shielded sprayers or wiper applicators in cotton. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Allow at least 7 days between application and harvest. Do not treat more than 10 percent of the total field area to be harvested.

Spot Treatment

Apply as a spot treatment in cotton prior to boll opening.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Do not spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "ANNUAL WEEDS", and "PERENNIAL WEEDS RATE SECTIONS" of this label. For cotton regrowth inhibition, apply 1 pint to 2 quarts of this product per acre.

Up to 2 quarts of this product may be applied after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

TANK MIXTURES: Tank mix with DEFTM 6, Folex™, Ginstar, or Prep™ to provide additional enhancement of cotton leaf drop.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Do not apply preharvest for cotton grown for seed, as a reduction in germination or vigor may occur. THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF THIS PRODUCT TO COTTON IS PROHIBITED. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

9.4 Fallow Systems

Apply during the fallow period prior to planting or emergence of any crop listed on this label; for non-listed crops, applications must be made at least 30 days prior to planting.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Aid-to-Tillage, Chemical Fallow

Use as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Tank mixtures of this product with 2,4-D and dicamba may be used for a broader weed control spectrum. Applications up to 2 quarts per acre may be made by aerial application onto fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

RESTRICTIONS: Do not apply dicamba tank mixtures by air in California. Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

Preplant Fallow Beds

This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label prior to planting.

TANK MIXTURES: Use 12 fluid ounces of this product, plus 2 to 3 fluid ounces of Goal™ 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 3" -- common cheeseweed, chickweed, groundsel; 6" -- London rocket, shepherd's-purse.

Use 16 fluid ounces of this product, plus 2 to 3 fluid ounces of Goal™ 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 6" -- common cheeseweed, groundsel, mare's tail (*Conyza canadensis*), 12" -- chickweed, London rocket, shepherd's-purse.

Aid-to-Tillage

Apply in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 12 fluid ounces of this product in 3 to 10 gallons of water per acre before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

Tank mixtures with residual herbicides may result in reduced product performance.

9.5 Grain Sorghum (Milo)

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Over-the-Top Wiper Application, Preharvest.

Preplant, At-Planting, Preemergence

Apply alone or in tank-mixture before, during or after planting grain sorghum. Applications must be made prior to crop emergence.

TANK MIXTURES: Apply with one of the following pesticide products as a tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, provided that the specific product being used is labeled for application prior to planting grain sorghum.

Atrazine	IntRRo
Bicep II Magnum	Lariat
Bullet	Micro-Tech
Dual II Magnum	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply a tank mixture of this product at 2 pints per acre plus the products listed above. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For other labeled annual weeds listed on this label, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased for acceptable weed control.

Spot Treatment, Over-the-Top Wiper Application

Apply as a spot treatment in grain sorghum before heading. Apply with wiper applicators to control or suppress tall weeds. For additional instructions, see "Wiper Applicators" in the "SELECTIVE EQUIPMENT" section of this label.

RESTRICTIONS: For spot treatment, do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Do not spray or to allow spray to drift outside target area to avoid unwanted crop destruction. Allow at least 40 days between wiper application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

Hooded Sprayers

Apply with hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Milo must be at least 12 inches tall, measured without extending leaves. Treat before milo, sends tillers between the drill rows. If tillers are sprayed, the main plant may be damaged or destroyed. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage, which shall be the sole responsibility of the applicator. Do not graze or feed milo forage or fodder following hooded sprayer applications. Do not apply more than 1 quart of this product per acre per application and no more than 3 quarts per acre per year for hooded sprayer applications.

Preharvest

Apply for weed control prior to harvest after sorghum grain has reached 30 percent grain moisture or less.

RESTRICTIONS: Do not apply more than 2 quarts of this product per acre. As with other herbicides that cause sudden plant death, avoid preharvest applications of this product to milo infected with charcoal rot as lodging can occur. Allow a minimum of 7 days between application and harvest of grain sorghum. Preharvest application is not used for sorghum grown for seed, as a reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California

Post-Harvest

Apply after harvest of grain sorghum. Higher rates may be required to control large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product per acre for control, or 1.5 pints of this product per acre for suppression.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.6 Herbs and Spices

Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Cilantro (leaf), Cilantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Over-the-Top Wiper Application (Peppermint and Spearmint Only), Spot Treatment (Peppermint and Spearmint Only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Over-the-Top Wiper Application, Spot Treatment (Peppermint and Spearmint only)

Apply as a spot treatment or over the top of peppermint or spearmint with wiper applicators in spearmint and peppermint or apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution to a limited area.

RESTRICTIONS: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested should be treated at one time. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. For wiper applications, contact of the herbicide solution with the crop may result in discoloration, stunting, or destruction.

9.7 Oil Seed Crops

Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower.

Use directions for canola containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

Apply before, during or after planting oil seed crops listed in this section. Broadcast applications must be made prior to crop emergence. Wiper applicators or hooded sprayers may be used between the rows once the crop is established. See additional instructions of the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl may be applied before, during or after planting into conventionally tilled soil, a cover crop, established sod or previous crop residue.

RESTRICTIONS: For use with canola, do not apply more than 2 quarts of this product per acre. For use with sunflowers, do not apply more than 1 quart of this product per acre as a single preplant or preemergence application per year. Do not feed or graze sunflower forage following application of this product.

9.8 Soybeans

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest, Selective Equipment.

Use directions for soybeans containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply alone or in a tank-mixture before, during or after planting soybeans. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with one or more of the following products to provide residual weed control, a broader spectrum, or an alternate mode of action. Ensure that the specific product being used in the tank mixture is registered for application prior to planting soybean, and follow all label directions of all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Aim™	Domain™	Micro-Tech
Assure II™	Dual Magnum	Prowl
Authority™	Dual II Magnum	Pursuit™
Axiom	Firstate™	Pursuit Plus
Blanket	Flexstar™	Python
Boundary	Frontier™/Outlook™	Reflex™
Canopy™	Fusion™	Resource™
Canopy EXT™	Gauntlet™	Scepter™
Classic	Gangster	Scor™/Lexone™
Command™	IntRRo	Squadron™
Command Xtra™	Linex™	Steel™
2,4-D	Lorox/Linuron	Valor™
	Lorox Plus™	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when

weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Spot Treatment

Apply as a spot treatment prior to initial pod set in soybeans.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction.

Preharvest

Apply for weed control prior to harvest of soybeans after pods have set and lost all green color. Apply at rates given in the "ANNUAL WEEDSRATE SECTION" and "PERENNIAL WEEDSRATE SECTION". Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Do not apply more than 5 quarts of this product per acre for preharvest applications. Do not apply more than 2 quarts of this product per acre by air. Allow a minimum of 7 days between application and harvest of soybeans. Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last preharvest application. (If the application rate is 1 quart per acre or lower, the grazing restriction is reduced to 14 days after last preharvest application.) Preharvest application is not used for soybeans grown for seed, as a reduction in germination or vigor may occur.

Selective Equipment

Apply through shielded or hooded sprayers, wiper applicators or sponge bars in soybeans. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow at least 7 days between application and harvest.

9.9 Sugarcane

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus Spot Treatment and Sugarcane Ripening.

Preplant, At-Planting, Preemergence

Apply in or around sugarcane fields, or in fields prior to the emergence of plant cane.

RESTRICTIONS: Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Spot Treatment

Apply as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using spray-to-wet a spray-to-wet technique. Volunteer or diseased sugarcane should have at least 7 new leaves.

RESTRICTIONS: Do not allow spray contact with healthy sugarcane plants because severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

Fallow Treatments

Apply as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Applications up to 3 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used.

Hooded Sprayers

Apply using hooded sprayers for weed control between the rows of sugarcane. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional use instructions.

RESTRICTIONS: Crop injury may occur when the foliage of treated weeds contact the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction.

Sugarcane Ripening

Apply to foliage as a plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane. It is effective in both low and high-tonnage sugarcane.

When applied as directed under the conditions described, this product will hasten ripening and extend the period of high sucrose level in sugarcane.

As a result of leaf desiccation, improved trash bum can be expected.

Most of the sucrose increase is concentrated in the top nodes of the treated sugarcane stalk. In order to maximize sugar where topping is practiced during harvest, top at the base of the fourth leaf.

APPLICATION RATES:

NOTE: Use the higher rate within the used range when treating sugarcane under adverse ripening conditions or when less responsive varieties are to be treated.

Use the following application rates and timing instructions according to the State in which the sugarcane is grown.

FLORIDA—Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of LAST PLATOON CANE ONLY.

HAWAII—Apply 12 to 28 fluid ounces of this product per acre 4 to 10 weeks before harvest.

LOUISIANA—Apply 5 to 16 fluid ounces of this product per acre 3 to 7 weeks before harvest of PLATOON CANE ONLY.

PUERTO RICO—Apply 7 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

TEXAS—Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

PRECAUTIONS, RESTRICTIONS: Prior to application, consult your state sugarcane authority or local Repar-Glypho, LLC representative regarding the degree of sucrose response anticipated from the variety of sugarcane to be treated.

Application of this product may initiate development of shooting eyes. This product may not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2 to 3 weeks after application, this product may produce a slight yellowing to pronounce browning and drying of leaves, and a shortening of upper internodes. Spindle death may occur.

Rainfall within 6 hours after application may reduce effectiveness.

Application is not used for sugarcane grown for seed, as a reduction in germination or vigor may occur.

Do not feed or graze treated sugarcane forage following application. Do not plant subsequent crops in treated fields other than the following for 30 days after application: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybeans, squash (all types) or wheat.

9.10 Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL VEGETABLE CROPS LISTED ALPHABETICALLY IN THIS SECTIONS THAT FOLLOW. **SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.**

TYPES OF APPLICATIONS: Chemical Fallow, Preplan Fallow Beds, Preplan, At-Planting, Preemergence, Prior to Transplanting Vegetables, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest, Selective Equipment Applications (Nonbearing Ginseng), Over-the-Top Wiper Application (Rutabagas only), Spot Treatment or Preharvest (Dry Beans, Peas, Lentils, and Chickpeas only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

9.10.1 Brassica Vegetables

Broccoli, Chinese broccoli (gai lan), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

9.10.2 Bulb Vegetables

Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

9.10.3 Cucurbit Vegetables and Fruits

Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechimma, Chinese okra), Melons (all), *Momordica spp* (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon.

RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

9.10.4 Leafy Vegetables

Amaranth (Chinese spinach), Arugula Croquette), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Com salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

RESTRICTIONS: For Watercress, do not within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of crop injury.

9.10.5 Fruiting Vegetables

Eggplant, Groundcherry (*Physalis spp.*), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.

RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato and tomatillos grown on sandy soil, do not make hooded, shielded sprayer or wiper applications in row middles because of potential for crop injury.

9.10.6 Legume Vegetables (Succulent or Dried)

Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

Spot Treatment, or Preharvest (Dry Beans, Peas, Lentils and Chickpeas only)

Apply as an over-the-top broadcast spray or as a spot treatment to control labeled weeds in dry beans, peas, lentils or chick peas. For spot treatment, to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed, apply in 10 to 20 gallons of water per acre through ground broadcast spray equipment or use a 2 percent solution in a hand-held sprayer. For preharvest treatments, apply in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less).

PRECAUTIONS, RESTRICTIONS: Follow the limitations listed in the table below. Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not specified for treatment in this label. Do not feed treated vines and hay from these crops to livestock. Do not treat cowpeas or field (feed) peas, since these crop are considered to be grown as livestock feed. Preharvest application is not used for dry legumes grown for seed, as a reduction in germination or vigor may occur.

Crop	Maximum Rate (fluid ounces per Acre)	Minimum Pre- Harvest Interval (days)	Allowed in
Dry Beans	32	7	All States
Dry Peas, Lentils, and Chickpeas	26	14	Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, and Washington

9.10.7 Root and Tuber Vegetables

Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

Selective Equipment Applications (Non-Bearing Ginseng only)

Apply for general weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and backpack wands, lances, and orchard guns or with wiper application equipment. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Conduct applications so that there is no contact of this product with the ginseng plant. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Applications must be made at least one year prior to harvest.

Over-the-Top Wiper Application (Rutabagas only)

Use Wiper applicators over the top of rutabagas for the control of taller weeds. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

9.11 Miscellaneous Crops

Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: General Weed Control, Site Preparation, Spot Treatment (Asparagus).

Use directions for sugar beets containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

RESTRICTIONS: Avoid contact of this product with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to

further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any crop not listed on this label. See additional information in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

General Weed Control, Site Preparation

This product may be applied for general weed control or for site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove residues of this product from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Allow at least 21 days between residue removal and transplanting. Applications made at emergence will result in injury or death to emerged seedlings.

Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.

Spot Treatment (Asparagus)

Apply immediately after cutting, but prior to the emergence of new spears.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Post-Harvest (Asparagus)

Apply after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Direct contact of the spray with asparagus may result in serious crop injury.

10.0 TREE, VINE, AND SHRUB CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL TREE, VINE, AND SHRUB CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Preplant (site preparation) Broadcast Sprays, General Weed Control, Middles (between rows of trees, vines or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (shielded sprayers, wiper treatments), Directed Spray, Spot Treatment, Perennial Grass Suppression, Cut Stump.

This product may be applied with boom equipment, CDA equipment, shielded sprayers, hand-held and backpack wands, lances, orchard guns or with wiper applicator equipment, unless prohibited in the specific Crop sections that follow.

Apply in middles (between rows of trees or vines), strips (within rows of trees or vines), and for general weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 1 pint to 5 quarts per acre according to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label. Use the higher rates in the range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts per acre per year. See the "GENERAL INFORMATION" section of this label for more information on Maximum Annual Use Rates.

GENERAL PRECAUTIONS, RESTRICTIONS: Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used in order to minimize the potential for overspray or drift of this product onto the crop. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with the crop may be used. See additional instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

Allow a minimum of 3 days between application and transplanting.

Middles (between rows)

This product will control or suppress annual and perennial weeds and ground covers growing between rows of tree and vine crops listed on this label. If weeds are under drought stress, irrigate prior to application. Reduced weed control may result if weeds have been recently mowed at the time of application.

TANK MIXTURES: A tank mixture of this product plus Goal 2XL may be applied for annual weed control between rows (middles) of citrus crops, tree fruits, tree nuts and vine crops. This mixture is used when weeds are stressed or growing in dense populations. Application of 16 to 32 fluid ounces of this product per acre plus 3 to 12 fluid ounces of Goal 2XL per acre will control annual weeds with a maximum height, length or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). This tank mixture will also control common cheeseweed (malva) or hairy fleabane (*Conyza bonariensis*) with a maximum height, length or diameter of 3 inches. Read and follow all labels and directions for use of all products being used in the tank mixture.

Strips (In Rows)

TANK MIXTURES: This product may be applied within rows of tree or vine crops in tank mixtures with the following products, provided that the label of the specific product used permits the desired use:

Devrinol™ 50 DF	Simazine 4L
Direx™ 4L	Simazine 80W
Goal 2XL	Sim-Trol™ 4L
Karmex DF	Solican™ DF
Krovarl	Surflan™ AS
Prowl	Surflan 75w
Princep Caliber™90	

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for information about specific crops, rates, geographic restrictions and precautionary statements. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Perennial Grass Suppression

This product will suppress perennial grasses such as bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

For burndown of Bermudagrass, apply 1 to 2 quarts of this product in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 6 to 16 fluid ounces of this product per acre east of the Rocky Mountains and 16 fluid ounces of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the Bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces of this product per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

Cut Stump (tree crops)

Make cut stump applications of this product during site preparation or site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

Citrus Trees: Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo (Ugli), Tangor.

Fruit Trees: Apple, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince.

Nut Trees: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).

Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

PRECAUTIONS, RESTRICTIONS: DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

10.1 Berry Crops

Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Salal. [Optional Crop: Raspberry (black, red)].

TYPES OF APPLICATIONS: Those listed in Section 10.0, plus Spot Treatment in Cranberry Production and Post-Harvest Treatment in Cranberry Production.

RESTRICTIONS: To avoid damage, herbicide spray must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest of cranberries. Allow a minimum of 14 days between last application and harvest of other berry crops.

Spot Treatment in Cranberry Production

Use spot treatments to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in this label may be used. Drop water level to remove standing water in ditches prior to application. With hand-held sprayers, use 1- to 2-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off.

RESTRICTIONS: To achieve maximum weed control in treatments of dry ditches after water draw down, allow 2 or more days after treatment before reintroduction of water. Apply this product within 1 day after draw down to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not make applications by air. Do not apply directly to water. Use nozzles that emit medium- to large-sized droplets in order to minimize spray drift and avoid crop injury.

Post-Harvest Treatments in Cranberry Production

Apply this product may be made after the harvest of cranberries to control weeds growing in the field. Apply this product to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label may be used. If using hand-held sprayers, use a 0.5- to 1-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off. If using hand-held boom sprayers, apply 2 to 4 quarts of this product per acre.

RESTRICTIONS: Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not make applications by air. Do not apply directly to water. Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

10.2 Citrus

Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugh), Tangor.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only.

The following use instructions pertain to applications in Florida and Texas only: For burndown or control of the weeds listed below, apply this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 2 to 3 quarts of this product in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall, and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, use of this product in a tank mixture with KrovarTM or KarmexTM may improve weed control. Refer to the individual product labels for treatment information on specific crops, rates, geographic restrictions and precautionary statements.

Perennial weeds:

S = Suppression

B = Burndown

PC = Partial control

C = Control

WEED SPECIES	GLYPHOSATE 41% SL RATE PER ACRE			
	1 QT	2 QT	3 QT	5 QT
Bennudagrass	B	---	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	---	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	---	PC	C

10.3 Miscellaneous Tree Food Crops

Cactus (fruit and pads), Palm (heart, leaves), Palm (oil).

TYPES OF APPLICATIONS: Those listed in Section 10.0.

10.4 Non-Food Tree Crops

Pine, Poplar, Eucalyptus, Christmas trees.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

NOTE: unless otherwise directed, this product is not used for use as an over-the-top broadcast spray in plantations or other labeled tree crops.

Directed Spray, Spot treatment, Wiper Application

This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

PRECAUTIONS, RESTRICTIONS: Avoid contact of spray, drift or mist of this product with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Site preparation

Apply this product for weed control prior to planting non-food tree crops.

PRECAUTIONS: Precautions should be taken to protect non-target plants during site preparation applications.

10.5 Pome Fruit

Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince. TYPES OF

APPLICATIONS: Those listed in Section 10.0.

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in pome crops.

10.6 Stone Fruit

Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), Plumcot.

TYPES OF APPLICATIONS: Those listed in Section 10.0. For olive groves, apply only as a directed spray.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 17 days between last application and harvest in stone fruit crops.

Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. USE EXTREME CARE TO ENSURE THAT NO PART OF THE PEACH TREE IS CONTACTED WITH OVERSPRAY OR DRIFT OF THIS PRODUCT.

10.7 Tree Nuts

Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (black, English).

TYPES OF APPLICATIONS: Those listed in Section 10.0.

RESTRICTIONS: Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut.

10.8 Tropical and Subtropical Trees and Fruits

Ambarella, Atemoya, Avocado, Banana, Barbados cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates,

Durian, Feijoa, Figs, Governor's plum, Guava, Hama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genie), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots and leaves), Wax jambe.

TYPES OF APPLICATIONS: Those listed in Section 10.0, and as described below as a Bananacide (Bananas Only).

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit listed. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

Bananacide (Banana Only)

Apply to destroy banana plants infected with the Banana Bunchy Top Virus, as well as non-infected banana plants in order to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1/25 fluid ounce (1 mL) of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) within a 4-foot radius around a treated mat should be mechanically destroyed.

For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1/2 fluid ounce (15 mL) of this product per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for general weed control.

10.9 Vine Crops

Grapes (raisin, table, wine) Hops, Kiwi, Passion fruit.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

Apply this product for weed control only when green shoots, canes or foliage are not in the spray zone.

In the northeast and Great Lakes regions, apply this product to grape vineyards prior to the end of the bloom stage of grapes in order to avoid crop injury, or apply using shielded sprayers or wiper equipment.

RESTRICTIONS: Allow a minimum of 14 days between last application and harvest of vine crops. Do not use selective equipment in kiwi.

11.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELANDS

GENERAL USE INSTRUCTIONS: Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label for rates for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. Application rates of this product specified on this label for the control of tough weeds, or those specified on separate supplemental labeling for this product, supersede rates in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling.

11.1 Alfalfa, Clover, and Other Forage Legumes

Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types).

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Preharvest (except Kenaf and Leucaena), Spot Treatment, Over-the-Top Wiper Application, Renovation, Stand Removal.

Use directions for alfalfa with a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply before, during or after planting crops listed in this section. Applications must be made prior to crop emergence.

RESTRICTIONS: Remove domestic livestock before application. Do not feed or graze until crop has reached sufficient maturity.

Preharvest (Except Kenaf and Leucaena)

This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This product will control annual or perennial weeds, including quackgrass, when applied prior to crop harvest. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

PRECAUTIONS, RESTRICTIONS: Make only one application to an existing stand of crop per year. The treated crop and weeds can be harvested and fed to livestock according to the application rates and intervals defined below.

	Maximum single application rate (per acre)	Minimum interval between application and harvest or livestock grazing
Alfalfa	2 quarts	36 hours
All other legumes listed	3 pints	3 days

This application may destroy the alfalfa stand and may severely injure or destroy other labeled crops such as clover. Do not apply for preharvest application for alfalfa grown for seed, as a reduction in germination or vigor may result.

Spot Treatment, Over-the-Top Wiper Application

Apply as a spot treatment or over the top of crops listed in this section with wiper applicators. For wipers, see "Wiper Applicators" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field area should be treated at one time. Remove domestic livestock before application and wait 3 days after application before grazing livestock or harvesting.

Renovation, Stand Removal

Apply as a broadcast spray to remove established stands of alfalfa, clover, and other forage legumes listed on this label. If the crop is to be grazed or harvested for feed, use up to a maximum of 2 quarts per acre in alfalfa and up to 3 pints per acre in other labeled legumes. For complete removal of established stands of clover, it may be necessary to use treatment rates greater than this, as listed in the "PERENNIAL WEEDS RATE SECTION" of this label.

PRECAUTIONS, RESTRICTIONS: For applications up to 2 quarts per acre for alfalfa or 3 pints per acre for all other forage legumes, remove domestic livestock before application, follow the minimum harvest or grazing intervals in the table above under Preharvest in this section. For treatment rates above these levels, do not graze or harvest treated foliage for livestock feed or allow grazing. Crops listed in this label may be planted into the treated area at any time; for all other crops, wait 30 days or more between application and planting.

11.2 Conservation Reserve Program (CRP)

TYPES OF APPLICATIONS: Renovation (Rotating Out Of CRP), Site Preparation, Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application.

Renovation (Rotating Out of CRP), Site preparation

This product may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation recommendations.

RESTRICTIONS. Crops listed on this label may be planted into the treated area at any time; for any other crop, wait at least 30 days between application and planting.

Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application

Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land. Applications may be made using wiper applicators to control tall weeds or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: *Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant.* No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

11.3 Grass Seed or Sod Production

Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation, Shielded Sprayers, Over-the-Top Wiper Application, Spot Treatment, Creating Rows in Annual Ryegrass.

Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation

This product controls most existing vegetation prior to renovating turf or forage grass seed areas or establishing turf grass grown for sod. It may be used to destroy undesirable grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting or for renovation. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turf management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates of up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed for treatment in this label, applications must be made at least 30 days prior to planting. Applications must be made prior to the crop emergence in order to avoid crop injury.

Shielded Sprayers

Apply 1 to 3 quarts of this product as a broadcast spray in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields. See additional instructions on the use of shielded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. .

Over-the-Top Wiper Application

Apply this product for over the top of desirable grasses using wiper applicators for the control of tall weeds. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. .

Spot Treatment

Apply a 1- to 1.5-percent solution of this product using hand-held spray equipment to control *weeds* within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may also be used to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS, RESTRICTIONS: Apply this product prior to heading of grasses. Do not treat more than 10 percent of the total field area. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. Hand-held equipment maybe used to control sod remnants or other unwanted vegetation after sod is harvested.

Creating Rows in Annual Ryegrass

Apply 16 to 32 fluid ounces of this product per acre. Best results are obtained when applications are made before ryegrass reaches 6 inches in height. Use the higher rate when the ryegrass is greater than 6 inches in height. Set nozzle heights to allow the establishment of the desired row spacing. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are used.

PRECAUTIONS.: Take care not to spray or allow droplets, spray fines, or drift to settle outside the treatment area to avoid unwanted crop destruction.

Grower assumes all responsibility for crop losses from misapplication.

11.4 Pastures

Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label, including Bahiagrass, Bermudagrass, Bluegrass, *Bromegrass*, *Fescue*, *Guineagrass*, *Kikuyu grass*, *Orchardgrass*, Pangola grass, Ryegrass, Timothy, Wheatgrass.

TYPES OF APPLICATIONS: Preplant, Preemergence, Pasture Renovation, Spot Treatment, Over-the-Top Wiper Application, Postemergence Weed Control (Broadcast Treatments).

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be used to control perennial pasture species listed on this label prior to re-planting.

RESTRICTIONS: If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed on this label, applications must be made at least 30 days prior to planting.

Spot Treatment, Over-the-Top Wiper Application

This product may be applied in pastures as a spot treatment or over the top of desired grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

RESTRICTIONS: For spot treatment and wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acre, no more than 10 percent of the field may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing or harvesting for feed.

Postemergence Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto pasture grasses except for renovation use described above in this section. If replanting is needed due to severe stand reduction, wait at least 30 days after treatment prior to planting any crop not listed on this label.

11.5 Rangelands

TYPES OF APPLICATIONS: Postemergence.

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands. Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Apply 12 to 16 fluid ounces of this product to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are used where spring moisture is usually limited and fall germination allows for good weed growth.

For medusahead, apply 16 fluid ounces of this product at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

PRECAUTIONS, RESTRICTIONS: Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off. Do not use ammonium sulfate when spraying rangeland grasses with this product. No waiting period between treatment and feeding or livestock grazing is required. Do not apply more than 3 quarts of this product per acre per year.

12.0 ROUNDUP READY CROPS

The following instructions or those published separately on Repar-Glypho, LLC. supplemental labeling for this product include all applications of this product that can be made onto the specified Roundup Ready crops during the complete cropping season. DO NOT combine these treatment instructions with those for crops in the "ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)" section of this label that do not contain a glyphosate tolerant gene, unless otherwise directed in this product's labeling.

USE THIS PRODUCT FOR POSTEMERGENCE (IN-CROP) APPLICATION ONLY ON CROPS DESIGNATED AS CONTAINING A GLYPHOSATE TOLERANT GENE. Applying this product to crops that are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a glyphosate tolerant gene, since severe plant injury or destruction will result. The Roundup Ready designation indicates that the crop contains a patented gene that provides tolerance to this product. Information on Roundup Ready crops may be obtained from your seed supplier or Repar-Glypho, LLC. representative. Roundup Ready crops must be purchased from an authorized licensed seed supplier.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground broadcast applications. Apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications. All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling, in particular in the "Aerial Equipment" section. Apply this product in 3 to 15 gallons of water per acre. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

For proper stewardship of aerial applications over the top of Roundup Ready crops, read and follow all precautions and procedures contained in the use guide "A Guide to On-Target Aerial Application." Copies of this publication are available by calling 1-800-ROUNDUP (1-800-768-6387).

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE.

TANK MIXTURE: Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT for applications of this product over the top of Roundup Ready crops unless otherwise noted in this product label.

Unless otherwise directed, nonionic surfactant may be added to the spray solution for applications to Roundup Ready crops. The addition of certain surfactants to this product may result in some crop response including leaf necrosis, leaf chlorosis or leaf speckling due to the surfactant added to the spray mixture. Read and carefully observe cautionary statements and other information appearing on the surfactant label.

applied at or before the 4-trifoliolate growth stage. Refer to the following tables for maximum in-crop application rates during stand establishment (seeding year).

STAND ESTABLISHMENT (Seeding Year)	
Application Rates	
Prior to First Cutting	
From emergence up to 4 trifoliolate leaves	1 to 2 quarts per acre
From 5 trifoliolate leaves up to 5 days before first cutting	Up to 2 quarts per acre
After First Cutting	
In-crop application, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

Established Stands—Refer to the following table for directions and maximum application rates for in-crop applications to established stands of alfalfa (non-seeding year).

ESTABLISHED STANDS	
Application Rates	
In-crop applications, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

PRECAUTIONS, RESTRICTIONS: Where Roundup Ready alfalfa is grown with a companion or cover crop, or is overseeded with a second species, in-crop (over-the-top) applications of this product will eliminate the non-glyphosate tolerant species. Any single in-crop application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart. The combined total per year for all in-crop applications in both newly established (seeding year) and established stands (non-seeding year) must not exceed 6 quarts per acre. Remove domestic livestock before application. Wait a minimum of 5 days after application before grazing, cutting and feeding of forage and hay.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application.

Newly Seeded Stands and Stand Establishment: For control of emerged annual grasses and broadleaf weeds, this product may be applied at up to 2 quarts per acre in a tank mixture with the following herbicides. Application should be made after weeds have emerged but before the alfalfa growth or regrowth would interfere with spray coverage of the target weeds.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, bromoxynil, clethodim, imazamox, imazethapyr, sethoxydim.

Arrow, Buctril, Butoxone, Butyrac, Poast, Pursuit, Raptor, Select.

Buctril can only be used in newly seeded stands]

Dormant Application and Winter Treatment: For control of emerged annual grasses and broadleaf weeds, up to 2 quarts per acre of this product may be applied in a tank mixture with the following herbicides. Tank mixtures should be applied when the temperature for the day remains above freezing.

and carefully observe cautionary statements and other information appearing on the surfactant label.

Ammonium sulfate may be mixed with this product for applications to Roundup Ready crops. Refer to the "MIXING" section for instructions on the use of ammonium sulfate. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The following instructions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of this product is used to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed with multiple germination times, or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Rates of this product specified in the following sections, or in separate supplemental labeling published by Repar-Glypho, LLC, for this product on the control of tough weeds, supersede the general rate recommendations in the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" of this label.

12.1 Roundup Ready Alfalfa

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: Refer to the following table for the maximum application rates of this product.

Annual Maximum Application Rates	
Combined total per year for all applications, including preplant during year of establishment	8 quarts per acre
Combined total per year for in-crop applications for newly established and established stands	6 quarts per acre

GENERAL PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting alfalfa with a glyphosate tolerant gene, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

Apply this product may be made over the top of Roundup Ready alfalfa (in-crop) from emergence to 5 days prior to harvest. To maximize crop yield and quality potential, applications of this product should be made after weeds have emerged but before alfalfa growth or re-growth interferes with spray coverage of the target weeds.

Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" in this label for rates for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. In addition to those weeds listed in these sections, this product will suppress the parasitic weed Dodder (*Cuscuta spp*) in Roundup Ready alfalfa. Repeat applications may be necessary for complete control.

Stand Establishment—Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings may not contain a Roundup Ready gene and will not survive after the first application of this product. To eliminate the

Dormant Application

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: diuron, hexazinone, imazamox, metribuzin, pronamide, terbacil, Kerb 50-W, Lexone, Raptor, Sencor, Sinbar, Velpar AlfaMax.]

Winter Treatment

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, diuron, hexazinone, Butoxone, Diruon, Velpar AlfaMax.]

PRECAUTIONS, RESTRICTIONS: Tank mixtures of this product with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control. Applications must not be made to frozen or snow covered ground.

12.2 Roundup Ready Canola (Spring)

Roundup Ready spring canola is defined as those Roundup Ready canola varieties that are seeded in the spring and harvested in the fall and do not enter a winter dormancy period.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

Annual Maximum Application Rates

Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	1 quart per acre

GENERAL PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates."

Preplant, At-planting, Preemergence,

This product may be applied before, during or after planting, up to a maximum of 2 quarts per acre. Postemergence (In-crop)

This product may be applied postemergence to Roundup Ready spring canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds. .

Single Application - Apply 16 to 24 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar crop injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application - Apply 16 fluid ounces per acre to 1 - to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Apply sequential applications for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass, or when multiple application times are needed for adequate weed control.

RESTRICTIONS: No more than two in-crop broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 32 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

12.3 Roundup Ready Canola (Winter)

Roundup Ready Winter Canola is defined as those Roundup Ready canola varieties that are seeded in early fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.

DO NOT USE THIS PRODUCT ON WINTER CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA,

Annual Maximum Application Rates	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	1 quart per acre

KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rate."

Preplant, At-Planting, Preemergence

Apply before, during or after planting, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

This product may be applied to Roundup Ready winter canola varieties from emergence to canopy closure in the fall and prior to bolting in the spring. Applications made during or after bolting may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Some weeds with multiple germination times, or suppressed (stunted) weeds, or weeds that have overwintered may require sequential applications of this product for control. The second application should be made after some regrowth has occurred and at least 60 days after a previous application of this product.

Single Application—Apply 24 to 32 fluid ounces of this product per acre in the fall. Applications in the fall should be made when weeds are small and actively growing. Use the higher rate in the range when weed densities are high, when weeds have overwintered or when weeds become large and well established. Applications of greater than 16 fluid ounces per acre prior to the 6-leaf stage may result in reduced crop growth in the fall. Avoid spray overlaps. Spray overlaps may result in temporary yellowing and/or growth reduction.

Sequential Applications - Apply 16 to 32 fluid ounces of this product per acre to 2-leaf or larger canola in the fall, followed by a sequential application at the same rate and at a minimum interval of 60 days, but before bolting in the spring. Sequential applications are for early emerging annual weeds and winter emerging weeds such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product will control or suppress most perennial weeds. For some perennial weeds, sequential applications may be required to reduce competition with the crop.

RESTRICTIONS: No more than two over-the-top broadcast applications may be made from crop emergence up to the onset of bolting, and the total in-crop application must not exceed 64 fluid ounces of this product per acre. Applications of greater than 24 fluid ounces per acre prior to the 6-leaf stage

may result in reduced crop growth in the fall. Allow a minimum of 60 days between last application and harvest of canola grain. No waiting period is required between application and open grazing of livestock.

12.4 Roundup Ready Corn 2

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Spot treatment, Preharvest, Post-harvest.

Annual Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from emergence through 48 inches	3 quarts per acre
Maximum preharvest application (See PRECAUTIONS, RESTRICTIONS section for Preharvest)	1 quart per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-planting, Preemergence

Apply alone or in a tank-mixture before, during or after planting.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application. Read and follow label directions for all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to corn: 2,4-D, acetochlor, alachlor, atrazine, carfentrazone-ethyl, clopyralid, dicamba, diflufenzopyr, dimethenamid, dimethenamid-p, flufenacet, flumetsulam, flumiclorac pentyl ester, isoxaflutole, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, rimsulfuron.

Aim, Axiom, Balance, Balance PRO, Banvel, Bicep MAGNUM, Bicep II MAGNUM, Bicep Lite II MAGNUM, Bullet, Cinch, Cinch ATZ, Clarity, Define, Degree, Degree Xtra, Distinct, Dual II MAGNUM, Epic, Frontier, FulTime, Guardsman MAX, Harness, Harness Xtra, Harness Xtra 5.6L, Homer, Keystone, Keystone LA, Lariat, Linex, Lorox, Marksman, Me-Too-Lachlor II, Micro-Tech, Prowl, Python, Python II, Radius, Resolve, Resource, Stalwart, Stalwart Xtra, Surpass, TopNotch]

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

Apply alone or in tank mixtures postemergence (in-crop) to corn hybrids designated as Roundup Ready Corn 2. When applied as directed, this product controls annual grass and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. A postemergence application of this product should be made before the weeds reach a height of 4 inches or before they become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre should be made before the weeds reach a height of 4 inches.

This product may be broadcast over-the-top or applied with drop nozzles to Roundup Ready Corn 2 from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing).

whichever comes first. Drop nozzles are for optimum spray coverage and weed control when corn height is 24 to 30 inches. For corn heights 30 to 48 inches (free standing), apply this product only by ground application using drop nozzles aligned to avoid spraying into the whorls of the corn plants.

TANK MIXTURES: This product may be tank-mixed with the following products. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to corn. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to corn: atrazine, acetochlor, alachlor, clopyralid, dicamba, diflufenzopyr, flumetsulam, flumiclorac pentyl ester, foramsulfuron, halosulfuron-methyl, iodosulfuron-methyl-sodium, mesotrione, nicosulfuron, rimsulfuron, thifensulfuron methyl.

Banvel, Basis, Basis Gold, Bullet, Callisto, Clarity, Degree, Degree Xtra, Distinct, Equip, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Marksman, Micro-Tech, Option, Resolve, Resource, YUKON]

Tank Mix Partner	Maximum Height Of Corn For Application
Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6L	11 inches
Bullet* Micro-Tech*	5 inches
Atrazine	12 inches

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet* or Micro-Tech* herbicides.

PRECAUTIONS, RESTRICTIONS: Single in-crop applications of this product should not exceed 3 pints per acre. Sequential in-crop applications of this product from emergence through 48 inches in height must not exceed a total of 3 quarts per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product in-crop and harvest of corn forage or grain. (For applications at preharvest timing, see **Preharvest** section of this label). The use of additional surfactants and other additives, including fertilizers and micro-nutrients, in the spray solution are not used with this product and may result in increased potential for crop injury or reduced yield, and is not used for over-the-top applications, unless otherwise directed in supplemental labeling of fact sheets published separately by Repar-Glypho, LLC..

Preharvest

A single preharvest application of up to 32 fluid ounces per acre of this product may be applied for annual and perennial weed control prior to crop harvest. Make application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

RESTRICTIONS: Do not make a preharvest application of this product if more than a combined total of 64 fluid ounces of this product has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

Postharvest

Apply for weed control after corn harvest. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

12.5 Roundup Ready Cotton

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Selective Equipment (In-Crop), Preharvest.

<u>Annual Maximum Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to layby	4 quarts per acre
Maximum preharvest application rate	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." **ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.**

Preplant, At-Planting, Preemergence

Apply before, during or after planting. Tank mixtures with other herbicides listed in Section 8.3 of this label maybe used.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to the emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyriithiobac-sodium.

Caparol, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H20, Stalwart, Staple]

RESTRICTIONS: The maximum quantity of this product that maybe applied for all preplant, at-planting and preemergence applications combined is 5 quarts per acre per season.

Postemergence (in-crop)

Apply over the top Roundup Ready cotton at rates up to 1 quart per acre per application from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). **Applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss.**

TANK MIXTURES: This product may be tank-mixed with the following products and applied over the top of Roundup Ready cotton: Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clethodim, fluazifop-P-butyl, metolachlor, s-metolachlor, pyriithiobac-sodium, sethoxydim, quizalofop-P-ethyl, Assure II, Dual MAGNUM, Fusilade, Poast Plus, Select, Stalwart, Staple
Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop).
Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting to exposed cotton leaves.]

RESTRICTIONS: The maximum quantity of this product that maybe applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. No more than two over-the-top broadcast applications may be made from crop emergence through the 4-leaf (node) stage of development. No more than two applications should be made from the 5-leaf stage through layby. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications. Do not add additional surfactant or additives containing surfactant to this product (other than those contained in any tank-mix product) for over-the-top applications to roundup ready cotton.

Salvage Treatment. Apply after the 4-leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

Selective Equipment (In-crop)

Apply this product using precision post-directed or hooded sprayers at rates up to 1 quart per acre per application to Roundup Ready cotton through layby. At this stage, use post-directed equipment to direct the spray to the base of the cotton plants. Avoid contact of the herbicide spray with the cotton leaves to the maximum extent possible. To minimize spray contact maintain low spray pressure (less than 30 pounds per square inch) and place nozzles in a low position, directing a horizontal spray pattern under the cotton leaves and onto weeds in the row. For best results, make applications while weeds are small (less than 3 inches). See additional instructions on the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: This product can be tank-mixed with the following products for in-crop application using precision post-directed or hooded sprayers. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: carfentrazoneethyl, diuron, flunioxazin, fluometuron, linuron, pendimethalin, prometryn, pyriithiobac-sodium, Aim, Caparol, Chateau, Cotoran, Direx, Layby-Pro, Prowl H20, Staple, Valor
Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop)]

RESTRICTIONS: The maximum quantity of this product that may be applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

Preharvest

Apply for annual and perennial weed control as a broadcast treatment prior to crop harvest after 20 percent boll crack. Apply up to 2 quarts of this product per acre. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Preharvest application is not used for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION; Use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.6 Roundup Ready Flex Cotton

The use of the over-the-top applications described in this section on **other** than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (node) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this section are specific to, and should only be used with, varieties designated as **Roundup Ready Flex cotton**. Do Not combine the instructions in this section, with those in the "Roundup Ready Cotton" section (12.5) of this label, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instructions on labeling for this or other glyphosate-containing product.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Preharvest.

Annual Maximum Allowable Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to 60 percent open bolls	6 quarts per acre
Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyriithiobac-sodium.

Caparel, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H2O, Stalwart, Staple]

Postemergence (In-crop)

Apply to control annual grasses and broadleaf weeds listed on this label in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds is used. This product may be applied postemergence at rates up to 1.5 quarts per acre per application. In addition to broadcast applications, post-directed spray equipment may be used to achieve weed coverage.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready Flex cotton. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clethodim, fluazifop-Pbutyl, metolachlor, s-metolachlor, pyriithiobac-sodium, quizalofop-p-ethyl, sethoxydim, trifloxysulfuron-sodium.

Assure II, Dual MAGNUM, Envoke, Fusilade, Poast Plus, Select, Stalwart, Staple

Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop).

Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting to exposed cotton leaves.)

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 1.5 quarts per acre made using ground application equipment. In-crop application rates above 1 quart per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling or leaf necrosis. Do not exceed a maximum rate of 1 quart per acre of this product *when* making applications by air. Between layby and 60 percent open bolls, the maximum combined total rate of this product that may be applied is 2 quarts per acre. The maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre. Do not add additional surfactant or additives containing surfactant to this product for over-the-top applications to Roundup Ready Flex cotton.

Preharvest

Apply up to 2 quarts of this product may be applied for annual and perennial weed control as a broadcast treatment prior to harvest after 60 percent boll crack. **NOTE:** This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Do not apply preharvest for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION: use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Flex Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.7 Roundup Ready Soybeans

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Preharvest, Post-Harvest.

Annual Maximum Application Rates

Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from cracking through flowering (R2 stage soybeans)	3 quarts per acre
Maximum preharvest application rate	1 quart per acre

GENERAL PRECAUTIONS/RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates."

Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

This product may be applied before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D, Banvel or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to soybeans: alachlor, atrazine, r-arfentrazone-ethyl, chlorimuron ethyl, clethodim, clomazone, cloransulam-methyl, dimethenamid, dimethenamid-p, fenoxypyr, fluazifop-p-butyl, flufenacet, flumetsulam, flumiclorac pentyl ester, flumioxazin, fomesafen, imazaquin, imazethapyr, lactofen, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, sulfentrazone, tribenuron methyl, quizalofop P-ethyl.

Aim, Amplify, Assure II, Axiom, Authority, Blanket, Boundary, Canopy, Canopy EX, Classic, Cobra, Command, Command Xtra, Domain, Dual MAGNUM, Dual II MAGNUM, FirstRate, Flexstar, Frontier, Fusion, Gaugster, Gauntlet, INTRRO, Lexone, Linex, Linuron, Lorox, Lorox Plus, Me-TooLachlor, Micro-Tech, Outlook, Pendimax, Prowl, Prowl H2O, Pursuit, Pursuit Plus, Python, Reflex, Resource, Scepter, Select, Select MAX, Sencor, Spartan, Squadron, Steel, Valor]

Postemergence (In-crop)

For use of this product to control annual grasses and broadleaf weeds listed on this label in Roundup Ready soybeans. Applications of this product can be made in from emergence (cracking) through flowering (R2 stage soybeans). R2 stage soybeans ends when a pod reaches 5 millimeters (0.2 inches) in length at one of the four uppermost nodes on the main stem with a fully developed leaf. Refer to the "ANNUAL WEEDS RATE SELECTION" in this label for rate recommendations for specific annual weeds. In general, an initial application of 1 quart per acre on 2- to 8-inch tall weeds is used. Weeds will generally be 2 to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 2 quarts per acre in any single in-crop application for control of annual weeds and where dense weed populations exist.

A 1- to 2-quarts per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem mully. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds. **IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE ROUNDUP READY SOYBEAN CROP.** To control giant ragweed, 1 quart per acre of this product be applied when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready soybeans. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to soybeans: acifluorfen, bentazonechlorimuron ethyl, clethodim, imazethapyr, cloransulam-methyl, fenoxypop, flazifop-phutyl, flumiclorac pentyl ester, fomesafen, imazamox, imazethapyr, lactofen, pendimethalin, quizalofop P-ethyl, sethoxydem, thifensulfuron-methyl.

Arrow, Assure II, Basagran, Classic, Cobra, Extreme, FirstRate, Flexstar, Fusilade DX, Fusion, Harmony GT XP, Poast, Poast Plus, Pursuit, Pursuit Plus, Raptor, Reflex, Select, Select MAX, Synchrony STS, Targa, Ultra Blazer.]

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in-crop application is 2 quarts per acre. The maximum combined total of this product that can be applied during flowering is 2 quarts per acre.

Preharvest

Apply for weed control prior to harvest of soybeans after pods have set and lost all green color. Up to 1 quart per acre of this product can be applied by aerial or ground application.

PRECAUTIONS, RESTRICTIONS: Care should be taken to avoid excessive seed shatter loss due to ground application equipment. Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

Post-Harvest

Apply this product after harvest of Roundup Ready soybeans. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

12.8 Roundup Ready Sugar Beet

<u>Maximum Annual Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Emergence to 8-leaf stage	2.5 quarts per acre
Between 8-leaf stage and canopy closure	2 quarts per acre

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of sugar beet. Read and follow label directions of all products in the tank mixture.

{Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplan, at-planting and/or preemergence to sugar beet: dimethenamid, S-metolachlor.

Dual MAGNUM, Frontier]

Postemergence (In-Crop)

Apply this over-the-top of Roundup Ready sugar beets for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate recommendations for specific annual weeds. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready sugar beet. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to sugar beet. Read and follow label directions of all products in the tank mixture

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to sugar beet: clethodim, clopyralid, desmedipham, ethofumesate, phenmedipham, quizalofop-p-ethyl, sethoxydim, trisulfuron-methyl.

Assure II, Betamix, Betanex, Norton SC, Poast, Progress, Select, Stinger, Upbeet]

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 4.5 quarts per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 1.5 quarts per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 1 quart per acre. Allow a minimum of 30 days *between* last application and sugar beet harvest.

13.0 NON-CROP USES AROUND THE FARMSTEAD

TYPES OF APPLICATIONS: General Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stump, Habitat Management.

13.1 General Weed Control, Trim-And-Edge

Apply to control annual weeds, perennial weeds and woody brush found in any part of the farmstead, including around building foundations and equipment storage areas, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, rangeland, rights-of-way, shelterbelts, and prior to planting landscape ornamentals. Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" in this label for treatment rates. For application through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "Annual Weeds -- Hand-Held or Backpack Equipment" section of this label for further instructions.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product used is registered for treatment of these non-crop sites. Refer to the individual product labels for approved farmstead sites and application rates. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Arsenal	Plateau
Barricade 65WG	Princep DF
Crossbow L	Princep 4L
Diuron	Ronstar 50 WP
Dicamba	Sahara
Endurance	Simazine

Escort	Surflan AS
Goal 2XL	Surflan WDG
Karmex DF	Telar
Krovar I DF	Transline
Outrider	Vanquish
Oust	Velpar DF
Pendimethalin	Velpar L
Pendulum 3.3 EC	2,4-D
Pendulum WDG	

This product plus dicamba tank mixtures may not be applied by air in California.

13.2 Greenhouse/Shadehouse

To control weeds in and around greenhouses and shade houses.

PRECAUTIONS, RESTRICTIONS: Desirable vegetation must not be present during application. Air circulation fans must be turned off. Do not use in residential greenhouses.

13.3 Chemical Mowing

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Use 16 fluid ounces of this product per acre when treating Bermudagrass. Use 64 fluid ounces of this product per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

PRECAUTIONS, RESTRICTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

13.4 Cut Stump

TYPES OF APPLICATIONS: Treating cut stumps in any non-crop site listed on this label.

Apply this product for control of growth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Salt-cedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, brazilian	Willow
Pine, Austrian	

PRECAUTIONS: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

13.5 Habitat Management

TYPES OF APPLICATIONS: Habitat Restoration and Maintenance, Wildlife Food Plots.

Habitat Restoration and Maintenance

Apply to control other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species, including Roundup ready canola, may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage. For specific product instructions for Roundup Ready canola wildlife food plots, see the "ROUNDUP READY CANOLA" section of this label.

RESTRICTIONS: Do not process treated Roundup ready canola seeds from Roundup ready canola wildlife food plots for food. Do not graze or feed treated Roundup Ready canola from wildlife food plots to livestock. There are no rotational restrictions for planting any wildlife food species or for allowing native species to repopulate the area following applications of this product.

14.0 ANNUAL WEEDS RATE SECTION

When water carrier volumes are between 16 and 40 gallons per acre for ground applications and between 6 and 15 gallons per acre for aerial applications, the following use rates will control the annual weeds listed in the table below:

- 1 quart per acre – grass and broadleaf annual weeds less than 6 inches in height or circumference and vines less than 3 inches in length.
- 3 pints per acre – grass and broadleaf annual weeds 6 to 12 inches in height or circumference and vines 3 to 6 inches in length.
- 2 quarts per acre – grass and broadleaf annual weeds greater than 12 inches in height or circumference and vines greater than 6 inches in length.

WHEN WATER CARRIER VOLUMES ARE BETWEEN 3 AND 15 GALLONS PER ACRE FOR GROUND APPLICATIONS AND BETWEEN 3 AND 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS, USE THE RATES SPECIFIED FOR INDIVIDUAL WEEDS AS FOLLOW IN THE "ANNUAL WEEDS RATE TABLE (Alphabetically by Species)".

Apply to actively growing annual weeds. Annual weeds are generally easiest to control when they are small.

Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Maximum size refers to the maximum plant height, length of runners for vines, or circumference of rosette plants in inches.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment. This product may be used up to 64 fluid ounces per acre where heavy weed densities exist.

ANNUAL WEEDS RATE TABLE
(Alphabetically by Species)

WEED SPECIES	RATE (Fluid Ounces Per Acre)				
	16	24	32	40	48
	(Maximum Size (in inches))				
Ammannia, purple	3	6	12	-	18
Anode, spurred	-	2	3	5	8
Barley	18	18+	-	-	-
Barnyardgrass	-	3	6	7	9
Basle, fivehook	-	-	6	-	-
Beggarweed, Florida	-	5	8	-	-
Bittercress	12	20	-	-	-
Bluegrass, annual	10	-	-	-	-

Bluegrass, bulbous	6	-	-	-	-
Brome, downy ^{1,2}	6	12	-	-	-
Brome, Japanese	6	12	24	-	-
Browntop panicum	6	8	12	-	24
Buckwheat, wild ³	-	1	2	-	-
Burcucumber	-	6	12	-	18
Buttercup	12	20	-	-	-
Carolina geranium	-	-	4	-	9
Carpetweed	-	6	12	-	-
Cheat ²	6	20	-	-	-
Chervil	20	-	-	-	-
Chickweed	-	12	18	-	-
Cocklebur	12	18	24	-	36
Copperleaf, hophornbeam	-	2	4	-	6
Copperleaf, Virginia	-	2	4	-	6
Coreopsis, plains	-	6	12	-	18
Corn, volunteer	6	12	20	-	-
Corn speedwell	12	-	-	-	-
Crabgrass	3	6	12	-	-
Crowfootgrass	-	-	6	-	12
Cutleaf evening primrose	-	-	3	-	6
Devilsclaw (unicorn plant)	-	3	6	-	-
Dwarf dandelion	12	-	-	-	-
Eastern manna grass	8	12	-	-	-
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
Falsedandelion	-	20	-	-	-
Falseflax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	6	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail, giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	6	12	-	-	-
Goosegrass	-	3	6	-	12
Grain sorghum (mild)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundsel, common	-	6	10	-	-
Hemp sesbania	-	2	4	6	8
Henbit	-	-	6	-	12
Horseweed/ Maretail (<i>Conyza canadensis</i>)*	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnson grass, seedling	6	12	18	-	24
Junglerice	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia ⁴	-	3 to 6	12	-	-
Lambsquarters*	-	6	12	-	20

Little barley	6	12	-	-	-
London rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morningglory, annual (<i>Ipomoea spp</i>)	-	-	3	-	6
Mustard, blue	6	12	18	-	-
Mustard, tansy	6	12	18	-	-
Mustard, tumble	6	12	18	-	-
Mustard, wild	6	12	18	-	-
Nightshade, black	-	4	6	-	12
Nightshade, hairy	-	4	6	-	12
Oats	3	6	18	-	-
Pigweed, Palmer*	-	12	18	24	-
Pigweed species*	-	12	18	24	-
Prickly lettuce	-	6	12	-	-
Purslane	-	-	3	-	6
Ragweed, common*	-	6	12	-	18
Ragweed, giant*	-	6	12	-	18
Red rice	-	-	4	-	-
Rye, volunteer/cereal ²	6	18	18+	-	-
Ryegrass species*	-	-	6	-	12
Sandbur, field	6	12	-	-	-
Sandbur, longspine	6	12	-	-	-
Shattercane	6	12	20	-	-
Shepherd's-purse	6	12	-	-	-
Sicklepod	-	2	4	-	8
Signalgrass, broadleaf	-	3	6	7	9
Smartweed, ladythumb	-	-	6	-	9
Smartweed, Pennsylvania	-	-	6	-	9
Sowthistle, annual	-	-	6	-	12
Spanishneedles	-	-	6	-	12
Speedwell, purslane	12	-	-	-	-
Sprangletop	6	12	20	-	-
Spurge, prostrate	-	6	12	-	-
Spurge, spotted	-	6	12	-	-
Spurry, umbrella	6	-	-	-	-
Stinkgrass	-	12	-	-	-
Sunflower	12	18	-	-	-
Swinecress	-	5	12	-	-
Teaweed/ Prickly sida	-	2	4	-	6
Texas panicum	6	8	12	-	24
Thistle, Russian ⁵	-	6	12	-	-
Velvetleaf	-	-	6	-	12
Virginia pepperweed	-	18	-	-	-
Waterhemp*	-	-	6	-	12
Wheat ²	6	12	18	-	-
Wheat, (overwintered)	-	6	12	-	18
Wild oats	3	6	18	-	-
Wild proso millet	-	6	12	-	18
Witchgrass	-	12	-	-	-
Woolly cupgrass	-	6	12	-	-
Yellow rocket	-	12	20	-	-

- ¹ For control of downy brome in no-till systems, use 24 fluid ounces per acre.
- ² Performance is better if application is made before this weed reaches the boot stage of growth.
- ³ Use 24 fluid ounces per acre of this product to control wild buckwheat in the cotyledon to 2-leaf stage. Use 32 fluid ounces per acre to control 2- to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 32 fluid ounces followed by 32 fluid ounces of this product per acre.
- ⁴ Do not treat kochia in the button stage.
- ⁵ Control of Russian thistle may vary based on environmental conditions and spray coverage. Whenever possible, a tank mixture with 2,4-D as described below may improve control.
- * A glyphosate resistant biotype has been confirmed. For additional information, refer to the "WEED RESISTANCE MANAGEMENT" section of this label. You may also visit on the Internet www.weedscience.org or www.weedresistancemanagement.com, or contact your Repar-Glypho, LLC, representative.

14.1 Annual Weeds — Tank Mixtures with 2,4-D or Dicamba or Tordon 22K

Better control of certain tough weeds can be achieved by tank mixing this product with 0.25 pound of dicamba, 0.5 pound of 2,4-D or 1 to 2 fluid ounces of Tordon 22K per acre. These other herbicides, combined with the rates of this product specified in the "ANNUAL WEEDS RATE TABLE" above, will control the following weeds up to the maximum height or length indicated: 6" — prickly lettuce, mare's tail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Tordon 22K only); 12" — cocklebur, lambsquarters, pigweed, Russian thistle (2,4-D only). For better control of common ragweed, giant ragweed, Pennsylvania smartweed, or velvetleaf with a maximum height of 6-inches, tank mix this product with 0.5 pound of 2,4-D per acre.

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA.

14.2 Annual Weeds—Hand-Held or Backpack Equipment

For control of weeds listed in the "ANNUAL WEEDS RATE TABLE" above, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 -percent solution.

For best results, use a 2-percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 5-percent solution for annual and perennial weeds and a 5- to 10-percent solution for woody brush and trees.

14.3 Annual Weeds — Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

Applications of 24 to 28 fluid ounces of this product plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 28 ounces for control), Downy brome, *Green* foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass and Kochia (add 0.12 pound of dicamba for control).

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines.

15.0 PERENNIAL WEEDS RATE SECTION

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage. Best results are obtained when soil moisture is adequate for active weed growth.

PERENNIAL WEEDS RATE TABLE

(Alphabetically by Species)

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alfalfa	1 - 2	3 - 10	2%
Apply after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.			
Alligatorweed	4	3 - 20	1.5%
For partial control, apply when most of the plants are in bloom. Repeat applications will be required to achieve control.			
Anise (fennel) ¹	-	-	1-2%
Apply as a spray-to-wet treatment.			
Bahiagrass²	3-5	3-20	2%
Bentgrass	3-5	10-20	2%
For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is used for best results.			
Bermudagrass	3 - 5	3 - 20	2%
For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to achieve control.			
Bermudagrass, water (knotgrass)	1 - 1.5	5 - 10	2%
Apply 3 pints of this product in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.			
Fall applications only: Apply 1 quart this product in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length.			
This product is not registered in California for use on water Bermudagrass.			
Bindweed, field	0.5-5	3 - 20	2%
Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.			
For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in			

late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 2 quarts of this product plus 0.5 pound of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. Irrigate at least once to promote active bindweed growth.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky	1 - 2	3 - 40	2%
----------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas	3 - 5	3 - 40	2%
------------------------	--------------	---------------	-----------

Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenfern	3 - 4	3 - 40	1 - 1.5%
--------------------	--------------	---------------	-----------------

Apply to fully expanded fronds that are at least 18 inches long.

Bromegrass, smooth	1 - 2	3 - 40	2%
---------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf	---	3 - 20	2%
-----------------------------	------------	---------------	-----------

For control, apply 2 quarts of this product plus 0.5 pound of dicamba per acre. For partial control, apply 1 quart of this product plus ½ pound of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed²	2 - 3	3 - 40	2%
--------------------------------------	--------------	---------------	-----------

Cattail²	3 - 5	3 - 40	2%
----------------------------	--------------	---------------	-----------

Clover; red or white¹	3 - 5	3 - 20	2%
---	--------------	---------------	-----------

Also for control, apply 16 to 32 fluid ounces of this product plus 0.5 to 1 pound of 1,4-D, in 3 to 10 gallons of water per acre.

Cogongrass	3 - 5	10 - 40	2%
-------------------	--------------	----------------	-----------

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to achieve control.

Dallisgrass²	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Dandelion¹	3 – 5	3 – 40	2%
------------------------------	--------------	---------------	-----------

Also for control, apply 1 pint of this product plus 0.5 of 2,4-D, in 3 to 10 gallons of water per acre.

Dock, curly¹	3 – 5	3 – 40	2%
--------------------------------	--------------	---------------	-----------

Also for control, apply 1 to 2 pints of this product plus 0.5 to 1 pound of 2,4-D, in 3 to 10 gallons of water per acre.

Dogbane, hemp	4	3 – 40	2%
----------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue (except tall) ¹	3 – 5	3 – 20	2%
--	--------------	---------------	-----------

Fescue, tall	1 – 3	3 – 40	2%
---------------------	--------------	---------------	-----------

Apply 3 quarts of this product per acre when most plants have reached boot-to-early seedhead stage of development.

Fall applications only: Apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.

Guineagrass	2 - 3	3 - 40	1%
--------------------	--------------	---------------	-----------

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 2 quarts of this product per acre for control. In the flatwoods region of Florida, 3 quarts of this product per acre are required for control.

Horsenettle¹	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Horseradish	4	3 – 40	2%
--------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Iccplant¹	---	---	1.5 – 2%
-----------------------------	------------	------------	-----------------

Thorough coverage is necessary for best control.

Jerusalem artichoke¹	3 – 5	3 – 20	2%
--	--------------	---------------	-----------

Johnsongrass	0.5 – 3	3 – 40	1%
---------------------	----------------	---------------	-----------

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 1 quart of this product per acre.

For burndown of Johnsongrass, apply 1 pint of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression)--Apply a 1-percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass	2 – 3	3 – 40	2%
--------------------	--------------	---------------	-----------

Knapweed	4	3 – 40	2%
-----------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Lantana	--	--	1-1.25%
----------------	-----------	-----------	----------------

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Lespedeza	3 – 5	3-20	2%
------------------	--------------	-------------	-----------

Milkweed, common	3	3-40	2%
-------------------------	----------	-------------	-----------

Apply when most plants have reached the late bud to flower stage of growth.

Muhly, wirestem	1-2	3-40	2%
------------------------	------------	-------------	-----------

Use 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications, or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.

Mullein, common¹	3 – 5	3 – 20	2%
------------------------------------	--------------	---------------	-----------

Napiergrass²	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Nutsedge, purple or yellow	0.5-3	3-40	1 - 2%
-----------------------------------	--------------	-------------	---------------

Apply 3 quarts of this product per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for longterm control of ungerminated tubers.

Sequential applications: 1 to 2 quarts of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For partial control of existing plants, apply 1 pint to 2 quarts of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

Orchardgrass	1 – 2	3-40	2%
---------------------	--------------	-------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of this product in 3 to 10 gallons

of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications, and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

Pampasgrass	--	--	1.5 - 2%
--------------------	----	----	----------

Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Paragrass²	3 - 5	3 - 20	2 %
------------------------------	-------	--------	-----

Phragmites	3 - 5	10 - 40	1 - 2 %
-------------------	-------	---------	---------

For partial control and best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may result in reduced control. Due to the dense nature of this vegetation that may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to achieve control. Visual control symptoms will be slow to develop.

Poison hemlock	--	--	1 - 2%
-----------------------	----	----	--------

For hand-held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Thorough coverage is necessary for best control.

Pokeweed, common	1	3-40	2%
-------------------------	---	------	----

Apply to actively growing plants up to 24 inches tall.

Quackgrass	1 - 3	3 - 40	2%
-------------------	-------	--------	----

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of this product. Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.

In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

Redvine	0.75-2	5 - 10	2%
----------------	--------	--------	----

For suppression, apply 1.5 pints of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, giant	--	--	2%
--------------------	----	----	----

Best results are obtained when applications are made in late summer to fall.

Ryegrass, perennial	1 - 3	3 - 40	1%
----------------------------	-------	--------	----

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1 quart of this product per acre.

Smartweed, swamp'	3-5	3-40	2%
Also for control, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall.			
Sowthistle, perennial	2-3	3-40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
Spurge, leafy		3-10	2%
For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most plants are 12 inches tall.			
Starthistle, yellow	2	10-40	2%
Best results are obtained when applications are made during the rosette, bolting and early flower stages.			
Sweet potato, wild			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, artichoke			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, Canada	2-3	3-40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
For suppression in the spring, apply 1 quart of this product, or 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.			
Timothy²	2-3	3-40	2%
Torpedograss	4-5	3-40	2%
For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to achieve control. Fall treatments must be applied before frost.			
Trumpet creeper	2	5-10	2%
For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.			
Vaseygrass²	3-5	3-20	2%
Velvetgrass²	3-5	3-20	2%

Cherry; bitter, black, pin	2-3	3-40	1.5%
Coyote brush	--	--	1.5-2%
Apply when at least 50 percent of the new leaves are fully developed.			
Dogwood ¹	2-5	3-40	1-2%
Elderberry	2-3	3-40	1-1.5%
Elm ¹	2-5	3-40	1-2%
Eucalyptus	--	--	2%
For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.			
Florida holly (Brazilian	2-5	3-40	1-2%
Peppertree) 1			
Gorse ¹	2-5	3-40	1-2%
Hasardia ¹ *2	--	--	1-2%
Hawthorn	2-3	3-40	1-1.5%
Hazel	2-3	3-40	1-1.5%
Hickory ¹	2-5	3-40	1-2%
Honeysuckle	3-4	3-40	1-1.5%
Hornbeam,	2-5	3-40	1-2%
American ¹			
Kudzu	4-5	3-40	2%
Repeat applications may be required to achieve control.			
Locust, black ¹	2-4	3-40	1-2%
Madrone resprouts ¹	--	--	2%
Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.			
Manzanita ¹	2-5	3-40	1-2%
Maple, red	2-4	3-40	1-1.5%
Apply a 1- to 1.5-percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre.			
Maple, sugar	--	--	1-1.5%
Apply when at least 50 percent of the new leaves are fully developed			
Monkey flower ²	--	--	1-2%
Oak; black,white ¹	2-4	3-40	1-2%
Oak, post	3-4	3-40	1-1.5%
Oak; northern,	--	--	1-1.5%
Apply when at least 50 percent of the new pin leaves are fully developed			
Oak; southern red	2-3	3-40	1-1.5%
Persimmon ¹	2-5	3-40	1-2%
Pine	2-5	3-40	1-2%
Poison ivy/Poison oak	4-5	3-40	2%
Repeat applications may be required to achieve control. Fall treatments must be applied before leaves lose green color.			
Poplar, yellow ¹	2-5	3-40	1-2%
Redbud, eastern	2-5	3-40	1-2%

Rose, multiflora	2	3-40	1 %
Treatments should be made prior to leaf deterioration by leaf-eating insects.			
Russian olive ¹	2-5	3-40	1 -2%
Sage, black ¹			1%
Sage, white ¹	2-5	3-40	1 -2%
Sage brush, California ²			1%
Salmonberry	2-3	3-40	1 -1.5%
Salt-cedar	2-5	3-40	1 -2%
Sassafras ¹	2-5	3-40	1 -2%
Sourwood ¹	2-5	3-40	1 -2%
Sumac; poison, smooth, winged ¹	2-4	3-40	1 -2%
Sweetgum	2-3	3-40	1 -1.5%
Swordfern ¹	2-5	3-40	1 -2%
Tallowtree, Chinese ¹			1%
Tan oak resprouts ¹	—	--	2%
Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.			
Thimbleberry	2-3	3-40	1 -1.5%
Tobacco, tree ¹			1 -2%
Trumpet creeper	2-3	3-40	1 -1.5%
Vine maple ¹	2-5	3-40	1 -2%
Virginia creeper	2-5	3-40	1 -2%
Waxmyrtle, Southern ¹	2-5	3-40	1 -2%
Willow	3-4	3-40	1 -1.5%

¹ Partial Control

² Thorough coverage of foliage is necessary for best results.

17.0 LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.

To the extent consistent with applicable law, this warranty is subject to the conditions and limitations stated on this labeling.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

To the extent consistent with applicable laws, this Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

For in-crop (over-the-top) uses on Roundup Ready crops, crop safety and weed control performance are not

warranted by Repar-Glypho, LLC when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAWS, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Amplify, Bullet, Degree, Farmsource, Harness, Lariat, , Micro-Tech, Monsanto and Vine symbol, and Permit are trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

II. MAIN LABEL FOR INDUSTRIAL, TURF, ORNAMENTAL USES

GLYPHOSATE 41% SL

The complete broad-spectrum postemergence herbicide for non-crop, industrial, turf and ornamental weed control.

Complete Directions for Use

EPA Reg. No. 86004-U

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product. Use only according to label instructions.

Not all products on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

[Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REAPR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC. REPACKAGING OR TOLL REPACKAGING AGREEMENT.

[Non-Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

1	1.0 INGREDIENTS
2	2.0 IMPORTANT PHONE NUMBERS
3	3.0 PRECAUTIONARY STATEMENTS 3.1 Hazards to Humans and Domestic Animals 3.2 Environmental Hazards 3.3 Physical or Chemical Hazards
4	4.0 STORAGE AND DISPOSAL
5	5.0 GENERAL INFORMATION (How This Product Works) 5.1 Weed Resistance Management 5.2 Management Recommendations for Glyphosate Resistant Weed Biotypes
6	6.0 MIXING 6.1 Mixing with Water 6.2 Tank Mixing 6.3 Tank Mixing Procedure 6.4 Mixing for Hand-Held Sprayers 6.5 Surfactants 6.6 Colorants or Dyes
7	7.0 APPLICATION EQUIPMENT AND TECHNIQUES 7.1 Aerial Equipment 7.2 Ground Broadcast Equipment 7.3 Hand-Held or Backpack Equipment 7.4 Selective Equipment 7.5 Injection Systems 7.6 CDA Equipment
8	8.0 SITE AND USE DIRECTIONS 8.1 Cut Stump 8.2 Forestry Site Preparation 8.3 General Non-crop Areas and Industrial Sites 8.4 Habitat Management 8.5 Hollow Stem Injection 8.6 Injection and Frill (Woody Brush and Trees) 8.7 Non-food Tree, Shrub, or Vine Production Sites 8.8 Railroads 8.9 Roadsides 8.10 Rangelands 8.11 Utility Sites 8.12 Conservation Reserve program (CRP) 8.13 Grass Seed or Sod Production 8.14 Pastures
9	9.0 WEEDS CONTROLLED 9.1 Annual Weeds 9.2 Perennial Weeds 9.3 Woody Brush and Trees
10	10.0 LIMIT OF WARRANTY AND LIABILITY

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....	41.0%
OTHER INGREDIENTS (Including surfactant).....	59.0%
	100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

(To be added later)

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

3.2 Environmental Hazards

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published REPAR-GLYPHO, I.I.C. Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. **PESTICIDE STORAGE:** Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

See container label for additional **CONTAINER DISPOSAL** instructions.

[Container Label Storage and Disposal Statements]:

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact Repar-Glypho, LLC.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

Refiners: Do not reuse this container except for refill in accordance with a valid Monsanto agreement for Repackaging or Toll Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration by burning, or, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

(How This Product Works)

Product Description: This product is a postemergent, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visual effects are gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at later growth stages approaching maturity.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates: The combined total of all treatments must not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate as the active ingredient, whether applied separately

or as mixtures. Calculate the application rates and ensure that the total use of this and other glyphosate containing products do not exceed stated maximum use rates. See the 'INGREDIENTS' section of this label for necessary information.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

5.1 Weed Resistance Management

GROUP	9	HERBICIDE
--------------	----------	------------------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using other cultural or mechanical practices.

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management practices:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Utilize the label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.

- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Repr-Glypho, LLC. representative, local retailer, or county extension agent.

5.2 Management Practices For Glyphosate Resistant Biotypes

Note: Appropriate testing is critical in order to determine if a weed is resistant to glyphosate. Contact your Repr-Glypho, LLC. representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

Control for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling for this product and can be obtained from your local retailer or Repr-Glypho, LLC. representative.

As the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Repr-Glypho, LLC. Company is not responsible for any losses that may result from the failure of this product to control glyphosate resistant weeds biotypes to the extent consistent with applicable laws.

The following good agronomic practices should be followed to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

6.0 MIXING

Spray solutions of this product should be mixed, stored, and applied using only clean stainless steel, aluminum, plastic, fiberglass, plastic or plastic-lined steel containers. Clean sprayer parts immediately after using this product by thoroughly flushing with water. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations.

6.1 Mixing with Water

NOTE: Product performance may be significantly reduced if water containing soil sediment is used as a carrier. Do not mix this product with water from ponds and ditches that is visibly muddy or murky.

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of clean water. Add the amount of this product near the end of the filling process and mix well. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

6.2 Tank Mixing

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

When this label gives directions for use of a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, or dicamba, the user is responsible for ensuring that the specific application is included on the label of the specific products used in the tank mixture. Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions and use according to the most restrictive label requirements. Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

To the extent consistent with applicable laws, buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly used in this label. Mixing this product with herbicides or other materials on this label may result in reduced performance.

6.3 Tank Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the amount of this product. If needed, add nonionic surfactant before completing the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" and "GENERAL INFORMATION" sections for additional precautions.

6.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product indicated in the following table with water:

Spray Solution

Desired Volume	Amount of GLYPHOSATE 41% SL					
	0.5%	1 %	1.5%	2%	5%	10%
1 gal	0.7 oz	1.3 oz	2 oz	2.7 oz	6.5 oz	13 oz
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

For backpack, knapsack or pump-up sprayers, the appropriate amount of this product be mixed with water in a larger container, and then used to fill the sprayer.

6.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

6.6 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to spray solutions of this product. However, they can reduce product performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system. Avoid direct application to any body of water.

APPLY SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial application.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversion" sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

- Application height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sunsets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use rates of this herbicide in 3 to 25 gallons of water per acre unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by ReapR-Glypho, LLC. for this product.

Drift reduction additives may be used. When a drift reduction additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application --to avoid streaking and uneven or overlapped application, use appropriate marking devices.

Aircraft Maintenance; Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may prevent corrosion.

7.2 Ground Broadcast Equipment

Apply at rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label for this product. As density of weeds increases, spray volume should be increased to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

7.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For low-volume directed spray applications, spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

7.4 Selective Equipment

This product may be diluted in water and applied through shielded applicators, hooded sprayers, wiper applicators or sponge bars to weeds listed on this label that are growing in any specified non-crop site.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators and Sponge Bars

Wiper applicators are devices that physically wipe the appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetations.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust the height of the applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper applicators when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from the use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators—Solutions ranging from 33 to 75 percent of this product in water may be used.

For Panel Applicators and pressure-feed systems--Solutions ranging from 33 to 100 percent of this product in water may be used.

When applied, this product **CONTROLS** the following weeds:

Corn, volunteer	Sicklepod
Panicum, Texas	Spanishneedles
Rye, common	Starbur, bristly Shattercane

When applied, this product **SUPPRESSES** the following weeds:

Beggarweed, Florida	Ragweed, common
Bermudagrass	Ragweed, giant
Dogbane, hemp	Smutgrass
Dogfennel	Sunflower
Guineagrass	Thistle, Canada
Johnsongrass	Thistle, musk
Milkweed	Vaseygrass
Nightshade, silverleaf	Velvetleaf
Pigweed, redroot	

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems.

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 INDUSTRIAL, TURF AND ORNAMENTAL SITES

Unless otherwise specified, applications may be made to control any weeds listed in the "WEEDS CONTROLLED" section 9.0 of this label. Refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" section for detailed instructions on different application methods.

8.1 Cut Stumps

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100-

percent solution of this product to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Salcedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow Pine, Austrian

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2 Forestry Site Preparation

This product is used for the control or partial control of woody brush, trees and herbaceous *weeds* in forestry. This product is also used in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is used in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

Use higher rates of this product for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

This product may be used in a tank mix with the following products for forestry site preparation.

Arsenal Applicators Concentrate
Chopper
Escort or Escort XP
Garlon 3A
Garlon 4A
Landmark XP
Oust or Oust XP
Westar

For control of herbaceous weeds, use the lower tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher tank mixture rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release, unless otherwise specified on this label for this product.

8.3 General Non-crop Areas, Industrial Sites

Use in areas such as airports, apartment complexes commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, rights-of-way, roadsides, shadehouses, sod or turf seed farms, storage areas, sports complexes, substations, turfgrass areas, utility sites, warehouse areas, and wildlife management areas

General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general non-crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product is registered for use on such non-crop sites. Refer to these products' labels for approved non-crop sites and application rates.

2,4-D	Goal 2XL	Princep™ 4L
Arsenal™	Karmex™ DF	Princep™ DF
atrazine	Krovar™ I DF	Ronstar™ 50WP
Barricade™ 65WG	Landmark II MP	Sahara™
Clarity	Milestone	simazine
Crossbow L	Oust	Surflan™ AS
dicamba	Oust XP	Surflan WDG
diuron	Outrider	Telar™
Endurance™	Pendulum WDG	Transline
Escort™	Pendulum™ 3.3 EC	Vanquish™
Gallery 75 DF	pendimethalin	Velpar DF
Garlon 4	Plateau™	Velpar L
Garlon™ 3A	Poast	

This product plus dicamba tank mixtures may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 1 to 2 quarts of this product plus 2 to 4 ounces of Oust or Oust XP per acre.

Bahiagrass	Johnsongrass
Bermudagrass	Poorjoe
Broomsedge	Quackgrass
Dallisgrass	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	
Fescue, tall	

Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 8 to 64 fluid ounces of this product per acre. Apply 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 16 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust or Oust XP in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate was greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

8.4 Habitat Management

Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.5 Hollow Stem Injection

This product may be applied through hand-held injection devices that deliver amounts of this product into targeted hollow-stem plants growing in any non-crop site specified on this label.

For control of the following hollow-stem plants, follow the use instructions below:

Japanese Knotweed, *Polygonum cuspidatum*

Inject 5 mL per stem of this product between second and third internode.

Bohemian Knotweed, *Polygonum bohemicum*

Inject 5 mL per stem of this product between the second and third internode.

Giant Hogweed, *Heracleum mantegazzianum*

Inject one leaf cane per plant 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Poison Hemlock, *Conium maculatum*

Inject one leaf cane per plant 10 to 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Field horsetail, *Equisetum arvense*

Inject one segment above the root crown with 0.5 mL per stem of this product. Use a small syringe that calibrates to this rate.

Canada Thistle, *Cirsium arvense*

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowed removed as 0.5 mL per stem of this product is injected into the stem.

Note: The combined total for all treatments must not exceed 10.6 quarts of this product per acre. At 5 mL per stem, 7 quarts should treat approximately 1300 stems per acre.

8.6 Injection and Frill (Woody Brush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 0.04 fluid ounce (1 mL of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50- to 100-percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100-percent concentration of this product. For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:

Control

Oak
Poplar
Sweetgum
Sycamore

Partial Control

Black gum
Dogwood
Hickory
Maple, red

8.7 Site Preparation for Planting of Non-Food Trees and Shrubs

Apply this product for general weed control prior to planting or around established ornamentals, or any woody tree, shrub, or vine species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce or yew, in any production site.

UNLESS OTHERWISE DIRECTED, THIS PRODUCT IS NOT FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be taken to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

TYPES OF APPLICATIONS: Site Preparation, Post-directed, Trim-and-edge, Wiper Application Site Preparation

This product may be used prior to planting any tree, shrub, or vine in an ornamental, nursery, or production setting, including Christmas tree species.

Post-directed, Trim-and-edge

This product may be used as a post-directed spray around established woody species, or to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a production setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Wiper Application

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs, or vines. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

8.8 Railroads

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along railroads.

Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

TANK MIXTURES: This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments, provided that the specific product is registered for use on such non-crop sites:

Arsenal	Oust XP
atrazine	Outrider
dicamba	Sahara DG
diuron	simazine
Escort	Spike™ 80 DF
Escort XP	Telar DF
Garlon 3A	Transline
Garlon 4	Vanquish
Hyvar™ X	Velpar DF
Hyvar X-L	Velpar L
Krovar I DF	2,4-D
Oust	

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 quarts per acre when using low volume and 10 quarts of this product per acre when using high volume as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.75- to 2-percent solution of this product when using high-volume spray-to-wet applications. Apply a 5- to 10-percent solution of this product when using low volume directed sprays for spot treatment.

TANK MIXTURES: This product may be mixed with the products listed above in this section for enhanced control of woody brush and trees along railroads, provided that the specific product is registered for use on such non-crop sites.

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestein, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestein, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpet creeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not done, since severe injury may occur.

8.9 Roadsides

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails, Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

TANK MIXTURES: This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments:

atrazine	Outrider
Clarity	Pendulum 3.3 EC
Crossbow I.	Pendulum WDG
dicamba	Plateau
diuron	Princep DF
Endurance	Princep 4L
Escort	Roustar 50WP
Escort XP	Sahara
Gallery 75DF	simazine
Krovar I DF	Surflan

Landmark 11 MP	Telar
Landmark MP	Vanquish
Landmark XP	2,4-D
Oust	

See the "MIXING" section of this label for general instructions for tank mixing.

Release of Bermudagrass or Bahiagrass

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

TANK MIXTURES: This product may also be tank-mixed with Outrider, Oust or Oust XP for residual control. These tank mixtures may delay greenup.

Apply 8 to 64 fluid ounces of this product in a tank mix with 0.75 to 1.3 ounces of Outrider per acre., Read and follow all label directions for Outrider. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust or Oust XP in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Oust or Oust XP per acre on Bermudagrass and no more than 0.5 ounce of Oust or Oust XP per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in 10 to 40 gallons of water per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestein, silver	Trumpet creeper
Fescue, tall	Vaseygrass

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 9 to 32 fluid ounces of this product with 0.75 to 1.3 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Bluestein, silver	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Trumpet creeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not done, since severe injury may occur.

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product *per* acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 6 fluid ounces of this product with 0.75 to 2 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weed greater than 6 inches in height. Use only on well-established bahiagrass.

A tank mixture of this product plus Oust or Oust XP may be used. Apply 6 fluid ounces of this product plus 0.25 ounce of Oust or Oust XP per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.10 Rangelands

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands, pastures, and industrial sites. Preventing weed seed production is critical to the successful control of annual weed grasses invading these perennial grass sites. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Bromus: This product may be used to control or suppress downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*), cheatgrass (*Bromus secalinus*), cereal rye and jointed goatgrass found in rangelands pastures and industrial sites. Apply 8 to 16 fluid ounces of this product per acre on a broadcast basis. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead: To control or suppress medusahead, apply 16 fluid ounces of this product per acre at the 3-leaf stage when plants are actively growing. Delaying applications beyond this stage will result in reduced or unacceptable control. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses. Applications may be made in the fall or spring.

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

Spot Treatment, Wiper Application

This product may be applied in rangeland, pastures or industrial sites as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals. For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire site or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acres, no more than 10 percent of the total site may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

8.11 Utility Sites

In utilities, this product is for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals,

flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product is also for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

For control of herbaceous weeds, use the lower tank mixture rates. For control of dense stands or woody brush and trees, use the higher tank mixture rates.

Bare Ground, Trim-and-Edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products provided that the specific product used is registered for application on these sites. Refer to the individual product's labels for approved sites and application rates.

2,4-D	Outrider
Arsenal	pendimethalin
atrazine	Plateau
Barricade 65WG	Princep
dicamba	Ronstar 50WP
diuron	Sahara
Endurance	simazine
Escort	Surflan AS
Escort XP	Surflan WDG
Garlon 3A	Telar DF
Garlon 4	Transline
Krenite	Vanquish
Krovar I DF	Velpar DF
Oust	Velpar L
Oust XP	

Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray incompatibility problems. For side trimming treatments, this product may be used alone or in a tank mixture with Garlon 4.

8.12 Conservation Reserve Program (CRP)

This product may be used for renovation (rotating out of CRP), site preparation, postemergence weed control in dormant CRP grasses, or wiper application on CRP land.

Renovation (Rotating out of CRP), Site Preparation

This product may be used to prepare CRP land for crop production. Refer to Federal, State or local use guides for CRP renovation recommendations.

Postemergence Weed Control in Dormant CRP Grasses, Wiper Application

Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land.

Applications may be made using wiper applicators to control tall weeds, or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 8 to 12 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Use the lower rate when treating weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

8.13 Grass Seed or Sod Production

This product may be used in grass seed and sod production for preplant, at-planting, preemergence, removal of established stands, renovation, site preparation, shielded spraying, wiper application, spot treatment, and creating rows in annual ryegrass.

Preplant, Preemergence, At-Planting, Removal of Established Stands, Renovation, Site Preparation

This product controls most existing vegetation for purposes of renovating turf or forage grass seed areas or for establishing turfgrass grown for sod. It may also be used to destroy remaining undesired grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting, or for renovation purposes. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species. Use the lower rate when grass seed or sod production is below 6 inches in height (or runner length). Use the higher rate when grass or sod is greater than 6 inches in height.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Applications must be made prior to crop emergence in order to avoid crop injury.

Shielded Sprayers

Apply 1 to 3 quarts of this product in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Use the lower rate when the weeds are below 6 inches in height (or runner length). Use the higher rate when the weeds are greater than 6 inches in height. Uniform planting in straight rows aids in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by the protective shields.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction.

Wiper Application

This product may be applied over the top of desirable grasses using wiper applicators for the control of tall weeds.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction.

Spot Treatment

Apply a 1.5-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

The crop sprayed in the treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

Creating Rows in Annual Ryegrass

Apply 1 to 2 pints of this product per acre. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height. Use the higher rate when ryegrass is greater than 6 inches in height.

Set nozzle heights to allow the establishment of the desired row spacing. Use low-pressure nozzles, or drop nozzles designed to target the application over a narrow band.

Take care not to spray or allow spray to drift outside target area in order to avoid unwanted crop destruction.

8.14 Pastures

This product may be applied to any pasture grass (*Gramineae* family), including bahiagrass, Bermudagrass, bluegrass, brome, fescue, guineagrass, kikuyugrass, orchardgrass, pangola grass, ryegrass, timothy, and wheatgrass. Application can be made as a spot treatment, wiper application, preplant, preemergence, pasture renovation, or postemergent broadcast.

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be applied to control perennial pasture species listed on this label prior to replanting.

For preplant, preemergence or pasture renovation application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When preplant, preemergence or pasture renovation application methods are made using rates above 3 quarts of this product per acre, no more than 10 percent of the total pasture may be treated at any one time. Use the lower rate when weeds are below 6 inches in height (or runner length). Use the higher rate when weeds are greater than 6 inches in height.

If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment, Wiper Application

This product may be applied in pastures as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper application are made using rates above 3 quarts of this product per acre, no more than 10 percent of the total pasture may be treated at any one time. Use the lower rate when weeds are below 6 inches in height (or runner length). Use the higher rate when weeds are greater than 6 inches in height. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

Postemergent Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Use the lower rate when weeds are below 6 inches in height (or runner length). Use the higher rate when weeds are greater than 6 inches in height. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts of this product per acre per year onto pasture grasses except for renovation uses as described previously in this section.

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for rates to the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results.

9.1 Annual Weeds

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions. Use the higher rate for certain species regardless of the weed size at application. Treat tough-to-control weeds early when they are relatively small. This product may be tank mixed provided the tank-mix product is registered for use on the target site. Refer to the individual product labels for approved sites and application rates. For spray-to-wet applications, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds.

For low volume directed spray applications, use a 4- to 7-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or

where there are multiple spouts.

WEED SPECIES

Anoda, spurred	Lamb's-quarters*
Barley*	Little barley*
Barnyardgrass*	London rocket*
Bittercress*	Mayweed
Bassia, fivehook	Medusahcad*
Black nightshade*	Morningglory (<i>Ipomoea spp</i>)
Bluegrass, annual*	Mustard, blue*
Bluegrass, bulbous*	Mustard, tansy*
Brome, downy*	Mustard, tumble*
Brome, Japanese*	Mustard, wild*
Browntop panicum*	Oats
Buttercup*	Pigweed*
Carolina foxtail*	Plains/Tickseed coreopsis*
Carolina geranium Castor bean	Prickly lettuce*
Cheatgrass*	Puncturevine
Cheeseweed (<i>Malva parviflora</i>)	Purslane, common
Chervil*	Ragweed, common*
Chickweed*	Ragweed, giant
Cocklebur*	Red rice
Copperleaf, hophornbeam	Russian thistle
Com*	Rye*
Com speedwell*	Ryegrass*
Crabgrass*	Sandbur, field*
Dwarf dandelion*	Shepherd's-purse*
Eastern manna grass*	Sicklepod
Eclipta*	Signalgrass, broadleaf*
Fall panicum*	Smartweed, ladythumb*
Falsedandelion*	Smartweed, Pennsylvania*
Falseflax, smallseed*	Sowthistle, annual
Fiddleneck	Sowthistle, annual
Field pennycress*	Spanishneedles
Filaree	Speedwell, purslane*
Fleabane, annual*	Sprangletop*
Fleabane, hairy (<i>Conyza bonariensis</i>)*	Spurge, annual
Fleabane, rough*	Spurge, prostrate*
Florida pulley	Spurge, spotted*
Foxtail*	Spurry, umbrella*
Goatgrass, jointed*	Starthistle, yellow
Goosegrass	Stinkgrass*
Grain sorghum (milo)*	Sunflower*
Groundsel, common*	Teaweed/ Prickly sida
Hemp sesbania	Texas panicum*
Henbit	Velvetleaf
Horseweed/Marestail (<i>Conyza canadensis</i>)	Virginia copperleaf
Itchgrass*	Virginia pepperweed*
Johnsongrass, seedling	Wheat*
Junglerice	Wild oats*
Knotweed	Witchgrass*
Kochia	Woolly cupgrass*
	Yellow rocket

*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 1 pint of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure through coverage of foliage and treat when weeds are in an early growth stage.

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate.

Ensure thorough coverage when using spray-to-wet treatments with hand-held equipment. For best results, use a 1.5-percent solution on tough to control weeds such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

For low volume directed spray applications, use a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Allow 7 or more days after application before tillage.

WEED SPECIES	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2 - 4	1 - 2
Bahiagrass	3 - 5	2
Beachgrass, European (<i>Ammophila arena</i>)	-	5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4 - 5	2
Bluegrass, Kentucky	2	2
Blueweed, Texas	4 - 5	2
Brackenfern	3 - 4	1 - 1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf	-	2
Canarygrass, reed	2 - 3	2
Cattail	3 - 5	2
Clover, red, white	3 - 5	2
Cogongrass	3 - 5	2
Dallisgrass	3 - 5	2
Dandelion	3 - 5	2
Dock, curly	3 - 5	2
Dogbane, hemp	4	2
Fescue (except tall)	3 - 5	2
Fescue, tall	1 - 3	2
German ivy	2 - 4	1 - 2
Guineagrass	3	1
Horsenettle	3 - 5	2
Horseradish	4	2
Iceplant	2	1.5-2
Jerusalem artichoke	3 - 5	2
Johnsongrass	2 - 3	1
Kikuyugrass	2 - 3	2
Knapweed	4	2
Lantana	-	1 - 1.25
Lespedeza	3 - 5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3 - 5	2
Napiergrass	3 - 5	2
Nightshade, silverleaf	2	2

Nutsedge, purple, yellow	3	1 - 2
Orchardgrass	2	2
Pampasgrass	3 - 5	1.5 - 2
Paragrass	3 - 5	2
Pepperweed, perennial	4	2
Phragmites*	3 - 5	1 - 2
Poison hemlock	2 - 4	1 - 2
Quackgrass	2 - 3	2
Reed, giant	4 - 5	2
Ryegrass, perennial	2 - 3	1
Smartweed, swamp	3 - 5	2
Spurge, leafy*	-	2
Sweet potato, wild*	-	2
Thistle, artichoke	2 - 3	1 - 2
Thistle, Canada	2 - 3	2
Timothy	2 - 3	2
Torpedograss*	4 - 5	2
Trumpetcreeper*	2 - 3	2
Vaseygrass	3 - 5	2
Velvetgrass	3 - 5	2
Wheatgrass, western	2 - 3	2

* Partially Controlled

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

For best results when using hand-held equipment, use a 1.5-percent solution on harder-to-control woody brush and trees. For low volume directed-spray applications, apply a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50-percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple spouts.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

WEED SPECIES	Rate (QT/A)	Hand-Held % Solution
Alder	3-4	1-1.5
Ash*	2-5	1-2
Aspen, quacking	2-3	1-1.5
Bearclover (Bearnat)*	2-5	1-2
Beech*	2-5	1-2
Birch	2	1
Blackberry	3-4	1-1.5
Blackgum	2-5	1-2
Bracken	2-5	1-2
Broom; French, Scotch	2-5	1.5-2
Buckwheat, California*	2-4	1-2
Cascara	2-5	1-1.5
Catclaw*	-	1-1.5
Ceanothus*	2-5	1-2
Chamise*	2-5	1
Cherry; bitter, black, pin	2-3	1-1.5
Coyote brush	3-4	1.5-2
Deerweed	2-5	1
Dogwood*	2-5	1
Elderberry	2	1
Elm*	2-5	1-2
Eucalyptus	--	2
Gorse*	2-5	1-2
Hasardia*	2-4	1-2
Hawthorn	2-3	1-1.5
Hazel	2	1
Hickory*	2-5	1-2
Honeysuckle	3-4	1-1.5
Hornbeam, American*	2-5	1-2
Kudzu	4	2
Locust, black*	2-4	1-2
Madrone resprouts*	--	2
Manzanita*	2-5	1-2

Maple, red	2-4	1-1.5
Maple, sugar	—	1-1.5
Monkey flower*	2-4	1-2
Oak; black, white*	2-4	1-2
Oak, post	3-4	1-1.5
Oak; northern, pin	2-4	1-1.5
Oak, Scrub*	2-4	1-1.5
Oak; southern red	2-3	1-1.5
Peppertree, Brazilian (Florida holly)*	2-5	1-2
Persimmon*	2-5	1-2
Pine	2-5	1-2
Poison ivy	4-5	2
Poison oak	4-5	2
Poplar, yellow*	2-5	1-2
Redbud, eastern	2-5	1-2
Rose, multiflora	2	1
Russian olive*	2-5	1-2
Sage, black	2-4	1
Sage, white*	2-4	1-2
Sage brush, California	2-4	1
Salmonberry	2	1
Salcedar*	2-5	1-2
Sassafras*	2-5	1-2
Sourwood*	2-5	1-2
Sumac; laurel, poison, smooth, sugarbur	2-4	1-2
Sweetgum	2-3	1-1.5
Swordfern*	2-5	1-2
Tallowtree, Chinese	-	1
Tan oak resprouts*	-	2
Thimbleberry	2	1
Tobacco, tree*	2-4	1-2
Toyon*	-	2
Trumpet creeper	2-3	1-1.5
Vine maple*	2-5	1-2
Virginia creeper	2-5	1-2
Waxmyrtle, southern*	2-5	1-2
Willow	3	1
Yerbasanta*	-	2

* Partially Controlled

10.0 LIMIT OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, this warranty is also subject to the conditions and limitations stated on this labeling. Repar – Glypho, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.

Buyer and all users shall promptly notify Repar – Glypho, LLC of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the

application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

Repar – Glypho, LLC does not warrant any product reformulated or repackaged from this product except in accordance with Repar – Glypho, LLC's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

EPA Reg. No. 86004-U
Repar-Glypho, LLC.
Silver Springs, MD 20910

III. SUPPLEMENTAL LABELING

Table of Contents: Supplemental labeling

	Name
A	FOR USE IN DORMANT ALFALFA
B	FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER FIRST CUTTING
C	FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS
D	FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARETAIL, <i>Conyza canadensis</i>) IN COTTON, CORN, AND SOYBEANS
E	HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROL IN ROUNDUP READY CANOLA
F	FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRENCO COUNTY
G	FOR AERIAL APPLICATION IN ARKANSAS ONLY
H	FOR CONTROLLING BARNYARDGRASS (ECHINOCHLOA CRUS-GALLI) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY
I	FOR WEED CONTROL APPLICATION IN SEED PRODUCTION OF ALFALFA WITH THE ROUNDUP READY® GENE
J	FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY
K	BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS
L	FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS
M	GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS
N	FOR GROUND AND AERIAL APPLICATION TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY
O	GLYPHOSATE 41% SL HERBICIDE FOR CONIFER RELEASE

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **GLYPHOSATE 41% SL** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **GLYPHOSATE 41% SL** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

GLYPHOSATE 41% SL Herbicide

EPA Reg. No. 86004-U

Keep out of the reach of children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any other manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIREABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for **GLYPHOSATE 41% SL** for essential product performance information.

[INSERT SPECIFIC DIRECTIONS FOR USE TEXT FROM THE FOLLOWING SECTIONS]

Read the "Limit of Warranty and Liability" in the label booklet for **GLYPHOSATE 41% SL** before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

REPAR-GLYPHO, LLC. COMPANY
SILVER SPRING, MD 20910

A. FOR USE IN DORMANT ALFALFA

DIRECTIONS FOR USE

This product will control or suppress many weeds, including quackgrass, downy brome, and cheatgrass in dormant alfalfa.

Apply 8 to 12 fluid ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with **GLYPHOSATE 41% SL**.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product can cause crop injury.

B. FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER THE FIRST CUTTING

DIRECTIONS FOR USE

This product may be applied at 16 fluid ounces per acre to control the weeds listed below and most other winter annual grass and broadleaf weeds in established coastal Bermudagrass pastures.

Annual bluegrass	Oats
Cheat	Ryegrass, Italian
Crabgrass	Sandbur, field
Henbit	Sunflower
Johnsongrass, seedling	Wheat
Little barley	Wild mustard

TIMING OF APPLICATION

Applications prior to spring growth: Apply this product in either late winter or early spring but before new coastal Bermudagrass growth begins in the spring. Applications to new growth can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 60 days after making this application before grazing or harvesting the treated area.

Application following the first cutting: Apply this product after the first Bermudagrass cutting when the Bermudagrass has not yet begun to regrow. Applications made after regrowth has begun can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 28 days after making this application before grazing or harvesting the treated area.

NOTE: ONLY ONE APPLICATION PER YEAR MAY BE MADE TO ANY ONE FIELD. A SPRING APPLICATION PRIOR TO GROWTH AND AN APPLICATION FOLLOWING THE FIRST CUTTING MAY NOT BE MADE ON THE FIELD DURING THE SAME YEAR.

C. FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS (WHEAT AND BARLEY)

DIRECTIONS FOR USE

Refer to the **GLYPHOSATE 41% SL** label for rates and weeds controlled.

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 2 to 3 gallons of water per acre.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, draft, or splash onto desirable vegetation because minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray to drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets.

In aerial applications, do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Ensure uniform application. Use appropriate marking devices when applying herbicides by air.

Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR IS MOST SUSCEPTABLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

D. FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARETAIL, *Conyza Canadensis*) IN CROP AND NON-CROP AREAS.

DIRECTIONS FOR USE

For ground applications, use 10 to 20 gallons of water per acre. For aerial applications, use 3 to 15 gallons of water per acre. **For tank-mix uses, read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental or fact sheets published separately for all herbicides used.**

Cotton:

Preplant:

For the control of horseweed, apply this product (32 fluid ounces per acre) in a tank-mix with Clarity® (8 fluid ounces per acre). This application must be made 21 to 35 days before planting and before horseweed reaches 6 inches in height. In order to avoid crop injury, a minimum interval of 21 days during which there is at least 1 inch cumulative rainfall must be observed between Clarity application and planting of cotton.

2,4-D may be included in the tank-mixture with this product. Refer to the 2,4-D product label for the time intervals that are required between application and planting and other geographic use restrictions.

Post-directed (Roundup Ready® Cotton and Roundup Ready Flex cotton® varieties only):

Management of early season weed competition and the development of a crop height differential between cotton and horseweed is often achieved by a combination of preplant burndown and post emergence over-the-top and/or directed applications of this product. These measures enhance the development of a height differential that is necessary to successfully make post-directed treatments. In-crop post-directed applications of MSMA (2 pounds active ingredient per acre) tank-mixed with diuron (0.5 to 0.75 pounds active ingredient per acre) should be made when the temperature is 80°F or higher.

Soybeans:

Preplant:

It is strongly encouraged that horseweed be controlled prior to planting. Apply a tank mixture of this product (32 fluid ounces per acre) with 2,4-D (0.5 pounds a.i. per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

In-crop (Roundup Ready Soybean varieties only):

This treatment should be used as a salvage treatment only for a horseweed infestation that was not controlled prior to planting. At the time of treatment, horseweed should not exceed 6 inches in height. Apply a tank mixture of this product (32 fluid ounces per acre) with FirstRate® (0.3 ounces per acre). Application should be made between full emergence of the first trifoliate leaf and 50% flowering stage of soybeans.

Corn:

Preplant, At-Planting, Preemergence:

Apply a tank mixture of this product (32 fluid ounces per acre) plus 2,4-D (0.5 pounds i.e. or per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Atrazine (1 to 2 pounds active ingredients per acre) may be included in the tank-mixture to provide residual control. Refer to the atrazine product label for specific use instructions.

In-crop (Roundup Ready Corn hybrids only):

Apply a tank-mixture of this product (32 fluid ounces per acre) plus Clarity (8 to 16 ounces per acre) or 2,4-D (0.5 to 1.0 pounds of inactive ingredients per acre) between corn emergence and the 5-leaf stage of growth (approximately 8 inches tall).

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Tree and Vine Crops:

Orchards (Pome Fruit, Stone Fruit and Tree Nuts).

Apply two quarts of this product plus 2,4-D (18 fluid ounces per acre Dri-Clean® or 2 pints per acre Orchard Master® CA) before marestail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is suggested. Follow labeling for 2,4-D product with regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply.

Vine Crops (grapes only):

Apply 2 quarts of this product per acre plus 2,4-D (18 fluid ounces per acre Dri-Clean®) before marestail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is suggested. Follow Dri-Clean® labeling in regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply. Also, residual herbicides such as diuron may provide additional preemergence control.

General Non-crop Areas:

Apply a tank mixture of this product (1 quart per acre) before horseweed exceeds 6 inches in height. Control may be enhanced by making applications when horseweed is still in the rosette stage of growth. This product may be tank-mixed with the following products provided that the specific product is registered for use on the target site. Refer to these product labels for approved sites and application rates. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive precautionary statements for each product in the mixture.

- 2,4-D
- Crossbow™
- Dicamba
- Gallery™ 75 DF
- Krovar™ I DF
- Landmark™ II MP
- Landmark™ MP
- Milestone™
- Overdrive®
- Telar™ DF
- Vanquish™
- Velpar® DF

E. HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROL IN ROUNDUP READY CANOLA

DIRECTIONS FOR USE

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A ROUNDUP READY GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELEWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA.

Maximum Allowable Combined Application Quantities Per Season	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application for emergence to 6-leaf stage	2 quarts per acre

Preplant, At-planting Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Weeds Controlled: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" in this label

Single Application- Apply 16 to 32 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application- Apply 32 fluid ounces per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but not later than the 6-leaf stage. Sequential applications are used for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass or when controlling weeds with multiple application times.

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in the Roundup Ready crops. No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 64 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

F. FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRESNO COUNTY, CALIFORNIA

DIRECTIONS FOR USE

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this supplemental labeling and in the product label booklet. Refer to Aerial Equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the product label for additional information. Refer to the individual use site section of the product label, or to other supplemental labeling or technical fact sheets published separately for this product by Repar-Glypho, LLC., for specific use instructions.

AVOID DRIFT-- DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetations after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops:

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).

3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.
5. Apply by air only to nonresidential areas.

When applied under the conditions described, this product controls annual and perennial weeds as listed in the label booklet

When tank mixing product with 2,4-D, only 2,4-D amine formulations may be used for aerial applications in California. Tank mixtures 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and alfalfa and pasture renovation applications only. This product, when tank mixed with dicamba, may not be applied by air in California.

ADDITIONAL INFORMATION FOR FRESNO COUNTY, CALIFORNIA

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California.

North:	Fresno County line
South:	Fresno County line
East:	State Highway 99
West:	Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Directions

A written recommendation MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation MUST state the proximity or surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rate of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to Fresno County Agricultural Commissioner.

Applications at Night- Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

G. FOR AERIAL APPLICATION IN ARKANSAS ONLY.

DIRECTIONS FOR USE

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when winds are in excess of 10 mph.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 mph is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 mph toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

H. FOR CONTROLLING BARNYARDGRASS (*ECHINOCHLOA CRUS-GALLI*) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY

DIRECTIONS FOR USE

Renovation Treatment

USE INSTRUCTIONS: This product may be applied as a renovation treatment in rice crops to control barnyardgrass infestations using ground broadcast spray or hand-held equipment. Renovation is defined as

herbicide treatment that will produce crop and weed destruction in an entire field or contiguous area treated within a field. Follow the application methods and treatment rates in the label booklet for **GLYPHOSATE 41% SL** herbicide.

PRECAUTIONS, RESTRICTIONS: Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction. The rice straw and stubble from the treated area, including a 25-foot buffer zone on all sides, shall not be used for grazing, animal bedding or any feed purposes.

Aerial applications are not permitted for rice renovation using this supplemental label.

I. FOR WEED CONTROL APPLICATIONS IN SEED PRODUCTION OF ROUNDUP READY® ALFALFA
--

General Information

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUITS OR CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for GLYPHOSATE 41% SL for essential product performance information.

The Roundup Ready designation indicates that the alfalfa contains a patented gene that provides tolerance to this product. Information on Roundup Ready Alfalfa varieties may be obtained from your seed supplier or Repar-Glypho, LLC representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

See the "ROUNDUP READY CROPS" section of the GLYPHOSATE 41% SL label booklet for general precautionary instructions for use in Roundup Ready crops. Do NOT combine the instructions in this supplemental label with other uses listed in the "PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS" section of the **GLYPHOSATE 41% SL** label booklet intended for alfalfa varieties that do not contain a glyphosate tolerance gene.

DIRECTIONS FOR USE

This product will control many troublesome emerged weeds with over-the-top applications in Roundup Ready alfalfa grown for seed. Over-the-top applications may be made from emergence through the late vegetative stage, and spot treatments may be made from early bud stage through seed harvest.

For ground applications with broadcast equipment, apply this product in 3 to 40 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial application: Use rates of this product in 3 to 15 gallons of spray solution per acre.

DO NOT EXCEED 2 QUARTS OF THIS PRODUCT PER ACRE WHEN MAKING APPLICATIONS BY AIR. FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATION IN THAT STATE. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready alfalfa. Follow the cleaning procedures specified on the label of the product(s) used. Alfalfa is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence and Post-harvest of seed

Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Total Preplant, At-planting and Preemergence Applications	2 quarts per acre
Total in-crop application rate from emergence through the late vegetative stage*	6 quarts per acre
Spot-treatment during early bud stage through seed harvest (See the "Spot Treatment after late vegetative stage" section and the "PRECAUTIONS AND RESTRICTIONS" section of this label for complete instructions)	Apply spray-to-wet; do not apply to the point of runoff

*There are no rotational crop restrictions following applications of this product. For any crop NOT listed in the label booklet, applications must be at least 30 days prior to planting.

Over-the-top applications: Broadcast applications of this product may be made using ground or aerial application equipment over the top of Roundup Ready alfalfa from emergence through the late vegetative stage. Do not make broadcast applications of this product between the initiation of alfalfa budding and the harvest of seed. Any single over-the-top broadcast application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart.

Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings are susceptible and will not survive or thrive after the first application of this product. A single application of at least 1 quart per acre of this product should be applied at or before the 3 to 4 trifoliate growth stage to eliminate the effects of stand gaps created by the loss of non-Roundup Ready plants.

Spot Treatment after late vegetative stage: For late emerging weeds, this product may be applied as a spot treatment in Roundup Ready alfalfa grown for seed, during the early bud stage through seed harvest. Applications made during this stage may result in reduced seed yield and quality and are the responsibility of the grower. Make applications on a spray-to-wet basis. Do not spray to the point of runoff. If a spot treatment is made after the late vegetative stage, harvested seed must not be used for alfalfa sprout production.

Post-harvest applications: Following harvest of Roundup Ready alfalfa seed, the stand may be managed for forage and hay production. Refer to the Roundup Ready alfalfa section of the label booklet GLYPHOSATE 41% SL for rates and timing of applications for forage and hay production.

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" in the GLYPHOSATE 41% SL label booklet. Some weeds with multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some re-growth of weeds has occurred.

In addition to those weeds listed in the GLYPHOSATE 41% SL label booklet, this product will suppress or control the parasitic weed, Dodder (*Cuscuta spp.*) in Roundup Ready alfalfa seed production. Repeat applications may be necessary for complete control.

Tank mixtures with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control and are NOT for over-the-top applications of this product.

RESTRICTIONS: Do not make over-the-top broadcast applications of this product between the initiation of alfalfa budding and the harvest of Roundup Ready alfalfa seed. The use of harvested Roundup Ready alfalfa seed is not suitable for, and is not for, production of alfalfa sprouts.

Read the "LIMIT OF WARRANTY OR LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using. For (in-crop) over-the-top uses on Roundup Ready crop varieties, crop safety and weed control performance are not warranted by Repar-Glypho, LLC, when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

J. FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY

DIRECTIONS FOR USE

GENERAL INFORMATION

The use of the over-the-top applications described in this label on *other* than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (nodes) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this supplemental label are specific to, and should only be used with varieties designated as **Round Ready Flex cotton**. **DO NOT** combine the instructions in this supplemental label with those in the "Roundup Ready Cotton" or "Roundup Ready Flex Cotton" sections found in the label booklet for **GLYPHOSATE 41% SL**, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instruction on labeling for this or other glyphosate-containing product. See "Maximum Use Rate" in the "GENERAL INFORMATION" section of the label booklet for **GLYPHOSATE 41% SL** for additional information.

ROUNDUP READY FLEX COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop), Preharvest

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all application	8 quarts per acre
Calculate the combined rate to be used for all preplant, in-crop and preharvest applications, to ensure that the total does not exceed the maximum allowed rate per acre year shown above.	
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to 60% percent open bolls	6 quarts per acre
Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 fluid ounces per acre

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of the **GLYPHOSATE 41% SL** label booklet for general precautionary instructions for use in Roundup Ready crops.

Preplant, At-planting, Preemergence

This product may be applied before, during or after planting Roundup Ready Flex cotton.

Postemergence (In-crop)

When applied in accordance with this label, GLYPHOSATE 41% SL will control labeled annual grasses and broadleaf weeds in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds. This product may be applied by ground application equipment at rates up to 2 quarts per acre per application postemergence to Roundup Ready Flex cotton. In addition to broadcast applications, post-directed equipment may be used to achieve weed coverage.

NOTE: For specific rates of application and instructions, refer to the "ANNUAL" and "PERENNIAL WEEDS RATE TABLES" in the label booklet for **GLYPHOSATE 41% SL**.

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 2 quarts per acre made using ground application equipment. **In-crop application rates above 1 quart per acre made alone, or with the addition of other crop chemical products containing surfactant, may cause a crop response, including leaf speckling or leaf necrosis.** Do not exceed a maximum rate of 3 pints per acre of this product when making applications by air. Between layby and 60 percent open bolls, the maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre.

Preharvest

Apply this product for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready Flex cotton after 60 percent boll crack. Up to 2 quarts of this product may be applied using either aerial or ground spray equipment. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of Roundup Ready Flex cotton. Do not apply this product over-the-top beyond first bloom to cotton grown for seed.

PRECAUTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY FLEX COTTON, HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

K. BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS.

DIRECTIONS FOR USE

NOTE: IF IMPROPERLY APPLIED, THIS PRODUCT HAS THE POTENTIAL TO CAUSE SEVERE CHRISTMAS TREE INJURY. FOLLOW ALL LABEL DIRECTIONS.

This product may be applied as a broadcast spray over established Christmas trees. Ensure that adequate buffers are maintained to prevent drift onto nearby desirable crops or vegetation. Read the entire "APPLICATION"

"EQUIPMENT AND TECHNIQUES" section of the GLYPHOSATE 41% SL label booklet for additional application precautions

This application is approved for the following Christmas tree species:

Douglas fir (*Pseudotsuga menziesii*)

Fir species (*Abies* spp.)

Spruce species (*Picea* spp.)

Applications may be made only after trees have completed at least a full growing season since planting or transplanting. Applications should not be made within 1 full year prior to tree harvest.

Applications may only be made in the fall after the formation of final conifer resting buds. Final resting buds must be fully hardened and in the dormant stage. Applications made at any other time may result in unacceptable Christmas tree injury.

Avoid spray pattern overlap, as injury may occur.

Apply 1 quart of this product per acre in 5 to 30 galls of water per acre.

NOTE: DO NOT ADD SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO THIS PRODUCT AS SEVERE CHRISTMAS TREE INJURY MAY RESULT.

This product may be used at rates from 1 to 2 quarts per acre in some areas. Consult your local Repar-Glypho, LLC representative or Glycel 41 % SL supplier for specific guidance if you require rates greater than 1 quart per acre.

Drift control additives may increase Christmas tree injury.

The use of other herbicides tank mixed with this product is not to be used because severe Christmas tree injury may result.

L. FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

GLYPHOSATE 41% SL Tolerant Tall Fescue Selections For Seed Protection

Use this product on [INSERT BRAND NAME] tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 4 to 16 fluid ounces per acre as a postemergence spray on **GLYPHOSATE 41% SL** tolerant tall fescue selections. See the label booklet for application instructions, rates, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the **GLYPHOSATE 41% SL** label booklet for a complete list of weeds controlled or suppressed.

NOTE: The rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with **GLYPHOSATE 41% SL** tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

Timing of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy and in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

M. GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the GLYPHOSATE 41% SL label booklet for rates. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quart per acre of this product may be used for improved results. Apply 10 quarts when weeds are dense.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any rates between 5 and 10 quarts of this product may be used in a tank mixture with these products.

<u>Tank-mix. Product*</u>	<u>Rate per Acre</u>
Arsenal TM *	0.5 to 4 pints
Barvel	1 to 4 pints
2, 4-D	0.5 to 1 pound
Garlon TM 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar TM X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar TM I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust TM	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike TM 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

*Follow all use directions for non-crops areas on this labeling of these potential tank mixture partners

- 1) Arsenal is not approved for use in the state of California.
- 2) Refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.
- 3) Read and carefully observe the label claims, cautionary statements, use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mixture products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Read "LIMITS OF WARRANTY AND LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

To the extent consistent with applicable law, when used in combination, the liability of Repar-Glypho, LLC shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Repar-Glypho, LLC product in such combination use.

TMArsenal is a trademark of American Cyanamid Company.

TMBanvel is a trademark of Sandoz Crop Protection Corporation.

TMGarlon and Spike are trademarks of Dow AgroSciences LLC.

TMHyvar, Krovar and Oust are trademarks of E. I DuPont de Nemours and Company.

N. FOR GROUND AND AERIAL APPLICATIONS TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY.
--

DIRECTIONS FOR USE

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Nonionic surfactants which are labeled for use with herbicides may be used to improve wetting of foliage. Do not reduce rates of GLYPHOSATE 41% SL when adding surfactant. Read and carefully observe surfactant rates, cautionary statements, and other information appearing on the surfactant label.

TIMING OF APPLICATION: Apply this product as a broadcast spray when plants are actively growing for partial control of undesirable vegetation listed on this label. Best results are obtained when application is made in the spring to early summer when brush species are at a high moisture content and flowering.

This product may be used for:

- Aid to burning treatment to establish and maintain fuel breaks
- Establishing fire perimeters and black lines
- Aid to prescribed burning
- Along fire roads and rights-of-way

APPLICATION RECOMMENDATION: Apply 2 quarts of this product per acre for partial control of the following emerged brush and chaparral species:

Ceanothus	Sage
Ceanothus spp.	Salvia spp.
Chamise	Scrub oak
Adenostroma fasciculatum	Quercus dumosa

Ground applications should be applied in 3 to 40 gallons of total spray solution per acre.

Aerial applications (helicopter only) should be applied in 3 to 15 gallons of total spray solution per acre.

Avoid direct application to any body of water.

AVOID DRIFT- DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

For aerial application of this product, please see the supplemental label directions for aerial application in California.

O. GLYPHOSATE 41% SL HERBICIDE FOR CONIFER RELEASE

DIRECTIONS FOR USE

AERIAL APPLICATION

This product may be applied using aerial spray equipment for conifer release treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of the label booklet for GLYPHOSATE 41% SL herbicide for information on how to properly spray this product by air.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

CONIFER RELEASE

For release, apply only where conifers have been established for more than one year. Vegetation should not be disturbed prior to treatment or until symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of the label booklet for GLYPHOSATE 41% SL herbicide.

For release of the following conifer species:

Douglas Fir	Pines*
<i>Pseudotsuga menziesii</i>	<i>Pinus</i> spp.
Fir	Spruce
<i>Abies</i> spp.	<i>Picea</i> spp.
Hemlockf	
<i>Tsuga</i> spp.	

*Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 1.5 to 2 quarts of this product per acre except in Washington and region, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of this product per acre before any major leaf drop of deciduous species.

For release of western hemlock apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly Pine	Slash Pine
<i>Pinus taeda</i>	<i>Pinus elliotti</i>
Eastern white pine	
<i>Pinus strobus</i>	

Late Season Application – Apply 1.5 to 2 quarts of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will result in potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later application. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label direction will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash	Oak, white
<i>Fraxinus spp.</i>	<i>Quercus alba</i>
Cherry, black	Persimmon
<i>Prunus serotina</i>	<i>Diospyros spp.</i>
Cherry, pin	Poplar, yellow
<i>Prunus pensylvanica</i>	<i>Liriodendron tulipifera</i>
Elm	Sassafras
<i>Ulmus spp.</i>	<i>Sassafras albidum</i>
Hawthorn	Sourwood
<i>Crataegus spp.</i>	<i>Oxydendrum arboreum</i>
Locust, black	Sumac, poison
<i>Robinia pseudoacacia</i>	<i>Rhus vernix</i>
Maple, red	Sumac, smooth
<i>Acer rubra</i>	<i>Rhus glabra</i>
Oak, black	Sumac, winged
<i>Quercus velutina</i>	<i>Rhus copallina</i>
Oak, post	Sweetgum
<i>Quercus stellata</i>	<i>Liquidambar styraciflua</i>
Oak, southern red	
<i>Quercus falcata</i>	

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

GLYPHOSATE 41% SL HERBICIDE PLUS OUST™ TANK MIXTURES FOR CONIFER RELEASE FROM HERBACEOUS WEEDS

To release loblolly pines from herbaceous weeds, tank mixtures of this product with Oust will provide control of annual weeds listed in the “WEEDS CONTROLLED” section of the label booklet for **GLYPHOSATE 41% SL** herbicide and the Oust label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of GLYPHOSATE 41% SL herbicide with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines.

This product plus Oust tank mixtures may not be applied by air in California.

This tank mixture may be applied using aerial equipment. When applying by air, use the selected rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use low rates of both products. Use higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	Johnsongrass**
<i>Paspalum notatum</i>	<i>Sorghum halepense</i>
Broomsedge	Poorjoe**
<i>Andropogon virginicus</i>	<i>Diodia teres</i>
Dock, curly	Trumpet creeper*
<i>Rumex crispus</i>	<i>Campsis radicans</i>
Dogfennel	Vaseygrass

<i>Eupatorium capilliflorum</i>	<i>Paspalum urvillei</i>
Fescue, tall	Vervain, blue
<i>Festuca arundinacea</i>	<i>Verbena hastata</i>
*Suppression at the higher rates only	
**Control at the higher rates	

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water, insects or disease.

Read and observe the cautionary statements and all other information appearing on the labels of herbicides used.

Oust is a trademark of E.I. DuPont de Nemours and Company.

MASTER LABEL FOR EPA REG. NO. 86004- U

Registered Brand Names:
GLYPHOSATE 41 % SL

Table of Contents for Master Label

	Main Label for Food Crop Uses	Page 2
	Main Label for Industrial, Turf, & Ornamental Uses	Page 69
	Supplemental Labeling	Page 99

See each label part for more detailed table of contents

I. MAIN LABEL FOR FOOD CROP USES

GLYPHOSATE 41% SL

Herbicide

Complete Directions for Use EPA Reg. No. 86004-

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Herbicide for Roundup Ready® Crops.

Selective broad-spectrum weed control in Roundup Ready crops. Non-selective, broad-spectrum weed control for many agricultural systems and farmsteads.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the entire label before using this product.

Use only according to label instructions.

[Container Label Statement (5 gallons or smaller):]

Do not use this container for residential treatments.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC REPACKAGING OR TOLL REPACKAGING AGREEMENT.

Non-refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

I. Main Label For Food Crop Uses

CONTENTS

1.	1.0	INGREDIENTS
2.	2.0	IMPORTANT PHONE NUMBERS
3.	3.0	PRECAUTIONARY STATEMENTS
	3.1	Hazards to Humans and Domestic Animals
	3.2	Environmental Hazards
	3.3	Physical or Chemical Hazards
4	4.0	STORAGE AND DISPOSAL
5	5.0	GENERAL INFORMATION (How This Product Works)
6	6.0	Weed Resistance Management
	6.1	General Weed Management Recommendations
	6.2	Management Recommendations for Glyphosate Resistant Biotype
7	7.0	MIXING
	7.1	Mixing with Water
	7.2	Tank Mixtures
	7.3	Tank Mixing Procedures
	7.4	Mixing for Hand-Held Sprayers
	7.5	Surfactants
	7.6	Ammonium Sulfate
	7.7	Colorants or Dyes
	7.8	Drift Reduction Additives
8	8.0	APPLICATION EQUIPMENT AND TECHNIQUES
	8.1	Aerial Equipment
	8.2	Ground Broadcast Equipment
	8.3	Hand-Held or Backpack Equipment
	8.4	Selective Equipment
	8.5	Injection Systems 8.6 CDA Equipment
9	9.0	ANNUAL AND PERENNIAL CROPS (Alphabetical)
	9.1	Cereal and Grain Crops
	9.2	Corn
	9.3	Cotton
	9.4	Fallow Systems
	9.5	Grain Sorghum (Milo)
	9.6	Herbs and Spices
	9.7	Oil Seed Crops
	9.8	Soybeans
	9.9	Sugarcane
	9.10	Vegetable Crops 9.11 Miscellaneous Crops
10	10.0	TREE, VINE, AND SHRUB CROPS (Alphabetical)
	10.1	Berry Crops
	10.2	Citrus
	10.3	Miscellaneous Tree Food Crops
	10.4	Non-Food Tree Crops

- 10.5 Pome Fruit
- I. Main Label For Food Crop Uses
 - 10.6 Stone Fruit
 - 10.7 Tree Nuts
 - 10.8 Tropical and Subtropical Trees and Fruits
 - 10.9 Vine Crops
- 11 11.0 **PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS**
 - 11.1 Alfalfa, Clover, and Other Forage Legumes
 - 11.2 Conservation Reserve Programs (CRP)
 - 11.3 Grass Seed or Sod Production
 - 11.4 Pastures
 - 11.5 Rangelands
- 12 12.0 **ROUNDUP READY CROPS**
 - 12.1 Roundup Ready Alfalfa
 - 12.2 Roundup Ready Canola (Spring Varieties)
 - 12.3 Roundup Ready Canola (Winter Varieties)
 - 12.4 Roundup Ready Corn 2
 - 12.5 Roundup Ready Gene Cotton
 - 12.6 Roundup Ready Flex Cotton
 - 12.7 Roundup Ready Gene Soybeans
 - 12.8 Roundup Ready Gene Sugar Beets
- 13 13.0 **NON-CROP USES AROUND THE FARMSTEAD**
 - 13.1 General Weed Control and Trim-And-Edge
 - 13.2 Greenhouse/Shade house
 - 13.3 Chemical Mowing
 - 13.4 Cut Stumps
 - 13.5 Habitat Management
- 14 14.0 **ANNUAL WEEDS RATE SECTION**
 - 14.1 Annual Weeds - Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K
 - 14.2 Annual Weeds - Hand-Held or Backpack Equipment
 - 14.3 Annual Weeds - Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems
- 15 15.0 **PERENNIAL WEEDS RATE SECTION**
- 16 16.0 **WOODY BRUSH AND TREES RATE SECTION**
- 17 17.0 **LIMIT OF WARRANTY AND LIABILITY**

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....41.0%

OTHER INGREDIENTS:.....59.0%
100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

No license granted under any non-U.S. patent(s).

I. Main Label For Food Crop Uses

2.0 IMPORTANT PHONE NUMBERS

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

I. MAIN LABEL FOR FOOD CROP USES

3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Repar-Glypho, LLC. Supplemental Labeling. Supplemental labeling may be found on the cdms.net or greenbook.net websites or obtained by contacting your authorized Repar-Glypho, LLC. retailer or Repar-Glypho, LLC. Company representative.

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

I. Main Label For Food Crop Uses

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. PESTICIDE STORAGE: Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

I. Main Label For Food Crop Uses

[Container Label Storage and Disposal Statements]:

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact Repar-Glypho, LLC. Company at 1-301-562-7330.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

Refiners: Do not reuse this container except for refill in accordance with a valid Repar-Glypho, LLC. Agreement for Repackaging or Toll Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Repar-Glypho, LLC. container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

(How This Product Works)

Product Description: This product is a postemergence, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

I. Main Label For Food Crop Uses

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds effects may not be visible for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" for specific weed recommendations.

Always use the higher product application rate within the recommended range when weed growth is heavy or dense or when weeds are growing in an undisturbed (non-cultivated) area.

Reduced weed control may result from treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by this herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates: The maximum application or use rates stated throughout this product's labeling are given in units of volume (fluid ounces, pints, or quarts) of this product per acre. However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as text mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of this product (6 pounds of glyphosate acid) per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product (8 pounds of glyphosate acid) per acre per year.

I. Main Label For Food Crop Uses

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

6.0 WEED RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE
-------	---	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using other cultural or mechanical practices.

6.1 General Weed Management Recommendations

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management recommendations:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Utilize the recommended label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Repar-Glypho, LLC representative, local retailer, or county extension agent.

6.2 Management Recommendations For Glyphosate Resistant Biotypes

Note: Contact your Repar-Glypho, LLC representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

I. Main Label For Food Crop Uses

Control recommendations for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling or fact sheets for this product and can be obtained from your local retailer or Repar-Glypho, LLC representative.

Since the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Repar-Glypho, LLC, is not responsible for any losses that may result from the failure of this product to control glyphosate resistant weeds biotypes.

The following good agronomic practices are recommended to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes:

7.0 MIXING

Spray solutions of this product should be mixed, stored and applied using only clean stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: PRODUCT PERFORMANCE MAY BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

7.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Begin filling the mixing tank or spray tank with clean water. Add the recommended amount of this product near the end of the filling process and mix gently. Use caution to avoid siphoning back into the carrier source. During mixing, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

7.2 Tank Mixtures This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and follow all label directions of all products in the tank mixture.

I. Main Label For Food Crop Uses

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC. for this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT recommended for applications of this product unless otherwise noted in this product label, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC..

When a tank mixture with a generic active ingredient, such as diuron, atrazine, 2,4-D or dicamba, is recommended in this label, the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank mixture.

Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture, and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Use according to the most restrictive label requirements.

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance. For best results, it is recommended that tank mixtures with this product be applied at a minimum spray volume rate of 10 gallons per acre.

7.3 Tank Mixing Procedure

Prepare tank mixtures of this product as follows:

1. Place a 20- to 35-mesh screen or wetting basket over the filling port of the tank.
2. Through the screen, fill the tank one-half full with water and start gentle agitation.
3. If ammonium sulfate is to be used, add it slowly through the screen into the tank, and continue adding water into the tank through the screen. If dry ammonium sulfate is being used, ensure that it is completely dissolved in the tank before adding other products.
4. If a wettable powder is used, make a slurry with the water, and add it SLOWLY through the screen into the tank while continuing gentle agitation.
5. If a flowable formulation is used, premix one part flowable with one part water, and add the diluted mixture SLOWLY through the screen into the tank while continuing gentle agitation.
6. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water, and add the diluted mixture slowly through the screen into the tank while continuing gentle agitation.
7. Continue filling the tank with water through the screen and add the required amount of this product near the end of the filling process.
8. If a nonionic surfactant is used, add it to the tank before completing the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquids (this product) followed by surfactant.

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying.

Keep by-pass and return lines on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

7.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product as indicated in the following table in water:

I. Main Label For Food Crop Uses

Spray Solution

Amount of **GLYPHOSATE 41 % SL**

Desired Volume	0.5%	1%	1.5%	2%	5%	10%
1 gal	0.7 oz.	1.3 oz.	2 oz.	2.7 oz.	6.5 oz	13 oz.
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in backpack sprayers, it is recommended that the appropriate amount of this product be mixed with water in a larger container and then fill the sprayer with the mixed solution.

7.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

7.6 Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates recommended in this label. Lower rates will result in reduced performance.

7.7 Colorants or Dyes

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce performance. Use colorants or dyes according to the manufacturer's recommendations.

7.8 Drift Reduction Additives

Drift reduction additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet Applicator (CDA) equipment. When a drift reduction additive is used, read and carefully observe the precautions, limitations, and all other information appearing on the additive label. Use of drift reduction additives can affect spray coverage, which can reduce product performance.

8.0 APPLICATION EQUIPMENT AND TECHNIQUES Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment: Aerial—Fixed wing and helicopter.

Ground Broadcast Spray--Boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held or Backpack Spray Equipment--Backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment-- Shielded and hooded sprayers, wiper applicators and sponge bars.

I. Main Label For Food Crop Uses

Injection Systems--Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA)---Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING THE DESIRED VOLUMES.

8.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED IN THIS LABEL.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling. Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 2 quart per acre. Refer to the individual use area sections of this label for further instructions.

FOR AERIAL APPLICATION IN ARKANSAS OR CALIFORNIA, OR SPECIFIC COUNTIES THEREIN, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE OR COUNTY FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

This product, when tank mixed with dicamba, may not be applied by air in California.

Ensure uniform application - to avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target movement during aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Comply with all State regulations where applicable.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (read the following sections on "Wind", "Temperature and Humidity", and "Temperature Inversions").

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing spray pressure.
- **Number of nozzles:** Use the minimum number of nozzles that will provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

I. Main Label For Food Crop Uses

- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind speed, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke produced by a ground source or ^{an} aircraft smoke generator can also identify temperature inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water. **Aircraft Maintenance**

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may help prevent corrosion.

I. Main Label For Food Crop Uses

8.2 Ground Broadcast Equipment

Apply the product rates recommended in this label in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid generating a fine mist. For best results with ground application equipment, use flat spray nozzles. Check spray pattern for uniform distribution.

8.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled on a spray-to-wet basis; do not spray to the point of run-off. Spray coverage should be uniform and complete. Use coarse sprays only. For recommended rates and timing, refer to the "ANNUAL WEEDS HAND-HELD OR BACKPACK EQUIPMENT" section of this product label.

8.4 Selective Equipment

This product may be diluted in water and applied through shielded sprayers, hooded sprayers, wiper applicators or sponge bars to weeds growing in any non-crop site specified on this label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants), and wipers may be used over-the-top of crops only when specifically recommended in this product's labeling. Such equipment must be capable of preventing all crop contact with the herbicide solutions and operated without leakage of spray mists or dripping onto crop.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of this product with desirable vegetation may result in unwanted plant damage or destruction. Crop injury may occur when the foliage of treated weeds comes into contact with leaves of the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Sprayers

This product, when applied at recommended rates under the conditions described in the following paragraphs for shielded and hooded sprayers, will control those weeds listed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Adjust the shields on these sprayers to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is capable of completely enclosing the spray pattern. If necessary, extend the front and rear flaps of the hooded applicator downward to reach the ground in deep furrows. **EXTREME CARE MUST BE TAKEN TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be configured and operated in a manner that minimizes bouncing and avoids raising the hoods up off the ground at any time. If the hood is raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operating this equipment on rough or sloping terrain where the spray hoods might be raised up off the ground surface.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure low-drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is recommended. Spray volume should be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- Spray hoods must be operated on the ground or skimming across the ground surface.

1. Main Label For Food Crop Uses

- Leave at least an 8-inch untreated strip over the drill row. (For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.)
- Operate at ground speeds of no greater than 5 miles per hour to avoid bouncing of the spray hoods.
- Apply when wind speeds are 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Wiper Applicators

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators: Use solutions ranging from 33 to 75 percent of this product in water. **For**

Panel Applicators: Use solutions ranging from 33 to 100 percent of this product in water.

8.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products for use in injection systems.

8.6 CDA Equipment

The rate of this product applied per acre by vehicle-mounted controlled droplet applicator (CDA) equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply in 2 to 15 gallons of water per acre.

For control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of

I. Main Label For Food Crop Uses

perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet applicators produce a spray pattern that is not easily visible. Extreme care must be taken to avoid spray or drift onto the foliage or any other green tissue of desirable vegetation, as damage or destruction of the plant may result.

9.0 ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the "ROUNDUP READY CROPS" section of this label or separately published Repar-Glypho, LLC, Supplemental Labeling for this product for instructions on treating Roundup Ready crops.

TYPES OF APPLICATIONS:

Chemical Fallow, Preplant Fallow Beds, Preplant, At-Planting, Preemergence, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest.

GENERAL USE INSTRUCTIONS:

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", AND WOODY BRUSH AND TREES RATE SECTIONS" in this label. Recommended rates of this product that are emphasized in this product's labeling to control tough weeds take precedence over the general rates in the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS". Repeat applications may be made up to a maximum of 8 quarts of this product per acre per year.

Post-directed hooded sprayers and wiper applicators capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted in the individual crop sections that follow, wipers may also be used above certain crops to control tall weeds. Refer to the "SELECTIVE EQUIPMENT" section of this label for essential precautions regarding crop injury. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified in this product's labeling. Refer to the "AERIAL EQUIPMENT" section of this label for additional information.

GENERAL PRECAUTIONS, RESTRICTIONS:

Avoid contact of this herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops as severe crop injury or destruction may result. When making at-planting and preemergence applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death of emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any crops not listed on this label. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

I. Main Label For Food Crop Uses

For broadcast postemergence treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

9.1 Cereal and Grain Crops

LABELED CROPS: Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Rye, Quinoa, Tef, Teosinte, Triticale, Wheat (all types), Wild rice.

PRECAUTIONS, RESTRICTIONS: Do not treat rice fields or levees when flooded.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Red Rice Control Prior to Planting Rice, Spot Treatment (Except Rice), Over-The-Top Wiper Application (Feed Barley and Wheat only), Preharvest (Wheat And Feed Barley Only).

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting of cereal crops. Applications must be made prior to crop emergence.

TANK MIXTURES: In wheat, a tank mix with Aim may be used.

Red Rice Control Prior to Planting Rice

USE INSTRUCTIONS: Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are at the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

PRECAUTIONS, RESTRICTIONS: Avoid spraying during conditions of low humidity, as reduced control may result. Do not treat rice fields or levees when the fields contain floodwater. Do not flood treated fields for 8 days following application.

Spot Treatment (Except Rice)

USE INSTRUCTIONS: This product may be applied as a spot treatment in cereal crops. Apply this product before heading in small grains.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Over-the-Top Wiper Application (Feed Barley and Wheat only)

USE INSTRUCTIONS: Wiper applications may be used in wheat and feed barley. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth.

PRECAUTIONS, RESTRICTIONS: Allow at least 35 days between application and harvest. Do not use roller applicators. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Preharvest (Feed Barley And Wheat Only)

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of feed barley or wheat. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard-dough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest.

For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1 quart of this product per acre. Allow 7 days between application and harvest or grazing. Preharvest application is not recommended for barley or wheat grown for seed, as a reduction in germination or vigor may occur.

1. Main Label For Food Crop Uses

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of cereal crops. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.2 Corn

TYPES OF CORN: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest.

Use directions for corn containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting corn. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with the following products to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Ensure that the specific product is registered for application prior to planting corn. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

2,4-D	Distinct™	Leadoff™
Airm®	Dual Magnum™	Alachlor
Atrazine	Dual II Magnum™	Linex™/Lorox™
Axiom™	Epic™	Keystone LA
Balance™ / Balance PRO	Frontier™/Outlook™	Marksman™
Banvel™/Clarity™	Fultime™	Micro-Tech®
Bicep Magnum™	Guardsman™/Guardsman MAX	Prowl™
Bicep II Magnum™	Harness®	Python™
Bullet®	Harness Xtra	Resource®
Degree®	Harness Xtra 5.6l	Simazine Surpass
Degree Xtra®	Hornet	Topnotch™
	Lariat®	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

PRECAUTIONS, RESTRICTIONS: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. The area covered by this recommendation includes Illinois and Indiana south of Route 50, and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

I. Main Label For Food Crop Uses

Hooded Sprayers

USE INSTRUCTIONS: This product may be applied with hooded sprayers for weed control between rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instructions for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Corn must be at least 12 inches tall, measured without extending leaves. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. Do not apply more than 1 quart of this product per acre for each application and no more than 3 quarts per acre per year for hooded sprayer applications.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment prior to Bilking of corn.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

USE INSTRUCTIONS: Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 2 quarts of this product per acre.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Preharvest application is not recommended for corn grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required to control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.3 Cotton

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Selective equipment, Spot treatment, Preharvest.

Use directions for cotton containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with the following products provided that the specific product being used is registered for application prior to planting cotton. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Caparol®	Direx	Prowl
Clarity™	Dual Magnum™	Prowl H2O
Command	Dual II Magnum™	Staple®
Cotoran®	Karmex	Stalwart
Cotton PRO®	Meturon®	Zorial®

PRECAUTIONS, RESTRICTIONS: For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

I. Main Label For Food Crop Uses

Selective Equipment

USE INSTRUCTIONS: This product may be applied through hooded sprayers, shielded sprayers or wiper applicators in cotton. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Allow at least 7 days between application and harvest.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment in cotton prior to boll opening.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

USE INSTRUCTIONS: This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "ANNUAL WEEDS", and "PERENNIAL WEEDS RATE SECTIONS" of this label. For cotton regrowth inhibition, apply 1 pint to 2 quarts of this product per acre.

Up to 2 quarts of this product may be applied after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

TANK MIXTURES: This product may be tank mixed with DEFTTM 6, FolexTM, Ginstar, or PrepTM to provide additional enhancement of cotton leaf drop.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Preharvest application is not recommended for cotton grown for seed, as a reduction in germination or vigor may occur. THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF THIS PRODUCT TO COTTON IS PROHIBITED. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

9.4 Fallow Systems

LABELED CROPS: This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label; for non-listed crops, applications must be made at least 30 days prior to planting.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Aid-to-Tillage, Chemical Fallow

USE INSTRUCTIONS: This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Tank mixtures of this product with 2,4-D and dicamba may be used for a broader weed control spectrum. Applications up to 2 quarts per acre may be made by aerial application onto fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

PRECAUTIONS, RESTRICTIONS: Do not apply dicamba tank mixtures by air in California. Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

Preplant Fallow Beds

USE INSTRUCTIONS: This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label prior to planting.

TANK MIXTURES: Use 12 fluid ounces of this product, plus 2 to 3 fluid ounces of GoalTM 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 3" -- common cheeseweed, chickweed, groundsel; 6" -- London rocket, shepherd's-purse.

Use 16 fluid ounces of this product, plus 2 to 3 fluid ounces of GoalTM 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 6" -- common cheeseweed, groundsel, mare's tail (*Coryza canadensis*), 12" -- chickweed, London rocket, shepherd's-purse.

I. Main Label For Food Crop Uses

Aid-to-Tillage

USE INSTRUCTIONS: This product may be used in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 12 fluid ounces of this product in 3 to 10 gallons of water per acre before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

PRECAUTIONS, RESTRICTIONS: Tank mixtures with residual herbicides may result in reduced product performance.

9.5 Grain Sorghum (Milo)

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Over-the-Top Wiper Application, Preharvest.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in tank-mixture before, during or after planting grain sorghum. Applications must be made prior to crop emergence.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, provided that the specific product being used is labeled for application prior to planting grain sorghum.

Atrazine	IntRRo
Bicep II Magnum	Lariat
Bullet	Micro-Tech
Dual II Magnum	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply a tank mixture of this product at 2 pints per acre plus the products listed above. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased for acceptable weed control.

Spot Treatment, Over-the-Top Wiper Application

USE INSTRUCTIONS: This product may be applied as a spot treatment in grain sorghum before heading. This product may be applied with wiper applicators to control or suppress tall weeds. For additional instructions, see "Wiper Applicators" in the "SELECTIVE EQUIPMENT" section of this label.

PRECAUTIONS, RESTRICTIONS: For spot treatment, do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction. Allow at least 40 days between wiper application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

Hooded Sprayers

USE INSTRUCTIONS: This product may be applied with hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Milo must be at least 12 inches tall, measured without extending leaves. Treat before mile, sends tillers between the drill rows. If tillers are sprayed, the main plant may be damaged or destroyed. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage, which shall be the sole responsibility of the applicator. Do not graze or feed milo forage or fodder following hooded sprayer applications. Do not apply more than 1 quart of this product per acre per application and no more than 3 quarts per acre per year for hooded sprayer applications.

I. Main Label For Food Crop Uses

Preharvest

USE INSTRUCTIONS: This product may be applied for weed control prior to harvest after sorghum grain has reached 30 percent grain moisture or less.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 2 quarts of this product per acre. As with other herbicides that cause sudden plant death, avoid preharvest applications of this product to milo infected with charcoal rot as lodging can occur. Allow a minimum of 7 days between application and harvest of grain sorghum. Preharvest application is not recommended for sorghum grown for seed, as a reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of grain sorghum. Higher rates may be required to control large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product per acre for control, or 1.5 pints of this product per acre for suppression.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.6 Herbs and Spices

LABELED CROPS: Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Cilantro (leaf), Cilantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Over-the-Top Wiper Application (Peppermint and Spearmint Only), Spot Treatment (Peppermint and Spearmint Only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Over-the-Top Wiper Application, Spot Treatment (Peppermint and Spearmint only)

USE INSTRUCTIONS: This product may be applied as a spot treatment or over the top of peppermint or spearmint with wiper applicators in spearmint and peppermint. Apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution to a limited area.

PRECAUTIONS, RESTRICTIONS: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested should be treated at one time. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. For wiper applications, contact of the herbicide solution with the crop may result in discoloration, stunting, or destruction.

I. Main Label For Food Crop Uses

9.7 Oil Seed Crops

LABELED CROPS: Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower.

Use directions for canola containing a glyphosate tolerant gene, are in the "ROUNDUP READY CROPS" section of this label.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

USE INSTRUCTIONS: This product may be applied before, during or after planting oil seed crops listed in this section. Broadcast applications must be made prior to crop emergence. Wiper applicators or hooded sprayers may be used between the rows once the crop is established. See additional instructions of the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl may be applied before, during or after planting into conventionally tilled soil, a cover crop, established sod or previous crop residue.

PRECAUTIONS, RESTRICTIONS: For use with canola, do not apply more than 2 quarts of this product per acre. For use with sunflowers, do not apply more than 1 quart of this product per acre as a single preplant or preemergence application per year. Do not feed or graze sunflower forage following application of this product.

9.8 Soybeans

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest, Selective Equipment.

Use directions for soybeans containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting soybeans. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with the following products to provide residual weed control, a broader spectrum, or an alternate mode of action. Ensure that the specific product being used in the tank mixture is registered for application prior to planting soybean, and follow all label directions of all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Aim™	Domain™	Micro-Tech
Assure II™	Dual Magnum	Prowl
Authority™	Dual II Magnum	Pursuit™
Axiom	Firststate™	Pursuit Plus
Blanket	Flexstar™	Python
Boundary	Frontier™/Outlook™	Reflex™
Canopy™	Fusion™	Resource™
Canopy EXT™	Gauntlet™	Scepter™
Classic	Gangster	Sencor™/Lexone™
Command™	IntRRo	Squadron™
Command Xtra™	Linex™	Steel™
2,4-D	Lorox/Linuron	Valor™
	Lorox Plus™	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when

I. Main Label For Food Crop Uses

weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment prior to initial pod set in soybeans.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction.

Preharvest

USE INSTRUCTIONS: This product may be applied for weed control prior to harvest of soybeans after pods have set and lost all green color. Apply at rates given in the "ANNUAL WEEDSRATE SECTION" and "PERENNIAL WEEDSRATE SECTION". Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 5 quarts of this product per acre for preharvest applications. Do not apply more than 2 quarts of this product per acre by air. Allow a minimum of 7 days between application and harvest of soybeans. Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last preharvest application. (If the application rate is 1 quart per acre or lower, the grazing restriction is reduced to 14 days after last preharvest application.) Preharvest application is not recommended for soybeans grown for seed, as a reduction in germination or vigor may occur.

Selective Equipment

USE INSTRUCTIONS: This product may be applied through shielded or hooded sprayers, wiper applicators or sponge bars in soybeans. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Allow at least 7 days between application and harvest.

9.9 Sugarcane

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus Spot Treatment. Sugarcane Ripening.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied in or around sugarcane fields, or in fields prior to the emergence of plant cane.

PRECAUTIONS, RESTRICTIONS: Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using spray-to-wet or a spray-to-wet technique. Volunteer or diseased sugarcane should have at least 7 new leaves.

PRECAUTIONS, RESTRICTIONS: Avoid spray contact with healthy sugarcane plants since severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

Fallow Treatments

USE INSTRUCTIONS: This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Applications up to 3 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used.

I. Main Label For Food Crop Uses

Hooded Sprayers

USE INSTRUCTIONS: This product may be applied using hooded sprayers for weed control between the rows of sugarcane. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional use instructions.

PRECAUTIONS, RESTRICTIONS: Crop injury may occur when the foliage of treated weeds contact the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Sugarcane Ripening

USE INSTRUCTIONS: This product may be applied to foliage as a plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane. It is effective in both low and high-tonnage sugarcane.

When applied as directed under the conditions described, this product will hasten ripening and extend the period of high sucrose level in sugarcane.

As a result of leaf desiccation, improved trash burn can be expected.

Most of the sucrose increase is concentrated in the top nodes of the treated sugarcane stalk. In order to maximize sugar where topping is practiced during harvest, top at the base of the fourth leaf.

APPLICATION RATES:

Use the following application rates and timing instructions according to the State in which the sugarcane is grown.

NOTE: Use the higher rate within the recommended range when treating sugarcane under adverse ripening conditions or when less responsive varieties are to be treated.

FLORIDA— Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of LAST PLATOON CANE ONLY.

HAWAII— Apply 12 to 28 fluid ounces of this product per acre 4 to 10 weeks before harvest.

LOUISIANA— Apply 5 to 16 fluid ounces of this product per acre 3 to 7 weeks before harvest of PLATOON CANE ONLY.

PUERTO RICO— Apply 7 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

TEXAS— Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

PRECAUTIONS, RESTRICTIONS: Prior to application, consult your state sugarcane authority or local Repar-Glypho, LLC, representative regarding the degree of sucrose response anticipated from the variety of sugarcane to be treated.

Application of this product may initiate development of shooting eyes. This product may not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2 to 3 weeks after application, this product may produce a slight yellowing to pronounce browning and drying of leaves, and a shortening of upper internodes. Spindle death may occur.

Rainfall within 6 hours after application may reduce effectiveness.

Application is not recommended for sugarcane grown for seed, as a reduction in germination or vigor may occur.

Do not feed or graze treated sugarcane forage following application. Do not plant subsequent crops in treated fields other than the following for 30 days after application: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybeans, squash (all types) or wheat.

Do not apply for enhanced ripening to any crops other than sugarcane.

I. Main Label For Food Crop Uses

9.10 Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL VEGETABLE CROPS LISTED ALPHABETICALLY IN THIS SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Chemical Fallow, Preplan Fallow Beds, Preplan, At-Planting, Preemergence, Prior to Transplanting Vegetables, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest, Selective Equipment Applications (Nonbearing Ginseng), Over-the-Top Wiper Application (Rutabagas only), Spot Treatment or Preharvest (Dry Beans, Peas, Lentils, and Chickpeas only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

9.10.1 Brassica Vegetables

LABELED CROPS: Broccoli, Chinese broccoli (gai lan), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

9.10.2 Bulb Vegetables

LABELED CROPS: Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

9.10.3 Cucurbit Vegetables and Fruits

LABELED CROPS: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Melons (all), *Momordica spp* (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon.

PRECAUTIONS, RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

I. Main Label For Food Crop Uses

9.10.4 Leafy Vegetables

LABELED CROPS: Amaranth (Chinese spinach), Arugula Croquette), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

PRECAUTIONS, RESTRICTIONS: For Watercress, avoid applications within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of crop injury.

9.10.5 Fruiting Vegetables

LABELED CROPS: Eggplant, Groundcherry (*Physalis spp.*), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.

PRECAUTIONS, RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato and tomatillos grown on sandy soil, do not make hooded, shielded sprayer or wiper applications in row middles because of potential for crop injury.

9.10.6 Legume Vegetables (Succulent or Dried)

LABELED CROPS: Bean (*Lupinus*; includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*; includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*; includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*; includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

Spot Treatment, or Preharvest (Dry Beans, Peas, Lentils and Chickpeas only)

USE INSTRUCTIONS: This product may be applied as an over-the-top broadcast spray or as a spot treatment to control labeled weeds in dry beans, peas, lentils or chick peas. For spot treatment, to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed, apply in 10 to 20 gallons of water per acre through ground broadcast spray equipment or use a 2 percent solution in a hand-held sprayer. For preharvest treatments, apply in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less).

PRECAUTIONS, RESTRICTIONS: Follow the limitations listed in the table below. Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not specified for treatment in this label. Do not feed treated vines and hay from these crops to livestock. Do not treat cowpeas or field (feed) peas, since these crop are considered to be grown as livestock feed. Preharvest application is not recommended for dry legumes grown for seed, as a reduction in germination or vigor may occur.

Crop	Maximum Rate (fluid ounces per Acre)	Minimum Pre- Harvest Interval (days)	Allowed in
Dry Beans	32	7	All States
Dry Peas, Lentils, and Chickpeas	26	14	Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, and Washington

I. Main Label For Food Crop Uses

LABELED CROPS: Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

Selective Equipment Applications (Non-Bearing Ginseng only)

USE INSTRUCTIONS: This product may be applied for general weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and backpack wands, lances, and orchard guns or with wiper application equipment. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Conduct applications so that there is no contact of this product with the ginseng plant. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Applications must be made at least one year prior to harvest.

Over-the-Top Wiper Application (Rutabagas only)

USE INSTRUCTIONS: Wiper applicators may be used over the top of rutabagas for the control of taller weeds. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

9.11 Miscellaneous Crops

LABELED CROPS: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: General Weed Control, Site Preparation, Spot Treatment (Asparagus).

Use directions for sugar beets containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

PRECAUTIONS, RESTRICTIONS: Avoid contact of this product with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any crop not listed on this label. See additional information in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

General Weed Control, Site Preparation

USE INSTRUCTIONS: This product may be applied for general weed control or for site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove residues of this product from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Allow at least 21 days between residue removal and transplanting. Applications made at emergence will result in injury or death to emerged seedlings.

I. Main Label For Food Crop Uses

Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.

Spot Treatment (Asparagus)

USE INSTRUCTIONS: This product may be applied immediately after cutting, but prior to the emergence of new spears.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Post-Harvest (Asparagus)

USE INSTRUCTIONS: This product may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Direct contact of the spray with asparagus may result in serious crop injury.

10.0 TREE, VINE, AND SHRUB CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL TREE, VINE, AND SHRUB CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Preplant (site preparation) Broadcast Sprays, General Weed Control, Middles (between rows of trees, vines or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (shielded sprayers, wiper treatments), Directed Spray, Spot Treatment, Perennial Grass Suppression, Cut Stump.

This product may be applied with boom equipment, CDA equipment, shielded sprayers, hand-held and backpack wands, lances, orchard guns or with wiper applicator equipment, unless prohibited in the specific Crop sections that follow.

GENERAL USE INSTRUCTIONS:

This product may be applied in middles (between rows of trees or vines), strips (within rows of trees or vines), and for general weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 1 pint to 5 quarts per acre according to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label. Use the higher rates in the recommended range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts per acre per year. See the "GENERAL INFORMATION" section of this label for more information on Maximum Annual Use Rates.

GENERAL PRECAUTIONS, RESTRICTIONS:

Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used in order to minimize the potential for overspray or drift of this product onto the crop. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with the crop may be used. See additional instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

I. Main Label For Food Crop Uses

Allow a minimum of 3 days between application and transplanting.

Middles (between rows)

USE INSTRUCTIONS: This product will control or suppress annual and perennial weeds and ground covers growing between rows of tree and vine crops listed on this label. If weeds are under drought stress, irrigate prior to application. Reduced weed control may result if weeds have been recently mowed at the time of application.

TANK MIXTURES: A tank mixture of this product plus Goal 2XL may be applied for annual weed control between rows (middles) of citrus crops, tree fruits, tree nuts and vine crops. This mixture is recommended when weeds are stressed or growing in dense populations. Application of 16 to 32 fluid ounces of this product per acre plus 3 to 12 fluid ounces of Goal 2XL per acre will control annual weeds with a maximum height, length or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). This tank mixture will also control common cheeseweed (malva) or hairy fleabane (*Conyza bonariensis*) with a maximum height, length or diameter of 3 inches. Read and follow all labels and directions for use of all products being used in the tank mixture.

Strips (In Rows)

TANK MIXTURES: This product may be applied within rows of tree or vine crops in tank mixtures with the following products, provided that the label of the specific product used permits the desired use:

Devrinol™ 50 DF	Simazine 4L
Direx™ 4L	Simazine 80W
Goal 2XL	Sim-Trol™ 4L
Karmex DF	Solican™ DF
Krovarl	Surflan™ AS
Prowl	Surflan 75w
Princep Caliber™90	

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for information about specific crops, rates, geographic restrictions and precautionary statements. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Perennial Grass Suppression

This product will suppress perennial grasses such as bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

I. Main Label For Food Crop Uses

For burndown of Bermudagrass, apply 1 to 2 quarts of this product in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 6 to 16 fluid ounces of this product per acre east of the Rocky Mountains and 16 fluid ounces of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the Bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces of this product per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

Cut Stump (tree crops)

USE INSTRUCTIONS: Cut stump applications of this product may be made during site preparation or site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

Citrus Trees: Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo (Ugli), Tangor.

Fruit Trees: Apple, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince.

Nut Trees: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).

Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

PRECAUTIONS, RESTRICTIONS: DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

10.1 Berry Crops

LABELED CROPS: Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Salal, [Optional Crop: Raspberry (black, red)].

TYPES OF APPLICATIONS: Those listed in Section 10.0, plus Spot Treatment in Cranberry Production and Post-Harvest Treatment in Cranberry Production.

PRECAUTIONS, RESTRICTIONS: To avoid damage, herbicide spray must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest of cranberries. Allow a minimum of 14 days between last application and harvest of other berry crops.

Spot Treatment in Cranberry Production

USE INSTRUCTIONS: Spot treatments may be used to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in this label may be used. Drop water level to

I. Main Label For Food Crop Uses

remove standing water in ditches prior to application. With hand-held sprayers, use 1- to 2-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off.

PRECAUTIONS, RESTRICTIONS: To achieve maximum weed control in treatments of dry ditches after water draw down, allow 2 or more days after treatment before reintroduction of water. Apply this product within 1 day after draw down to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not make applications by air. Do not apply directly to water. Use nozzles that emit medium- to large-sized droplets in order to minimize spray drift and avoid crop injury.

Post-Harvest Treatments in Cranberry Production

USE INSTRUCTIONS: Application of this product may be made after the harvest of cranberries to control weeds growing in the field. Apply this product to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label may be used. If using hand-held sprayers, use a 0.5- to 1-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off. If using hand-held boom sprayers, apply 2 to 4 quarts of this product per acre.

PRECAUTIONS, RESTRICTIONS: Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not make applications by air. Do not apply directly to water. Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

10.2 Citrus

LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugh), Tangor.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only.

USE INSTRUCTIONS:

The following use instructions pertain to applications in Florida and Texas only: For burndown or control of the weeds listed below, apply the recommended rates of this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 2 to 3 quarts of this product in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall, and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, use of this product in a tank mixture with KrovarTM I or KarmexTM may improve weed control. Refer to the individual product labels for treatment information on specific crops, rates, geographic restrictions and precautionary statements.

Perennial weeds:

S = Suppression

B = Burndow

PC = Partial control

C = Control

WEED SPECIES	GLYPHOSATE 41% SL RATE PER ACRE			
	1 QT	2 QT	3 QT	5 QT
Bermudagrass	B	---	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	---	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	---	PC	C

I. Main Label For Food Crop Uses

10.3 Miscellaneous Tree Food Crops

LABELED CROPS: Cactus (fruit and pads), Palm (heart, leaves), Palm (oil).

TYPES OF APPLICATIONS: Those listed in Section 10.0.

10.4 Non-Food Tree Crops

LABELED CROPS: Pine, Poplar, Eucalyptus, Christmas trees, All other non-food tree crops.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

NOTE: unless otherwise directed, this product is not recommended for use as an over-the-top broadcast spray in plantations or other labeled tree crops.

Directed Spray, Spot treatment, Wiper Application

USE INSTRUCTIONS: This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

PRECAUTIONS, RESTRICTIONS: Avoid contact of spray, drift or mist of this product with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Site preparation

USE INSTRUCTIONS: This product may be used for weed control prior to planting non-food tree crops.

PRECAUTIONS, RESTRICTIONS: Precautions should be taken to protect non-target plants during site preparation applications.

10.5 Pome Fruit

LABELED CROPS: Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in pome crops.

10.6 Stone Fruit

LABELED CROPS: Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), Plumcot.

TYPES OF APPLICATIONS: Those listed in Section 10.0. For olive groves, apply only as a directed spray.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 17 days between last application and harvest in stone fruit crops.

Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. **USE EXTREME CARE TO ENSURE THAT NO PART OF THE PEACH TREE IS CONTACTED WITH OVERSPRAY OR DRIFT OF THIS PRODUCT.**

10.7 Tree Nuts

LABELED CROPS: Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (black, English)

TYPES OF APPLICATIONS: Those listed in Section 10.0.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut.

10.8 Tropical and Subtropical Trees and Fruits

LABELED CROPS: Ambarella, Atemoya, Avocado, Banana, Barbados cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates,

I. Main Label For Food Crop Uses

Durian, Feijoa, Figs, Governor's plum, Guava, Ilaça, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genie), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots and leaves), Wax jambe.

TYPES OF APPLICATIONS: Those listed in Section 10.0, and as described below as a Bananacide (Bananas Only).

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit listed. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

Bananacide (Banana Only)

USE INSTRUCTIONS: This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus, as well as non-infected banana plants in order to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1/25 fluid ounce (1 mL) of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) within a 4-foot radius around a treated mat should be mechanically destroyed.

For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1/2 fluid ounce (15 mL) of this product per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for general weed control.

10.9 Vine Crops

LABELED CROPS: Grapes (raisin, table, wine) Hops, Kiwi, Passion fruit.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

Apply this product for weed control only when green shoots, canes or foliage are not in the spray zone.

In the northeast and Great Lakes regions, apply this production grape vineyards prior to the end of the bloom stage of grapes in order to avoid crop injury, or apply using shielded sprayers or wiper equipment.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 14 days between last application and harvest of vine crops. Do not use selective equipment in kiwi.

11.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELANDS

GENERAL USE INSTRUCTIONS: Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label for rate recommendations for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. Recommended rates of this product specified on this label for the control of tough weeds, or those specified on separate supplemental labeling for this product, supersede rates recommended in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label. Additional information on the control of tough weeds can be found in Fact Sheets published by Repar-Glypho, LLC. for this product.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling.

I. Main Label For Food Crop Uses

11.1 Alfalfa, Clover, and Other Forage Legumes

LABELED CROPS: Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types).

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Preharvest (except Kenaf and Leucaena), Spot Treatment, Over-the-Top Wiper Application, Renovation, Stand Removal.

Use directions for alfalfa with a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting crops listed in this section. Applications must be made prior to crop emergence.

PRECAUTIONS, RESTRICTIONS: Remove domestic livestock before application. The crop may be fed or grazed as soon as it reaches sufficient maturity.

Preharvest (Except Kenaf and Leucaena)

USE INSTRUCTIONS: This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This product will control annual or perennial weeds, including quackgrass, when applied prior to crop harvest. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

PRECAUTIONS, RESTRICTIONS: Make only one application to an existing stand of crop per year. The treated crop and weeds can be harvested and fed to livestock according to the application rates and intervals defined below.

	Maximum single application rate (per acre)	Minimum interval between application and harvest or livestock grazing
Alfalfa	2 quarts	36 hours
All other legumes listed	3 pints	3 days

This application may destroy the alfalfa stand and may severely injure or destroy other labeled crops such as clover. Preharvest application is not recommended for alfalfa grown for seed, as a reduction in germination or vigor may result.

Spot Treatment, Over-the-Top Wiper Application

USE INSTRUCTIONS: This product may be applied as a spot treatment or over the top of crops listed in this section with wiper applicators. For wipers, see "Wiper Applicators" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label. Applications may be made in the same area at 30-day intervals.

PRECAUTIONS, RESTRICTIONS: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field area should be treated at one time. Remove domestic livestock before application and wait 3 days after application before grazing livestock or harvesting.

Renovation, Stand Removal

USE INSTRUCTIONS: This product may be applied as a broadcast spray to remove established stands of alfalfa, clover, and other forage legumes listed on this label. If the crop is to be grazed or harvested for feed, use up to a maximum of 2 quarts per acre in alfalfa and up to 3 pints per acre in other labeled legumes. For complete removal of established stands of clover, it may be necessary to use treatment rates greater than this, as listed in the "PERENNIAL WEEDS RATE SECTION" of this label.

I. Main Label For Food Crop Uses

PRECAUTIONS, RESTRICTIONS: For applications up to 2 quarts per acre for alfalfa or 3 pints per acre for all other forage legumes, remove domestic livestock before application, follow the minimum harvest or grazing intervals in the table above under Preharvest in this section. For treatment rates above these levels, do not graze or harvest treated foliage for livestock feed or allow grazing. Crops listed in this label may be planted into the treated area at any time; for all other crops, wait 30 days or more between application and planting.

11.2 Conservation Reserve Program (CRP)

TYPES OF APPLICATIONS: Renovation (Rotating Out Of CRP), Site Preparation, Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application.

Renovation (Rotating Out of CRP), Site preparation

USE INSTRUCTIONS: This product may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation recommendations.

PRECAUTIONS, RESTRICTIONS: Crops listed on this label may be planted into the treated area at any time; for any other crop, wait at least 30 days between application and planting.

Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application

USE INSTRUCTIONS: Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land. Applications may be made using wiper applicators to control tall weeds or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: *Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant.* No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

10.3 Grass Seed or Sod Production

LABELED CROPS: Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation, Shielded Sprayers, Over-the-Top Wiper Application, Spot Treatment, Creating Rows in Annual Ryegrass.

Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation

USE INSTRUCTIONS: This product controls most existing vegetation prior to renovating turf or forage grass seed areas or establishing turf grass grown for sod. It may be used to destroy undesirable grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting or for renovation. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turf management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates of up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed for treatment in this label, applications must be made at least 30 days prior to planting. Applications must be made prior to the crop emergence in order to avoid crop injury.

I. Main Label For Food Crop Uses

Shielded Sprayers

USE INSTRUCTIONS: Apply 1 to 3 quarts of this product as a broadcast spray in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields. See additional instructions on the use of shielded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Over-the-Top Wiper Application

USE INSTRUCTIONS: This product may be applied over the top of desirable grasses using wiper applicators for the control of tall weeds. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Spot Treatment

USE INSTRUCTIONS: Apply a 1- to 1.5-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may also be used to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS, RESTRICTIONS: Apply this product prior to heading of grasses. Do not treat more than 10 percent of the total field area. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. Hand-held equipment maybe used to control sod remnants or other unwanted vegetation after sod is harvested.

Creating Rows in Annual Ryegrass

USE INSTRUCTIONS: Apply 16 to 32 fluid ounces of this product per acre. Best results are obtained when applications are made before ryegrass reaches 6 inches in height. Use the higher rate when the ryegrass is greater than 6 inches in height. Set nozzle heights to allow the establishment of the desired row spacing. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

PRECAUTIONS, RESTRICTIONS: Take care not to spray or allow droplets, spray lines, or drift to settle outside the treatment area to avoid unwanted crop destruction.

Grower assumes all responsibility for crop losses from misapplication. 11.4 Pastures

LABELED CROPS: Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label, including Bahiagrass, Bermudagrass, Bluegrass, Bromegrass, Fescue, Guinea grass, Kikuyu grass, Orchardgrass, Pangola grass, Ryegrass, Timothy, Wheatgrass.

TYPES OF APPLICATIONS: Preplant, Preemergence, Pasture Renovation, Spot Treatment, Over-the-Top Wiper Application, Postemergence Weed Control (Broadcast Treatments).

Preplant, Preemergence, Pasture Renovation

USE INSTRUCTIONS: This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be used to control perennial pasture species listed on this label prior to re-planting.

I. Main Label For Food Crop Uses

PRECAUTIONS, RESTRICTIONS: If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed on this label, applications must be made at least 30 days prior to planting.

Spot Treatment, Over-the-Top Wiper Application

USE INSTRUCTIONS: This product may be applied in pastures as a spot treatment or over the top of desired grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: For spot treatment and wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acre, no more than 10 percent of the field may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing or harvesting for feed.

Postemergence Weed Control (Broadcast Treatments)

USE INSTRUCTIONS: This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto pasture grasses except for renovation use described above in this section. If replanting is needed due to severe stand reduction, wait at least 30 days after treatment prior to planting any crop not listed on this label.

11.5 Rangelands

TYPES OF APPLICATIONS: Postemergence.

USE INSTRUCTIONS: This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands. Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Apply 12 to 16 fluid ounces of this product to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are recommended where spring moisture is usually limited and fall germination allows for good weed growth.

For medusahead, apply 16 fluid ounces of this product at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

PRECAUTIONS, RESTRICTIONS: Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off. Do not use ammonium sulfate when spraying rangeland grasses with this product. No waiting period between treatment and feeding or livestock grazing is required. Do not apply more than 3 quarts of this product per acre per year.

I. Main Label For Food Crop Uses

12.0 ROUNDUP READY CROPS

The following instructions or those published separately on Repar-Glypho, LLC, supplemental labeling for this product include all applications of this product that can be made onto the specified Roundup Ready crops during the complete cropping season. DO NOT combine these treatment instructions with those for crops in the "ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)" section of this label that do not contain a glyphosate tolerant gene, unless otherwise directed in this product's labeling.

REPAR-GLYPHO, LLC, COMPANY RECOMMENDS USE OF THIS PRODUCT FOR POSTEMERGENCE (IN-CROP) APPLICATION ONLY ON CROPS DESIGNATED AS CONTAINING A GLYPHOSATE TOLERANT GENE. Applying this product to crops that are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a glyphosate tolerant gene, since severe plant injury or destruction will result. The Roundup Ready designation indicates that the crop contains a patented gene that provides tolerance to this product. Information on Roundup Ready crops may be obtained from your seed supplier or Repar-Glypho, LLC, representative. Roundup Ready crops must be purchased from an authorized licensed seed supplier.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground broadcast applications. Apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications. All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling, in particular in the "Aerial Equipment" section. Apply this product in 3 to 15 gallons of water per acre. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

For proper stewardship of aerial applications over the top of Roundup Ready crops, Repar-Glypho, LLC, recommends that growers and applicators read and follow all precautions and procedures contained in the use guide "A Guide to On-Target Aerial Application." Copies of this publication are available by calling 1-800-ROUNDUP (1-800-768-6387).

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE.

TANK MIXTURE: Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT recommended for applications of this product over the top of Roundup Ready crops unless otherwise noted in this product label, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC, for this product.

I. Main Label For Food Crop Uses

Unless otherwise directed, nonionic surfactant may be added to the spray solution for applications to Roundup Ready crops. The addition of certain surfactants to this product may result in some crop response including leaf necrosis, leaf chlorosis or leaf speckling due to the surfactant added to the spray mixture. Read and carefully observe cautionary statements and other information appearing on the surfactant label.

Ammonium sulfate may be mixed with this product for applications to Roundup Ready crops. Refer to the "MIXING" section for instructions on the use of ammonium sulfate. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The following instructions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of this product is recommended to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed with multiple germination times, or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Recommended rates of this product specified in the following sections, or in separate supplemental labeling published by Repar-Glypho, LLC, for this product on the control of tough weeds, supersede the general rate recommendations in the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" of this label.

12.1 Roundup Ready Alfalfa

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: Refer to the following table for the maximum application rates of this product.

Annual Maximum Application Rates	
Combined total per year for all applications, including preplant during year of establishment	8 quarts per acre
Combined total per year for in-crop applications for newly established and established stands	6 quarts per acre

GENERAL PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting alfalfa with a glyphosate tolerant gene, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

USE INSTRUCTIONS: Applications of this product may be made over the top of Roundup Ready alfalfa (in-crop) from emergence to 5 days prior to harvest. To maximize crop yield and quality potential, applications of this product should be made after weeds have emerged but before alfalfa growth or re-growth interferes with spray coverage of the target weeds.

Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" in this label for rate recommendations for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. In addition to those weeds listed in these sections, this product will suppress the parasitic weed Dodder (*Cuscuta spp*) in Roundup Ready alfalfa. Repeat applications may be necessary for complete control.

I. Main Label For Food Crop Uses

Stand Establishment—Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings may not contain a Roundup Ready gene and will not survive after the first application of this product. To eliminate the undesirable effects of stand gaps created by this loss of plants, a single application of this product per acre should be applied at or before the 4-trifoliate growth stage. Refer to the following tables for maximum in-crop application rates during stand establishment (seeding year).

STAND ESTABLISHMENT (Seeding Year)	
Application Rates	
Prior to First Cutting	
From emergence up to 4 trifoliate leaves	1 to 2 quarts per acre
From 5 trifoliate leaves up to 5 days before first cutting	Up to 2 quarts per acre
After First Cutting	
In-crop application, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

Established Stands—Refer to the following table for directions and maximum application rates for in-crop applications to established stands of alfalfa (non-seeding year).

ESTABLISHED STANDS	
Application Rates	
In-crop applications, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

PRECAUTIONS, RESTRICTIONS: Where Roundup Ready alfalfa is grown with a companion or cover crop, or is overseeded with a second species, in-crop (over-the-top) applications of this product will eliminate the non-glyphosate tolerant species. Any single in-crop application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart. The combined total per year for all in-crop applications in both newly established (seeding year) and established stands (non-seeding year) must not exceed 6 quarts per acre. Remove domestic livestock before application. Wait a minimum of 5 days after application before grazing, cutting and feeding of forage and hay.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application.

Newly Seeded Stands and Stand Establishment: For control of emerged annual grasses and broadleaf weeds, this product may be applied at up to 2 quarts per acre in a tank mixture with the following herbicides. Application should be made after weeds have emerged but before the alfalfa growth or regrowth would interfere with spray coverage of the target weeds.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, bromoxynil, clethodim, imazamox, imazethapyr, sethoxydim, Arrow, Buctril, Butoxone, Butyrac, Poast, Pursuit, Raptor, Select.

Buctril can only be used in newly seeded stands]

Dormant Application and Winter Treatment: For control of emerged annual grasses and broadleaf weeds, up to 2 quarts per acre of this product may be applied in a tank mixture with the following herbicides. Tank mixtures should be applied when the temperature for the day remains above freezing.

I. Main Label For Food Crop Uses

Dormant Application

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: diuron, hexazinone, imazamox, metribuzin, pronamide, terbacil.

Kerb 50-W, Lexone, Raptor, Sencor, Sinbar, Velpar AlfaMax.]

Winter Treatment

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, diuron, hexazinone.

Butoxone, Diruon, Velpar AlfaMax.]

PRECAUTIONS, RESTRICTIONS: Tank mixtures of this product with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control. Applications should not be made to frozen or snow covered ground.

12.2 Roundup Ready Canola (Spring)

Roundup Ready spring canola is defined as those Roundup Ready canola varieties that are seeded in the spring and harvested in the fall and do not enter a winter dormancy period.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

Annual Maximum Application Rates

Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	1 quart per acre

GENERAL PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "**Annual Maximum Application Rates.**"

Preplant, At-planting, Preemergence,

USE INSTRUCTIONS: This product maybe applied before, during or after planting, up to a maximum of 2 quarts per acre. Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready spring canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds. .

Single Application - Apply 16 to 24 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar crop injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application - Apply 16 fluid ounces per acre to 1 - to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications are recommended for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass, or when multiple application times are needed for adequate weed control.

I. Main Label For Food Crop Uses

PRECAUTIONS, RESTRICTIONS: No more than two in-crop broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 32 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

12.3 Roundup Ready Canola (Winter)

Roundup Ready Winter Canola is defined as those Roundup Ready canola varieties that are seeded in early fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.

<u>Annual Maximum Application Rates</u>	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	1 quart per acre

DO NOT USE THIS PRODUCT ON WINTER CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "**Annual Maximum Application Rate.**"

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied to Roundup Ready winter canola varieties from emergence to canopy closure in the fall and prior to bolting in the spring. Applications made during or after bolting may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Some weeds with multiple germination times, or suppressed (stunted) weeds, or weeds that have overwintered may require sequential applications of this product for control. The second application should be made after some regrowth has occurred and at least 60 days after a previous application of this product.

Single Application—Apply 24 to 32 fluid ounces of this product per acre in the fall. Applications in the fall should be made when weeds are small and actively growing. Use the higher rate in the recommended range when weed densities are high, when weeds have overwintered or when weeds become large and well established. Applications of greater than 16 fluid ounces per acre prior to the 6-leaf stage may result in reduced crop growth in the fall. Avoid spray overlaps. Spray overlaps may result in temporary yellowing and/or growth reduction.

Sequential Applications—Apply 16 to 32 fluid ounces of this product per acre to 2-leaf or larger canola in the fall, followed by a sequential application at the same rate and at a minimum interval of 60 days, but before bolting in the spring. Sequential applications are recommended for early emerging annual weeds and winter emerging weeds such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product will control or suppress most perennial weeds. For some perennial weeds, sequential applications may be required to reduce competition with the crop.

PRECAUTIONS, RESTRICTIONS: No more than two over-the-top broadcast applications may be made from crop emergence up to the onset of bolting, and the total in-crop application must not exceed 64 fluid ounces of this product per acre. Applications of greater than 24 fluid ounces per acre prior to the 6-leaf stage

1. Main Label For Food Crop Uses

may result in reduced crop growth in the fall. Allow a minimum of 60 days between last application and harvest of canola grain. No waiting period is required between application and open grazing of livestock.

12.4 Roundup Ready Corn 2

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Spot treatment, Preharvest, Post-harvest.

<u>Annual Maximum Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from emergence through 48 inches	3 quarts per acre
Maximum preharvest application (See PRECAUTIONS, RESTRICTIONS section for Preharvest)	1 quart per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application. Read and follow label directions for all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to corn: 2,4-D, acetochlor, alachlor, atrazine, carfentrazone-ethyl, clopyralid, dicamba, diflufenzopyr, dimethenamid, dimethenamid-p, flufenacet, flumetsulam, flumiclorac pentyl ester, isoxaflutole, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, rimsulfuron.

Aim, Axiom, Balance, Balance PRO, Banvel, Bicep MAGNUM, Bicep II MAGNUM, Bicep Lite II MAGNUM, Bullet, Cinch, Cinch ATZ, Clarity, Define, Degree, Degree Xtra, Distinct, Dual II MAGNUM, Epic, Frontier, FullTime, Guardsman MAX, Harness, Harness Xtra, Harness Xtra 5.6L, Homer, Keystone, Keystone LA, Lariat, Linex, Lorox, Marksman, Me-Too-Lachlor 11, Micro-Tech, Prowl, Python, Python 11, Radius, Resolve, Resource, Stalwart, Stalwart Xtra, Surpass, TopNotch]

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

USE INSTRUCTIONS: This product may be applied alone or in tank mixtures postemergence (in-crop) to corn hybrids designated as Roundup Ready Corn 2. When applied as directed, this product controls annual grass and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. A postemergence application of this product should be made before the weeds reach a height of 4 inches or before they become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre should be made before the weeds reach a height of 4 inches.

This product may be broadcast over-the-top or applied with drop nozzles to Roundup Ready Corn 2 from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing).

I. Main Label For Food Crop Uses

whichever comes first. Drop nozzles are recommended for optimum spray coverage and weed control when corn height is 24 to 30 inches. For corn heights 30 to 48 inches (free standing), apply this product only by ground application using drop nozzles aligned to avoid spraying into the whorls of the corn plants.

TANK MIXTURES: This product may be tank-mixed with the following products. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to corn. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to corn: atrazine, acetochlor, alachlor, clopyralid, dicamba, diflufenzopyr, flumetsulam, flumiclorac pentyl ester, foramsulfuron, halosulfuron-methyl, iodosulfuron-methyl-sodium, mesotrione, nicosulfuron, rimsulfuron, thifensulfuron methyl].

Banvel, Basis, Basis Gold, Bullet, Callisto, Clarity, Degree, Degree Xtra, Distinct, Equip, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Marksman, Micro-Tech, Option, Resolve, Resource, YUKON]

Tank Mix Partner	Maximum Height Of Corn For Application
Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6L	11 inches
Bullet* Micro-Tech*	5 inches
Atrazine	12 inches

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet* or Micro-Tech* herbicides.

PRECAUTIONS, RESTRICTIONS: Single in-crop applications of this product should not exceed 3 pints per acre. Sequential in-crop applications of this product from emergence through 48 inches in height must not exceed a total of 3 quarts per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product in-crop and harvest of corn forage or grain. (For applications at preharvest timing, see **Preharvest** section of this label). The use of additional surfactants and other additives, including fertilizers and micro-nutrients, in the spray solution are not recommended with this product and may result in increased potential for crop injury or reduced yield, and is not recommended for over-the-top applications, unless otherwise directed in supplemental labeling of fact sheets published separately by Repar-Glypho, LLC.

Preharvest

USE INSTRUCTIONS: A single preharvest application of up to 32 fluid ounces per acre of this product may be applied for annual and perennial weed control prior to crop harvest. Make application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

I. Main Label For Food Crop Uses

PRECAUTIONS, RESTRICTIONS: Do not make a preharvest application of this product if more than a combined total of 64 fluid ounces of this product has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

Postharvest

USE INSTRUCTIONS: This product may be applied for weed control after corn harvest. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

12.5 Roundup Ready Cotton

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Selective Equipment (In-Crop), Preharvest.

<u>Annual Maximum Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to layby	4 quarts per acre
Maximum preharvest application rate	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." **ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.**

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting. Tank mixtures with other herbicides listed in Section 8.3 of this label maybe used.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to the emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyriithiobac-sodium.

Caparol, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H2O, Stalwart, Staple]

PRECAUTIONS, RESTRICTIONS: The maximum quantity of this product that maybe applied for all preplant, at-planting and preemergence applications combined is 5 quarts per acre per season.

Postemergence (in-crop)

USE INSTRUCTIONS: This product may be applied over the top Roundup Ready cotton at rates up to 1 quart per acre per application from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). **Applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss.**

I. Main Label For Food Crop Uses

TANK MIXTURES: This product may be tank-mixed with the following products and applied over the top of Roundup Ready cotton: Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clethodim, fluazifop-P-butyl, metolachlor, s-metolachlor, pyriithiobac-sodium, sethoxydim, quizalofop-P-ethyl, Assure 11, Dual MAGNUM, Fusilade, Poast Plus, Select, Stalwart, Staple
Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop).
Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting to exposed cotton leaves.]

PRECAUTIONS, RESTRICTIONS: The maximum quantity of this product that maybe applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. No more than two over-the-top broadcast applications may be made from crop emergence through the 4-leaf (node) stage of development. No more than two applications should be made from the 5-leaf stage through layby. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications. Do not add additional surfactant or additives containing surfactant to this product (other than those contained in any tank-mix product) for over-the-top applications to roundup ready cotton.

Salvage Treatment. This treatment may be used after the 4-leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

Selective Equipment (In-crop)

USE INSTRUCTIONS: This product may be applied using precision post-directed or hooded sprayers at rates up to 1 quart per acre per application to Roundup Ready cotton through layby. At this stage, use post-directed equipment to direct the spray to the base of the cotton plants. Avoid contact of the herbicide spray with the cotton leaves to the maximum extent possible. To minimize spray contact maintain low spray pressure (less than 30 pounds per square inch) and place nozzles in a low position, directing a horizontal spray pattern under the cotton leaves and onto weeds in the row. For best results, make applications while weeds are small (less than 3 inches). See additional instructions on the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: This product can be tank-mixed with the following products for in-crop application using precision post-directed or hooded sprayers. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: carfentrazoneethyl, diuron, flumioxazin, fluometuron, linuron, pendimethalin, prometym, pyriithiobac-sodium, Aim, Caparol, Chateau, Cotoran, Direx, Layby-Pro, Prowl H2O, Staple, Valor
Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop)]

PRECAUTIONS, RESTRICTIONS: The maximum quantity of this product that may be applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

I. Main Label For Food Crop Uses

Preharvest

USE INSTRUCTIONS: This product may be applied for annual and perennial weed control as a broadcast treatment prior to crop harvest after 20 percent boll crack. Apply up to 2 quarts of this product per acre. **NOTE:** This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Preharvest application is not recommended for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION; Use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.6 Roundup Ready Flex Cotton

The use of the over-the-top applications described in this section on **other** than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (node) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this section are specific to, and should only be used with, varieties designated as Roundup Ready Flex cotton. Do Not combine the instructions in this section, with those in the "Roundup Ready Cotton" section (12.5) of this label, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instructions on labeling for this or other glyphosate-containing product.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Preharvest.

<u>Annual Maximum Allowable Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to 60 percent open bolls	6 quarts per acre
Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyridiobac-sodium.

I. Main Label For Food Crop Uses

Caparol, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H2O, Stalwart, Staple

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied to control annual grasses and broadleaf weeds listed on this label in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds is recommended. This product may be applied postemergence at rates up to 1.5 quarts per acre per application. In addition to broadcast applications, post-directed spray equipment may be used to achieve weed coverage.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready Flex cotton. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clethodim, fluazifop-Pbutyl, metolachlor, s-metolachlor, pyriithiobac-sodium, quizalofop-p-ethyl, sethoxydim, trifloxysulfuron-sodium.

Assure 11, Dual MAGNUM, Envoke, Fusilade, Poast Plus, Select, Stalwart, Staple

Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop).

Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting to exposed cotton leaves.)

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 1.5 quarts per acre made using ground application equipment. In-crop application rates above 1 quart per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling or leaf necrosis. Do not exceed a maximum rate of 1 quart per acre of this product *when* making applications by air. Between layby and 60 percent open bolls, the maximum combined total rate of this product that may be applied is 2 quarts per acre. The maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre. Do not add additional surfactant or additives containing surfactant to this product for over-the-top applications to Roundup Ready Flex cotton.

Preharvest

USE INSTRUCTIONS: Up to 2 quarts of this product may be applied for annual and perennial weed control as a broadcast treatment prior to harvest after 60 percent boll crack. **NOTE:** This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Preharvest application is not recommended for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION: use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Flex Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.7 Roundup Ready Soybeans

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Preharvest, Post-Harvest.

I. Main Label For Food Crop Uses

<u>Annual Maximum Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from cracking through flowering (R2 stage soybeans)	3 quarts per acre
Maximum preharvest application rate	1 quart per acre

GENERAL PRECAUTIONS/RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on **"Annual Maximum Application Rates."**

Refer to individual tank mixture product label for restrictions and precautions. Use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D, Banvel or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to soybeans: alachlor, atrazine, r-arfentrazone-ethyl, chlorimuron ethyl, clethodim, clomazone, cloransulam-methyl, dimethenamid, dimethenamid-p, fenoxypyr, fluazifop-p-butyl, flufenacet, flumetsulam, flumiclorac pentyl ester, flumioxazin, fomesafen, imazaquin, imazethapyr, lactofen, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, sulfentrazone, tribenuron methyl, quizalofop P-ethyl.

Aim, Amplify, Assure II, Axiom, Authority, Blanket, Boundary, Canopy, Canopy EX, Classic, Cobra, Command, Command Xtra, Domain, Dual MAGNUM, Dual II MAGNUM, FirstRate, Flexstar, Frontier, Fusion, Gangster, Gauntlet, INTRO, Lexone, Linex, Linuron, Lorox, Lorox Plus, Me-Toolachlor, Micro-Tech, Outlook, Pendimax, Prowl, Prowl H2O, Pursuit, Pursuit Plus, Python, Reflex, Resource, Scepter, Select, Select MAX, Sencor, Spartan, Squadron, Steel, Valor]

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be used to control annual grasses and broadleaf weeds listed on this label in Roundup Ready soybeans. Applications of this product can be made in from emergence (cracking) through flowering (R2 stage soybeans). R2 stage soybeans ends when a pod reaches 5 millimeters (0.2 inches) in length at one of the four uppermost nodes on the main stem with a fully developed leaf. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate recommendations for specific annual weeds. In general, an initial application of 1 quart per acre on 2- to 8-inch tall weeds is recommended. Weeds will generally be 2 to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 2 quarts per acre in any single in-crop application for control of annual weeds and where dense weed populations exist.

A 1- to 2-quarts per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, mare's tail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds. **IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE ROUNDUP READY SOYBEAN CROP.** To control giant ragweed, it is recommended that 1 quart per acre of this product be applied when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

I. Main Label For Food Crop Uses

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready soybeans. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to soybeans: acifluorfen, bentazonechlorimuron ethyl, clethodim, imazethapyr, cloransulam-methyl, fenoxypop, fluazifop-pbutyl, flumiclorac pentyl ester, fomesafen, imazamox, imazethapyr, lactofen, pendimethalin, quizalofop P-ethyl, sethoxyden, thifensulfuron-methyl.

Arrow, Assure II, Basagran, Classic, Cobra, Extreme, FirstRate, Flexstar, Fusilade DX, Fusion, Harmony GT XP, Poast, Poast Plus, Pursuit, Pursuit Plus, Raptor, Reflex, Select, Select MAX, Synchrony STS, Targa, Ultra Blazer.]

PRECAUTIONS, RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in-crop application is 2 quarts per acre. The maximum combined total of this product that can be applied during flowering is 2 quarts per acre.

Preharvest

USE INSTRUCTIONS: This product may be applied for weed control prior to harvest of soybeans after pods have set and lost all green color. Up to 1 quart per acre of this product can be applied by aerial or ground application.

PRECAUTIONS, RESTRICTIONS: Care should be taken to avoid excessive seed shatter loss due to ground application equipment. Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of Roundup Ready soybeans. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

12.8 Roundup Ready Sugar Beet

Maximum Annual Application Rates

Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Emergence to 8-leaf stage	2.5 quarts per acre
Between 8-leaf stage and canopy closure	2 quarts per acre

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions. Use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of sugar beet. Read and follow label directions of all products in the tank mixture.

I. Main Label For Food Crop Uses

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplan, at-planting and/or preemergence to sugar beet: dimethenamid, s-metolachlor.

Dual MAGNUM, Frontier]

Postemergence (In-Crop)

USE INSTRUCTIONS: This product may be applied over-the-top of Roundup Ready sugar beets for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate recommendations for specific annual weeds. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications maybe required to eliminate crop competition throughout the growing season.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready sugar beet. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to sugar beet. Read and follow label directions of all products in the tank mixture

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to sugar beet: clethodim, clopyralid, desmedipham, ethofumesate, phenmedipham, quizalofop-p-ethyl, sethoxydim, trisulfuron-methyl.

Assure II, Betamix, Betanex, Norton SC, Poast, Progress, Select, Stinger, Upbeet]

PRECAUTIONS, RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 4.5 quarts per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 1.5 quarts per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 1 quart per acre. Allow a minimum of 30 days *between* last application and sugar beet harvest.

13.0 NON-CROP USES AROUND THE FARMSTEAD

TYPES OF APPLICATIONS: General Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stump, Habitat Management.

13.1 General Weed Control, Trim-And-Edge

USE INSTRUCTIONS: This product may be used to control annual weeds, perennial weeds and woody brush found in any part of the farmstead, including around building foundations and equipment storage areas, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, rangeland, rights-of-way, shelterbelts, and prior to planting landscape ornamentals. Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" in this label for treatment rates. For application through backpack sprayers, handguns or other high-volume spray-to-wet applications, *see* the "Annual Weeds -- Hand-Held or Backpack Equipment" section of this label for further instructions.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product used is registered for treatment of these non-crop sites. Refer to the individual product labels for approved farmstead sites and application rates. For more information, *see* the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Arsenal	Plateau
Barricade 65WG	Princep DF
Crossbow I.	Princep 4L
Diuron	Ronstar 50 WP
Dicamba	Sahara
Endurance	Simazine

I. Main Label For Food Crop Uses

Escort	Surflan AS
Goal 2XL	Surflan WDG
Karmex DF	Telar
Krovar I DF	Transline
Outrider	Vanguard
Oust	Velpar DF
Pendimethalin	Velpar L
Pendulum 3.3 EC	2,4-D Pendulum WDG

This product plus dicamba tank mixtures may not be applied by air in California. 13.2 **Greenhouse/Shadehouse**

USE INSTRUCTIONS: This product may be used to control weeds in and around greenhouses and shade houses.

PRECAUTIONS, RESTRICTIONS: Desirable vegetation must not be present during application. Air circulation fans must be turned off. Do not use in residential greenhouses.

13.3 Chemical Mowing

USE INSTRUCTIONS: This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Use 16 fluid ounces of this product per acre when treating Bermudagrass. Use 64 fluid ounces of this product per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

PRECAUTIONS, RESTRICTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

13.4 Cut Stump

TYPES OF APPLICATIONS: Treating cut stumps in any non-crop site listed on this label.

USE INSTRUCTIONS: This product will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Salt-cedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, brazilian	Willow Pine, Austrian

PRECAUTIONS, RESTRICTIONS: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

13.5 Habitat Management

TYPES OF APPLICATIONS: Habitat Restoration and Maintenance, Wildlife Food Plots.

Habitat Restoration and Maintenance

USE INSTRUCTIONS: This product may be used to control exotic and other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

I. Main Label For Food Crop Uses

Wildlife Food Plots

USE INSTRUCTIONS: This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species, including Roundup ready canola, may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage. For specific product instructions for Roundup Ready canola wildlife food plots, see the "ROUNDUP READY CANOLA" section of this label.

PRECAUTIONS, RESTRICTIONS: Do not process treated Roundup ready canola seeds from Roundup ready canola wildlife food plots for food. Do not graze or feed treated Roundup Ready canola from wildlife food plots to livestock. There are no rotational restrictions for planting any wildlife food species or for allowing native species to repopulate the area following applications of this product.

14.0 ANNUAL WEEDS RATE SECTION

When water carrier volumes are between 16 and 40 gallons per acre for ground applications and between 6 and 15 gallons per acre for aerial applications, the following use rates will control the annual weeds listed in the table below:

- 1 quart per acre – grass and broadleaf annual weeds less than 6 inches in height or circumference and vines less than 3 inches in length.
- 3 pints per acre – grass and broadleaf annual weeds 6 to 12 inches in height or circumference and vines 3 to 6 inches in length.
- 2 quarts per acre – grass and broadleaf annual weeds greater than 12 inches in height or circumference and vines greater than 6 inches in length.

WHEN WATER CARRIER VOLUMES ARE BETWEEN 3 AND 15 GALLONS PER ACRE FOR GROUND APPLICATIONS AND BETWEEN 3 AND 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS, USE THE RATES SPECIFIED FOR INDIVIDUAL WEEDS AS FOLLOW IN THE "ANNUAL WEEDS RATE TABLE" (Alphabetically by Species).

Apply to actively growing annual weeds. Annual weeds are generally easiest to control when they are small.

Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Maximum size refers to the maximum plant height, length of runners for vines, or circumference of rosette plants in inches.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment. This product may be used up to 64 fluid ounces per acre where heavy weed densities exist.

ANNUAL WEEDS RATE TABLE
(Alphabetically by Species)

WEED SPECIES	RATE (Fluid Ounces Per Acre)				
	16	24	32	40	48
	(Maximum Size (in inches))				
Ammannia, purple	3	6	12	-	18
Anode, spurred	-	2	3	5	8
Barley	18	18+	-	-	-
Barnyardgrass	-	3	6	7	9
Basle, fivehook	-	-	6	-	-
Beggarweed, Florida	-	5	8	-	-
Bittercress	12	20	-	-	-
Bluegrass, annual	10	-	-	-	-

Bluegrass, bulbous	6	-	-	-	-
Brome, downy ^{1,2}	6	12	-	-	-
Brome, Japanese	6	12	24	-	-
Browntop panicum	6	8	12	-	24
Buckwheat, wild ³	-	1	2	-	-
Burcucumber	-	6	12	-	18
Buttercup	12	20	-	-	-
Carolina geranium	-	-	4	-	9
Carpetweed	-	6	12	-	-
Cheat ²	6	20	-	-	-
Chervil	20	-	-	-	-
Chickweed	-	12	18	-	-
Cocklebur	12	18	24	-	36
Copperleaf, hophornbeam	-	2	4	-	6
Copperleaf, Virginia	-	2	4	-	6
Coreopsis, plains	-	6	12	-	18
Corn, volunteer	6	12	20	-	-
Corn speedwell	12	-	-	-	-
Crabgrass	3	6	12	-	-
Crowfootgrass	-	-	6	-	12
Cutleaf evening primrose	-	-	3	-	6
Devilsclaw (unicorn plant)	-	3	6	-	-
Dwarf dandelion	12	-	-	-	-
Eastern mannagrass	8	12	-	-	-
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
Falsedandelion	-	20	-	-	-
Falseflax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	6	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail, giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	6	12	-	-	-
Goosegrass	-	3	6	-	12
Grain sorghum (mild)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundset, common	-	6	10	-	-
Hemp sesbania	-	2	4	6	8
Henbit	-	-	6	-	12
Horseweed/ Maretail (<i>Conyza canadensis</i>)*	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnsongrass, seedling	6	12	18	-	24
Junglerice	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia ⁴	-	3 to 6	12	-	-
Lambsquarters*	-	6	12	-	20
Little barley	6	12	-	-	-

London rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morningglory, annual (<i>Ipomoea spp</i>)	-	-	3	-	6
Mustard, blue	6	12	18	-	-
Mustard, tansy	6	12	18	-	-
Mustard, tumble	6	12	18	-	-
Mustard, wild	6	12	18	-	-
Nightshade, black	-	4	6	-	12
Nightshade, hairy	-	4	6	-	12
Oats	3	6	18	-	-
Pigweed, Palmer*	-	12	18	24	-
Pigweed species*	-	12	18	24	-
Prickly lettuce	-	6	12	-	-
Purslane	-	-	3	-	6
Ragweed, common*	-	6	12	-	18
Ragweed, giant*	-	6	12	-	18
Red rice	-	-	4	-	-
Rye, volunteer/cereal ²	6	18	18+	-	-
Ryegrass species*	-	-	6	-	12
Sandbur, field	6	12	-	-	-
Sandbur, longspine	6	12	-	-	-
Shattercane	6	12	20	-	-
Shepherd's-purse	6	12	-	-	-
Sicklepod	-	2	4	-	8
Signalgrass, broadleaf	-	3	6	7	9
Smartweed, ladysthumb	-	-	6	-	9
Smartweed, Pennsylvania	-	-	6	-	9
Sowthistle, annual	-	-	6	-	12
Spanishneedles	-	-	6	-	12
Speedwell, purslane	12	-	-	-	-
Sprangletop	6	12	20	-	-
Spurge, prostrate	-	6	12	-	-
Spurge, spotted	-	6	12	-	-
Spurry, umbrella	6	-	-	-	-
Stinkgrass	-	12	-	-	-
Sunflower	12	18	-	-	-
Swinecress	-	5	12	-	-
Teaweed/ Prickly sida	-	2	4	-	6
Texas panicum	6	8	12	-	24
Thistle, Russian ²	-	6	12	-	-
Velvetleaf	-	-	6	-	12
Virginia pepperweed	-	18	-	-	-
Waterhemp*	-	-	6	-	12
Wheat ²	6	12	18	-	-
Wheat, (overwintered)	-	6	12	-	18
Wild oats	3	6	18	-	-
Wild proso millet	-	6	12	-	18
Witchgrass	-	12	-	-	-
Woolly cupgrass	-	6	12	-	-
Yellow rocket	-	12	20	-	-

1. Main Label For Food Crop Uses

¹ For control of downy brome in no-till systems, use 24 fluid ounces per acre.

² Performance is better if application is made before this weed reaches the boot stage of growth.

³ Use 24 fluid ounces per acre of this product to control wild buckwheat in the cotyledon to 2-leaf stage. Use 32 fluid ounces per acre to control 2- to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 32 fluid ounces followed by 32 fluid ounces of this product per acre.

⁴ Do not treat kochia in the button stage.

⁵ Control of Russian thistle may vary based on environmental conditions and spray coverage. Whenever possible, a tank mixture with 2,4-D as described below may improve control.

* A glyphosate resistant biotype has been confirmed. For additional information, refer to the "WEED RESISTANCE MANAGEMENT" section of this label. You may also visit on the Internet www.weedscience.org or www.weedresistancemanagement.com, or contact your Repar-Glypho, I.I.C. representative.

14.1 Annual Weeds — Tank Mixtures with 2,4-D or Dicamba or Tordon 22K

Better control of certain tough weeds can be achieved by tank mixing this product with 0.25 pound of dicamba, 0.5 pound of 2,4-D or 1 to 2 fluid ounces of Tordon 22K per acre. These other herbicides, combined with the rates of this product specified in the "ANNUAL WEEDS RATE TABLE" above, will control the following weeds up to the maximum height or length indicated: 6" -- prickly lettuce, mare's tail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Tordon 22K only); 12" -- cocklebur, lambsquarters, pigweed, Russian thistle (2,4-D only). For better control of common ragweed, giant ragweed, Pennsylvania smartweed, or velvetleaf with a maximum height of 6-inches, tank mix this product with 0.5 pound of 2,4-D per acre.

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA.

14.2 Annual Weeds—Hand-Held or Backpack Equipment

For control of weeds listed in the "ANNUAL WEEDS RATE TABLE" above, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1-percent solution.

For best results, use a 2-percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 5-percent solution for annual and perennial weeds and a 5- to 10-percent solution for woody brush and trees.

14.3 Annual Weeds — Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

Applications of 24 to 28 fluid ounces of this product plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 28 ounces for control), Downy brome, Green foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass and Kochia (add 0.12 pound of dicamba for control).

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines.

I. Main Label For Food Crop Uses

15.0 PERENNIAL WEEDS RATE SECTION

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage. Best results are obtained when soil moisture is adequate for active weed growth.

PERENNIAL WEEDS RATE TABLE

(Alphabetically by Species)

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alfalfa	1 - 2	3 - 10	2%

Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

Alligatorweed	4	3 - 20	1.5%
----------------------	----------	---------------	-------------

For partial control, apply when most of the plants are in bloom. Repeat applications will be required to achieve control.

Anise (fennel) ¹	-	-	1-2%
------------------------------------	----------	----------	-------------

Apply as a spray-to-wet treatment.

Bahiagrass²	3-5	3-20	2%
-------------------------------	------------	-------------	-----------

Bentgrass	3-5	10-20	2%
------------------	------------	--------------	-----------

For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results.

Bermudagrass	3 - 5	3 - 20	2%
---------------------	--------------	---------------	-----------

For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to achieve control.

Bermudagrass, water (knotgrass)	1 - 1.5	5 - 10	2%
--	----------------	---------------	-----------

Apply 3 pints of this product in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart this product in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length.

This product is not registered in California for use on water Bermudagrass.

Bindweed, field	0.5-5	3 - 20	2%
------------------------	--------------	---------------	-----------

Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.

For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre

east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 2 quarts of this product plus 0.5 pound of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. Irrigate at least once to promote active bindweed growth.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky	1 - 2	3 - 40	2%
----------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas	3 - 5	3 - 40	2%
------------------------	--------------	---------------	-----------

Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenfern	3 - 4	3 - 40	1 - 1.5 %
--------------------	--------------	---------------	------------------

Apply to fully expanded fronds that are at least 18 inches long.

Bromegrass, smooth	1 - 2	3 - 40	2%
---------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf	---	3 - 20	2%
-----------------------------	------------	---------------	-----------

For control, apply 2 quarts of this product plus 0.5 pound of dicamba per acre. For partial control, apply 1 quart of this product plus ½ pound of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed²	2 - 3	3 - 40	2%
--------------------------------------	--------------	---------------	-----------

Cattail²	3 - 5	3 - 40	2%
----------------------------	--------------	---------------	-----------

Clover; red or white¹	3 - 5	3 - 20	2%
---	--------------	---------------	-----------

Also for control, apply 16 to 32 fluid ounces of this product plus 0.5 to 1 pound of 1,4-D, in 3 to 10 gallons of water per acre.

Cogongrass	3 - 5	10 - 40	2%
-------------------	--------------	----------------	-----------

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to achieve control.

I. Main Label For Food Crop Uses

Dallisgrass²	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Dandelion¹	3 – 5	3 – 40	2%
------------------------------	--------------	---------------	-----------

Also for control, apply 1 pint of this product plus 0.5 of 2,4-D, in 3 to 10 gallons of water per acre.

Dock, curly¹	3 – 5	3 – 40	2%
--------------------------------	--------------	---------------	-----------

Also for control, apply 1 to 2 pints of this product plus 0.5 to 1 pound of 2,4-D, in 3 to 10 gallons of water per acre.

Dogbane, hemp	4	3 – 40	2%
----------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue (except tall)¹	3 – 5	3 – 20	2%
---	--------------	---------------	-----------

Fescue, tall	1 – 3	3 – 40	2%
---------------------	--------------	---------------	-----------

Apply 3 quarts of this product per acre when most plants have reached boot-to-early seedhead stage of development.

Fall applications only: Apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.

Guineagrass	2 - 3	3 - 40	1%
--------------------	--------------	---------------	-----------

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 2 quarts of this product per acre for control. In the flatwoods region of Florida, 3 quarts of this product per acre are required for control.

Horsenettle¹	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Horseradish	4	3 – 40	2%
--------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Iceplant¹	---	---	1.5 – 2%
-----------------------------	------------	------------	-----------------

Thorough coverage is necessary for best control.

Jerusalem artichoke¹	3 – 5	3 – 20	2%
--	--------------	---------------	-----------

Johnsongrass	0.5 – 3	3 – 40	1%
---------------------	----------------	---------------	-----------

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 1 quart of this product per acre.

For burndown of Johnsongrass, apply 1 pint of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression)--Apply a 1-percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass	2 - 3	3 - 40	2%
--------------------	--------------	---------------	-----------

Knapweed	4	3 - 40	2%
-----------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Lantana	--	--	1-1.25%
----------------	-----------	-----------	----------------

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Lespedeza	3 - 5	3-20	2%
------------------	--------------	-------------	-----------

Milkweed, common	3	3-40	2%
-------------------------	----------	-------------	-----------

Apply when most plants have reached the late bud to flower stage of growth.

Muhly, wirestem	1-2	3-40	2%
------------------------	------------	-------------	-----------

Use 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications, or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.

Mullein, common¹	3 - 5	3 - 20	2%
------------------------------------	--------------	---------------	-----------

Napiergrass²	3 - 5	3 - 20	2%
--------------------------------	--------------	---------------	-----------

Nutsedge, purple or yellow	0.5-3	3-40	1 - 2%
-----------------------------------	--------------	-------------	---------------

Apply 3 quarts of this product per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for longterm control of ungerminated tubers.

Sequential applications: 1 to 2 quarts of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For partial control of existing plants, apply 1 pint to 2 quarts of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

Orchardgrass	1 - 2	3 - 40	2%
---------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of this product in 3 to 10 gallons

I. MAIN LABEL FOR FOOD CROP USES

of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications, and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

Pampasgrass	--	--	1.5 - 2%
--------------------	----	----	----------

Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Paragrass²	3 - 5	3 - 20	2%
------------------------------	-------	--------	----

Phragmites	3 - 5	10 - 40	1 - 2%
-------------------	-------	---------	--------

For partial control and best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may result in reduced control. Due to the dense nature of this vegetation that may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to achieve control. Visual control symptoms will be slow to develop.

Poison hemlock	--	--	1 - 2%
-----------------------	----	----	--------

For hand-held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Thorough coverage is necessary for best control.

Pokeweed, common	1	3-40	2%
-------------------------	---	------	----

Apply to actively growing plants up to 24 inches tall.

Quackgrass	1 - 3	3 - 40	2%
-------------------	-------	--------	----

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of this product. Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.

In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

Redvine	0.75-2	5 - 10	2%
----------------	--------	--------	----

For suppression, apply 1.5 pints of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, giant	--	--	2%
--------------------	----	----	----

Best results are obtained when applications are made in late summer to fall.

Ryegrass, perennial	1 - 3	3 - 40	1%
----------------------------	-------	--------	----

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1 quart of this product per acre.

I. MAIN LABEL FOR FOOD CROP USES

Smartweed, swamp'	3 - 5	3 - 40	2%
Also for control, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall.			
Sowthistle, perennial	2 - 3	3 - 40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
Spurge, leafy		3 - 10	2%
For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most plants are 12 inches tall.			
Starthistle, yellow	2	10 - 40	2%
Best results are obtained when applications are made during the rosette, bolting and early flower stages.			
Sweet potato, wild			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, artichoke			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, Canada	2 - 3	3 - 40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
For suppression in the spring, apply 1 quart of this product, or 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.			
Timothy²	2-3	3-40	2%
Torpedograss	4-5	3-40	2%
For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to achieve control. Fall treatments must be applied before frost.			
Trumpet creeper	2	5 - 10	2%
For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.			
Vaseygrass²	3-5	3-20	2%
Velvetgrass²	3-5	3-20	2%

Wheatgrass, western ²	2-3	3-40	2%
----------------------------------	-----	------	----

² Apply when most plants have reached the early heading stage of growth.

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

Unless otherwise directed, apply broadcast treatments in 3 to 40 gallons of water per acre. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alder	3-4	3-40	1-1.5%
Ash	2-5	3-40	1-2%
Aspen, quaking	2-3	3-40	1-1.5%
Bearmat (Bearclover)	2-5	3-40	1-2%
Beech	2-5	3-40	1-2%
Birch	2-3	3-40	1-1.5%
Blackberry	3-4	10-40	1-1.5%

Blackgum	2-5	3-40	1-2%
Bracken	2-5	3-40	1-2%
Broom; French, Scotch			1.5-2%
Buckwheat, California, 2			1-2%

Cascara ¹	2-5	3-40	1-2%
Catsclaw	.		1-1.5%
Ceanothus ¹	2-5	3-40	1-2%
Chamise ²			1%

I. MAIN LABEL FOR FOOD CROP USES

Cherry; biter, black, pin	2-3	3-40	1-1.5%
Coyote brush	--	--	1.5-2%
Apply when at least 50 percent of the new leaves are fully developed.			
Dogwood'	2-5	3-40	1-2%
Elderberry	2-3	3-40	1-1.5%
Elm'	2-5	3-40	1-2%
Eucalyptus	--	--	2%
For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.			
Florida holly (Brazilian Peppertree) 1	2-5	3-40	1-2%
Gorse'	2-5	3-40	1-2%
Hasardia*2	--	--	1-2%
Hawthorn	2-3	3-40	1-1.5%
Hazel	2-3	3-40	1-1.5%
Hickory'	2-5	3-40	1-2%
Honeysuckle	3-4	3-40	1-1.5%
Hornbeam,	2-5	3-40	1-2%
American'			
Kudzu	4-5	3-40	2%
Repeat applications may be required to achieve control.			
Locust, black'	2-4	3-40	1-2%
Madrone resprouts'	--	--	2%
Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.			
Manzanita'	2-5	3-40	1-2%
Maple, red	2-4	3-40	1-1.5%
Apply a 1- to 1.5-percent solution when at least 50 percent of the new leaves are fully developed.			
For partial control, apply 2 to 4 quarts of this product per acre.			
Maple, sugar	--	--	1-1.5%
Apply when at least 50 percent of the new leaves are fully developed.			
Monkey flower,2	--	--	1-2%
Oak: black,white'	2-4	3-40	1-2%
Oak, post	3-4	3-40	1-1.5%
Oak: northern,	--	--	1-1.5%
Apply when at least 50 percent of the new pin leaves are fully developed			
Oak: southern red	2-3	3-40	1-1.5%
Persimmon'	2-5	3-40	1-2%
Pine	2-5	3-40	1-2%
Poison ivy/Poison oak	4-5	3-40	2%
Repeat applications may be required to achieve control. Fall treatments must be applied before leaves lose green color.			
Poplar, yellow'	2-5	3-40	1-2%
Redbud, eastern	2-5	3-40	1-2%

1. MAIN LABEL FOR FOOD CROP USES

Rose, multiflora	2	3-40	1%
Treatments should be made prior to leaf deterioration by leaf-eating insects.			
Russian olive'	2-5	3-40	1-2%
Sage, black'			1%
Sage, white'	2-5	3-40	1-2%
Sage brush, California ²			1%
Salmonberry	2-3	3-40	1-1.5%
Salt-cedar	2-5	3-40	1-2%
Sassafras'	2-5	3-40	1-2%
Sourwood'	2-5	3-40	1-2%
Sumac; poison, smooth, winged'	2-4	3-40	1-2%
Sweetgum	2-3	3-40	1-1.5%
Swordfern'	2-5	3-40	1-2%
Tallowtree, Chinese'			1%
Tan oak resprouts'	--	--	2%
Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.			
Thimbleberry	2-3	3-40	1-1.5%
Tobacco, tree'			1-2%
Trumpet creeper	2-3	3-40	1-1.5%
Vine maple'	2-5	3-40	1-2%
Virginia creeper	2-5	3-40	1-2%
Waxmyrtle, Southern'	2-5	3-40	1-2%
Willow	3-4	3-40	1-1.5%
' Partial Control			

² Thorough coverage of foliage is necessary for best results.

17.0 LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the fullest extent permitted by law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture

II. MAIN LABEL FOR INDUSTRIAL, TURF, ORNAMENTAL USES

GLYPHOSATE 41 % SL

The complete broad-spectrum postemergence herbicide for non-crop, industrial, turf and ornamental weed control.

Complete Directions for Use

EPA Reg. No. 86004-

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS). DESIRABLE PLANTS AND TREES. BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product. Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

[Container Label Statement (5 gallons or smaller):]

Do not use this container for residential use.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

[Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REAPR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC. REPACKAGING OR TOLL REPACKAGING AGREEMENT.

[Non-Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

1	1.0 INGREDIENTS
2	2.0 IMPORTANT PHONE NUMBERS
3	3.0 PRECAUTIONARY STATEMENTS
	3.1 Hazards to Humans and Domestic Animals
	3.2 Environmental Hazards
	3.3 Physical or Chemical Hazards

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- 4 **4.0 STORAGE AND DISPOSAL**
- 5 **5.0 GENERAL INFORMATION**
(How This Product Works)
5.1 Weed Resistance Management
5.2 Management Recommendations for Glyphosate Resistant Weed Biotypes
- 6 **6.0 MIXING**
6.1 Mixing with Water
6.2 Tank Mixing
6.3 Tank Mixing Procedure
6.4 Mixing for Hand-Held Sprayers
6.5 Surfactants
6.6 Colorants or Dyes
- 7 **7.0 APPLICATION EQUIPMENT AND TECHNIQUES**
7.1 Aerial Equipment
7.2 Ground Broadcast Equipment
7.3 Hand-Held or Backpack Equipment
7.4 Selective Equipment
7.5 Injection Systems 7.6 CDA Equipment
- 8 **8.0 SITE AND USE DIRECTIONS**
8.1 Cut Stump
8.2 Forestry Site Preparation
8.3 General Non-crop Areas and Industrial Sites 8.4 Habitat Management
8.5 Hollow Stem Injection
8.6 Injection and Frill (Woody Brush and Trees)
8.7 Non-food Tree, Shrub, or Vine Production Sites
8.8 Railroads
8.9 Roadsides
8.10 Rangelands
8.11 Utility Sites
8.12 Conservation Reserve program (CRP) 8.13 Grass Seed or Sod Production
8.14 Pastures
- 9 **9.0 WEEDS CONTROLLED**
9.1 Annual Weeds
9.2 Perennial Weeds
9.3 Woody Brush and Trees

10 10.0 LIMIT OF WARRANTY AND LIABILITY

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....	41.0%
OTHER INGREDIENTS (including surfactant).....	<u>59.0%</u>
	100.0%

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

3.2 Environmental Hazards

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published REPAR GLYPHO, I.I.C Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. PESTICIDE STORAGE: Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

See container label for additional CONTAINER DISPOSAL instructions.

[Container Label Storage and Disposal Statements];

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact Repar-Glypho, LLC.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Refiners: Do not reuse this container except for refill in accordance with a valid Monsanto agreement for Repackaging or Toll Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration by burning, or, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

(How This Product Works)

Product Description: This product is a postemergent, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visual effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at later growth stages approaching maturity.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Annual Maximum Application Rates: The combined total of all treatments must not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate as the active ingredient, whether applied separately or as mixtures. Calculate the application rates and ensure that the total use of this and other glyphosate containing products does not exceed stated maximum use rates. See the "INGREDIENTS" section of this label for necessary information.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

5.1 Weed Resistance Management

GROUP	9	HERBICIDE
-------	---	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using other cultural or mechanical practices.

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management recommendations:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Utilize the recommended label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Reapri-Glypho, LLC, representative, local retailer, or county extension agent.

5.2 Management Recommendations For Glyphosate Resistant Biotypes

Note: Appropriate testing is critical in order to determine if a weed is resistant to glyphosate. Contact your Reapri-Glypho, LLC, representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

Control recommendations for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling or fact sheets for this product and can be obtained from your local retailer or Reapri-Glypho, LLC, representative.

Since the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Reapri-Glypho, LLC, Company is not responsible for any losses that may result from the failure of this product to control glyphosate resistant weeds biotypes.

The following good agronomic practices are recommended to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

6.0 MIXING

Spray solutions of this product should be mixed, stored, and applied using only clean stainless steel, aluminum, plastic, fiberglass, plastic or plastic-lined steel containers. Clean sprayer parts immediately after using this product by thoroughly flushing with water. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations.

6.1 Mixing with Water

NOTE: Product performance may be significantly reduced if water containing soil sediment is used as a carrier. Do not mix this product with water from ponds and ditches that is visibly muddy or murky.

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of clean water. Add the recommended amount of this product near the end of the filling process and mix well. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

6.2 Tank Mixing

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, or dicamba, the user is responsible for ensuring that the specific application is included on the label of the specific products used in the tank mixture. Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions and use according to the most restrictive label requirements. Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

6.3 Tank Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the recommended amount of this product. If needed, add nonionic surfactant before completing the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" and "GENERAL INFORMATION" sections for additional precautions.

6.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product indicated in the following table with water:

Spray Solution

Desired Volume	Amount of GLYPHOSATE 41% SL					
	0.5%	1 %	1.5%	2%	5%	10%
1 gal	0.7 oz	1.3 oz	2 oz	2.7 oz	6.5 oz	13 oz
2.5 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

For backpack, knapsack or pump-up sprayers, it is recommended that the appropriate amount of this product be mixed with water in a larger container, and then used to fill the sprayer.

6.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

6.6 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to spray solutions of this product. However, they can reduce product performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system. Avoid direct application to any body of water.

APPLY SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial application.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversion" sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- Application height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential, NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most *severe* when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sunsets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rates of this herbicide in 3 to 25 gallons of water per acre unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by Reap-Glypho, LLC, for this product.

Drift reduction additives may be used. When a drift reduction additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application --to avoid streaking and uneven or overlapped application, use appropriate marking devices.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Aircraft Maintenance: Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may prevent corrosion.

7.2 Ground Broadcast Equipment

Apply the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label or in separate supplemental labeling or fact sheets published by Reapri-Glypho, LLC. for this product. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

7.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For low-volume directed spray applications, spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

7.4 Selective Equipment

This product may be diluted in water and applied through shielded applicators, hooded sprayers, wiper applicators or sponge bars to weeds listed on this label that are growing in any specified non-crop site.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators and Sponge Bars

Wiper applicators are devices that physically wipe the appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust the height of the applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper applicators when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from the use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators—Solutions ranging from 33 to 75 percent of this product in water may be used.

For Panel Applicators and pressure-feed systems—Solutions ranging from 33 to 100 percent of this product in water may be used.

When applied as recommended, this product **CONTROLS** the following weeds:

Corn, volunteer	Sicklepod
Panicum, Texas	Spanishneedles
Rye, common	Starbur, bristly Shattercane

When applied as recommended, this product **SUPPRESSES** the following weeds:

Beggarweed, Florida	Ragweed, common
Bermudagrass	Ragweed, giant
Dogbane, hemp	Smutgrass
Dogfennel	Sunflower
Guineagrass	Thistle, Canada
Johnsongrass	Thistle, musk
Milkweed	Vaseygrass
Nightshade, silverleaf	Velvetleaf
Pigweed, redroot	

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically recommended.

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 INDUSTRIAL, TURF AND ORNAMENTAL SITES

Unless otherwise specified, applications may be made to control any weeds listed in the "WEEDS CONTROLLED" section of this label. Refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" section for detailed instructions on different application methods.

8.1 Cut Stumps

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100-

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

percent solution of this product to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Saltcedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow Pine, Austrian

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2 Forestry Site Preparation

This product is recommended for the control or partial control of woody brush, trees and herbaceous *weeds* in forestry. This product is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is recommended for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any recommended rate of this product may be used in a tank mix with the following products for forestry site preparation.

Arsenal Applicators Concentrate

Chopper
Escort or Escort XP
Garlon 3A
Garlon 4A
Landmark XP
Oust or Oust XP
Westar

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release, unless otherwise specified on this label or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC, for this product.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

8.3 General Non-crop Areas, Industrial Sites

Use in areas such as airports, apartment complexes commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, rights-of-way, roadsides, shadehouses, sod or turf seed farms, storage areas, sports complexes, substations, turfgrass areas, utility sites, warehouse areas, and wildlife management areas..

General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general non-crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product is registered for use on such non-crop sites. Refer to these products' labels for approved non-crop sites and application rates.

2,4-D	Goal 2XL	Princep™ 4L
Arsenal™	Karmex™ DF	Princep™ DF
atrazine	Krovar™ I DF	Ronstar™ 50WP
Barricade™ 65WG	Landmark II MP	Sahara™
Clarity	Milestone	simazine
Crossbow L	Oust	Surflan™ AS
dicamba	Oust XP	Surflan WDG
diuron	Outrider	Telar™
Endurance™	Pendulum WDG	Transline
Escort™	Pendulum™ 3.3 EC	Vanquish™
Gallery 75 DF	pendimethalin	Velpar DF
Garlon 4	Plateau™	Velpar L
Garlon™ 3A	Poast	

This product plus dicamba tank mixtures may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 1 to 2 quarts of this product plus 2 to 4 ounces of Oust or Oust XP per acre.

Bahiagrass	Johnsongrass
Bermudagrass	Poorjoe
Broomsedge	Quackgrass
Dallisgrass	Vaseygrass
Dock, curly	Vervain, blue Dogfennel
Fescue, tall	

Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Chemical Mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 8 to 64 fluid ounces of this product per acre. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 16 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust or Oust XP in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate was greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

8.4 Habitat Management

Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.5 Hollow Stem Injection

This product may be applied through hand-held injection devices that deliver recommended amounts of this product into targeted hollow-stem plants growing in any non-crop site specified on this label.

For control of the following hollow-stem plants, follow the use instructions below:

Japanese Knotweed, *Polygonum cuspidatum*

Inject 5 mL per stem of this product between second and third internode.

Bohemian Knotweed, *Polygonum bohemicum*

Inject 5 mL per stem of this product between the second and third internode.

Giant Hogweed, *Heracleum mantegazzianum*

Inject one leaf cane per plant 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Poison Hemlock, *Conium maculatum*

Inject one leaf cane per plant 10 to 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Field horsetail, *Equisetum arvense*

Inject one segment above the root crown with 0.5 mL per stem of this product. Use a small syringe that calibrates to this rate.

Canada Thistle, *Cirsium arvense*

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.5 mL per stem of this product is injected into the stem.

Note: The combined total for all treatments must not exceed 10.6 quarts of this product per acre. At 5 mL per stem, 7 quarts should treat approximately 1300 stems per acre.

8.6 Injection and Frill (Woody Brush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 0.04 fluid ounce (1 mL) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50- to 100-percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100-percent concentration of this product. For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:

<u>Control</u>	<u>Partial Control</u>
Oak	Black gum
Poplar	Dogwood
Sweetgum	Hickory
Sycamore	Maple, red

8.7 Non-food Tree, Shrub, or Vine Production Sites

This product may be used for general weed control prior to planting or around established ornamentals, or any woody tree, shrub, or vine species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce or yew, in any production site.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

UNLESS OTHERWISE DIRECTED, THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be taken to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

TYPES OF APPLICATIONS: Site Preparation, Post-directed, Trim-and-edge, Wiper Application Site Preparation

This product may be used prior to planting any tree, shrub, or vine in an ornamental, nursery, or production setting, including Christmas tree species.

Post-directed, Trim-and-edge

This product may be used as a post-directed spray around established woody species, or to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a production setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Wiper Application

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs, or vines. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

8.8 Railroads

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along railroads.

Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

TANK MIXTURES: This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments, provided that the specific product is registered for use on such non-crop sites:

Arsenal	Oust XP
atrazine	Outrider
dicamba	Sahara DG
diuron	simazine
Escort	Spike™ 80 DF
Escort XP	Telar DF
Garlon 3A	Transline
Garlon 4	Vanquish
Hyvar™ X	Velpar DF
Hyvar X-L	Velpar L
Krovar I DF	2,4-D
Oust	

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.75- to 2-percent solution of this product when using high-volume spray-to-wet applications. Apply a 5- to 10-percent solution of this product when using low volume directed sprays for spot treatment.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

TANK MIXTURES: This product may be mixed with the products listed above in this section for enhanced control of woody brush and trees along railroads, provided that the specific product is registered for use on such non-crop sites.

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestein, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestein, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpet creeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue Dogfennel

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may occur.

8.9 Roadsides

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails, Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

TANK MIXTURES: This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments:

atrazine	Outrider
Clarity	Pendulum 3.3 EC
Crossbow L	Pendulum WDG
dicamba	Plateau
diuron	Princep DF
Endurance	Princep 4L
Escort	Ronstar 50WP
Escort XP	Sahara
Gallery 75DF	simazine
Krovar 1 DF	Surflan

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Landmark 11 MP
Landmark MP
Landmark XP
Oust

Telar
Vanquish
2,4-D

See the "MIXING" section of this label for general instructions for tank mixing. **Release of Bermudagrass or Bahiagrass**

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

TANK MIXTURES: This product may also be tank-mixed with Outrider, Oust or Oust XP for residual control. These tank mixtures may delay greenup.

Apply 8 to 64 fluid ounces of this product in a tank mix with 0.75 to 1.3 ounces of Outrider per acre. Read and follow all label directions for Outrider.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust or Oust XP in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Oust or Oust XP per acre on Bermudagrass and no more than 0.5 ounce of Oust or Oust XP per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in 10 to 40 gallons of water per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass
Bluestein, silver
Fescue, tall

Johnsongrass
Trumpet creeper
Vaseygrass

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 9 to 32 fluid ounces of this product with 0.75 to 1.3 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass
Bluestein, silver
Broomsedge
Dallisgrass
Dock, curly
Dogfennel

Fescue, tall
Johnsongrass
Poojoe
Trumpet creeper
Vaseygrass
Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not recommended, since severe injury may occur.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product *per* acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 6 fluid ounces of this product with 0.75 to 2 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height. Use only on well-established bahiagrass.

A tank mixture of this product plus Oust or Oust XP may be used. Apply 6 fluid ounces of this product plus 0.25 ounce of Oust or Oust XP per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.10 Rangelands

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands, pastures, and industrial sites. Preventing weed seed production is critical to the successful control of annual grassy weeds invading these perennial grass sites. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Bromus: This product may be used to control or suppress downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*), cheatgrass (*Bromus secalinus*), cereal rye and jointed goatgrass found in rangelands pastures and industrial sites. Apply 8 to 16 fluid ounces of this product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead: To control or suppress medusahead, apply 16 fluid ounces of this product per acre at the 3-leaf stage when plants are actively growing. Delaying applications beyond this stage will result in reduced or unacceptable control. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses. Applications may be made in the fall or spring.

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. for applications *using* ground equipment, apply in 10 to 20 gallons of water per acre.

Spot Treatment, Wiper Application

This product may be applied in rangeland, pastures or industrial sites as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire site or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acres, no more than 10 percent of the total site maybe treated at any one time. To *achieve* maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

8.11 Utility Sites

In utilities, this product is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product is also recommended for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

Bare Ground, Trim-and-Edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products provided that the specific product used is registered for application on these sites. Refer to the individual product's labels for approved sites and application rates.

2,4-D	Outrider
Arsenal	pendimethalin
atrazine	Plateau
Barricade 65WG	Princep
dicamba	Ronstar 50WP
diuron	Sahara
Endurance	simazine
Escort	Surflan AS
Escort XP	Surflan WDG
Garlon 3A	Telar DF
Garlon 4	Transline
Krenite	Vanguard
Krovar I DF	Velpar DF
Oust	Velpar I.
Oust XP	

Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray incompatibility problems. For side trimming treatments, it is recommended that this product be used alone or in a tank mixture with Garlon 4.

8.12 Conservation Reserve Program (CRP)

This product may be used for renovation (rotating out of CRP), site preparation, postemergence weed control in dormant CRP grasses, or wiper application on CRP land.

Renovation (Rotating out of CRP), Site Preparation

This product may be used to prepare CRP land for crop production. Refer to Federal, State or local use guides for CRP renovation recommendations.

Postemergence Weed Control in Dormant CRP Grasses, Wiper Application

Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Applications may be made using wiper applicators to control tall weeds, or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 8 to 12 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

8.13 Grass Seed or Sod Production

This product may be used in grass seed and sod production for preplant, at-planting, preemergence, removal of established stands, renovation, site preparation, shielded spraying, wiper application, spot treatment, and creating rows in annual ryegrass.

Preplant, Preemergence, At-Planting, Removal of Established Stands, Renovation, Site Preparation

This product controls most existing vegetation for purposes of renovating turf or forage grass seed areas or for establishing turfgrass grown for sod. It may also be used to destroy remaining undesired grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting, or for renovation purposes. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Applications must be made prior to crop emergence in order to avoid crop injury.

Shielded Sprayers

Apply 1 to 3 quarts of this product in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aids in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by the protective shields.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Wiper Application

This product may be applied over the top of desirable grasses using wiper applicators for the control of tall weeds.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Spot Treatment

Apply a 1.5-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

The crop sprayed in the treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

Creating Rows in Annual Ryegrass

Apply 1 to 2 pints of this product per acre. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height. Use the higher rate within the recommended range when ryegrass is greater than 6 inches in height.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Set nozzle heights to allow the establishment of the desired row spacing. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

Take care not to spray or allow spray to drift outside target area in order to avoid unwanted crop destruction. Grower assumes all responsibility for crop losses resulting from misapplication of this product.

8.14 Pastures

This product may be applied to any pasture grass (*Gramineae* family), including bahiagrass, Bermudagrass, bluegrass, brome, fescue, guineagrass, kikuyugrass, orchardgrass, pangola grass, ryegrass, timothy, and wheatgrass. Application can be made as a spot treatment, wiper application, preplant, preemergence, pasture renovation, or postemergent broadcast.

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be applied to control perennial pasture species listed on this label prior to replanting.

if application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment, Wiper Application

This product may be applied in pastures as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper application are made using rates above 3 quarts of this product per acre, no more than 10 percent of the total pasture may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

Postemergent Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts of this product per acre per year onto pasture grasses except for renovation uses as described previously in this section.

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for recommended rates for the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results.

9.1 Annual Weeds

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions. Use the higher rate for tough-to-control species regardless of the weed size at application. Treat tough-to-control weeds early when they are relatively small. This product may be tank mixed provided the tank-mix product is registered for use on the target site. Refer to the individual product labels for approved sites and application rates. For spray-to-wet applications, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. For

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds.

For low volume directed spray applications, use a 4- to 7-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple spouts.

WEED SPECIES

Anoda, spurred	Lamb's-quarters*
Barley*	Little barley*
Barnyardgrass*	London rocket*
Bittercress*	Mayweed
Bassia, fivehook	Medusahead*
Black nightshade*	Morningglory (<i>Ipomoea spp</i>)
Bluegrass, annual*	Mustard, blue*
Bluegrass, bulbous*	Mustard, tansy*
Brome, downy*	Mustard, tumble*
Brome, Japanese*	Mustard, wild*
Browntop panicum*	Oats
Buttercup*	Pigweed*
Carolina foxtail*	Plains/Tickseed coreopsis*
Carolina geranium Castor bean	Prickly lettuce*
Cheatgrass*	Puncturevine
Cheeseweed (<i>Malva parviflora</i>)	Purslane, common
Chervil*	Ragweed, common*
Chickweed*	Ragweed, giant
Cocklebur*	Red rice
Copperleaf, hophornbeam	Russian thistle
Corn*	Rye*
Corn speedwell*	Ryegrass*
Crabgrass*	Sandbur, field*
Dwarf dandelion*	Shepherd's-purse*
Eastern manna grass*	Sicklepod
Eclipta*	Signalgrass, broadleaf*
Fall panicum*	Smartweed, ladythumb*
Falsedandelion*	Smartweed, Pennsylvania*
Falseflax, smallseed*	Sowthistle, annual
Fiddleneck	Sowthistle, annual
Field pennycress*	Spanishneedles
Filaree	Speedwell, purslane*
Fleabane, annual*	Sprangletop*
Fleabane, hairy (<i>Conyza bonariensis</i>)*	Spurge, annual
Fleabane, rough*	Spurge, prostrate*
Florida pulley	Spurge, spotted*
Foxtail*	Spurry, umbrella*
Goatgrass, jointed*	Starthistle, yellow
Goosegrass	Stinkgrass*
Grain sorghum (milo)*	Sunflower*
Groundsel, common*	Teaweed/ Prickly sida
Hemp sesbania	Texas panicum*
Henbit	Velvetleaf
Horseweed/Marestail (<i>Conyza canadensis</i>)	Virginia copperleaf
Itchgrass*	Virginia pepperweed*
Johnsongrass, seedling	Wheat*

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Junglerice	Wild oats*
Knotweed	Witchgrass*
Kochia	Woolly cupgrass*
	Yellow rocket

*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 1 pint of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure through coverage of foliage and treat when weeds are in an early growth stage.

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the recommended range.

Ensure thorough coverage when using spray-to-wet treatments with hand-held equipment. For best results, use a 1.5-percent solution on harder-to-control perennials such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

For low volume directed spray applications, use a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Allow 7 or more days after application before tillage.

WEED SPECIES	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2 - 4	1 - 2
Bahiagrass	3 - 5	2
Beachgrass, European (<i>Ammophila arenaria</i>)		5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4 - 5	2
Bluegrass, Kentucky	2	2
Blueweed, Texas	4 - 5	2
Brackenfern	3 - 4	1 - 1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf		2
Canarygrass, reed	2 - 3	2
Cattail	3 - 5	2
Clover; red, white	3 - 5	2
Cogongrass	3 - 5	2
Dallisgrass	3 - 5	2
Dandelion	3 - 5	
Dock, curly	3 - 5	2
Dogbane, hemp	4	2
Fescue (except tall)	3 - 5	2
Fescue, tall	1 - 3	2
German ivy	2 - 4	1 - 2
Guineagrass	3	1
Horsenettle	3 - 5	2
Horsradish	4	2
Iceplant	2	1.5-2

Jerusalem artichoke	3 - 5	2
Johnsongrass	2 - 3	1
Kikuyugrass	2 - 3	2
Knapweed	4	2
Lantana		1 - 1.25
Lespedeza	3 - 5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3 - 5	2
Napiergrass	3 - 5	2
Nightshade, silverleaf	2	2
Nutsedge: purple, yellow	3	1 - 2
Orchardgrass	2	2
Pampasgrass	3 - 5	1.5 - 2
Paragrass	3 - 5	2
Pepperweed, perennial	4	2
Phragmites*	3 - 5	1 - 2
Poison hemlock	2 - 4	1 - 2
Quackgrass	2 - 3	2
Reed, giant	4 - 5	2
Ryegrass, perennial	2 - 3	1
Smartweed, swamp	3 - 5	2
Spurge, leafy*	-	2
Sweet potato, wild*	-	2
Thistle, artichoke	2 - 3	1 - 2
Thistle, Canada	2 - 3	2
Timothy	2 - 3	2
Torpedograss*	4 - 5	2
Trumpetcreeper*	2 - 3	2
Vaseygrass	3 - 5	2
Velvetgrass	3 - 5	2
Wheatgrass, western	2 - 3	2

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In and areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

For best results when using hand-held equipment, use a 1.5-percent solution on harder-to-control woody brush and trees. For low volume directed-spray applications, apply a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50-percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple spouts.

Symptoms may not appear prior to frost or senescence with fall treatments.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

WEED SPECIES	Broadcast Rate (Q T / A)	Hand - Held Spray-to-Wet % Solution
Alder	3 - 4	1 - 1.5
Ash*	2 - 5	1 - 2
Aspen, quaking	2 - 3	1 - 1.5
Bearclover (Bearmat)*	2 - 5	1 - 2
Beech*	2 - 5	1 - 2
Birch	2	1
Blackberry	3 - 4	1 - 1.5
Blackgum	2 - 5	1 - 2
Bracken	2 - 5	1 - 2
Broom; French, Scotch	2 - 5	1.5 - 2
Buckwheat, California*	2 - 4	1 - 2
Cascara**	2 - 5	1 - 2
Catsclaw*		1 - 1.5
Ceanothus*	2 - 5	1 - 2
Chamise*	2 - 5	1
Cherry; bitter, black, pin	2 - 3	1 - 1.5
Coyote brush	3 - 4	1.5 - 2
Deerweed	2 - 5	1
Dogwood*	2 - 5	1 - 2
Elderberry	2	1
Elm*	2 - 5	1 - 2
Eucalyptus	--	2
Gorse*	2 - 5	1 - 2
Hasardia*	2 - 4	1 - 2
Hawthorn	2 - 3	1 - 1.5
Hazel	2	1
Hickory*	2 - 5	1 - 2
Honeysuckle	3 - 4	1 - 1.5
Hornbeam, American*	2 - 5	1 - 2
Kudzu	4	2
Locust, black*	2 - 4	1 - 2
Madrone resprouts*	--	2
Manzanita*	2 - 5	1 - 2
Maple, red	2 - 4	1 - 1.5
Maple, sugar	--	1 - 1.5
Monkey flower**	2 - 4	1 - 2
Oak; black, white*	2 - 4	1 - 2
Oak, post	3 - 4	1 - 1.5
Oak; northern, pin	2 - 4	1 - 1.5
Oak, Scrub*	2 - 4	1 - 1.5
Oak; southern red	2 - 3	1 - 1.5
Peppertree, Brazilian (Florida holly)*	2 - 5	1 - 2
Persimmon*	2 - 5	1 - 2
Pine	2 - 5	1 - 2
Poison ivy	4 - 5	2
Poison oak	4 - 5	2
Poplar, yellow*	2 - 5	1 - 2

Redbud, eastern	2 - 5	1 - 2
Rose, multiflora	2	1
Russian olive*	2 - 5	1 - 2
Sage, black	2 - 4	1
Sage, white*	2 - 4	1 - 2
Sage brush, California	2 - 4	1
Salmonberry	2	1
Saltcedar*	2 - 5	1 - 2
Sassafras*	2 - 5	1 - 2
Sourwood*	2 - 5	1 - 2
Sumac; laurel, poison, smooth, sugarbur	2 - 4	1 - 2
Sweetgum	2 - 3	1 - 1.5
Swordfern*	2 - 5	1 - 2
Tallowtree, Chinese	—	1
Tan oak resprouts*	-	2
Thimbleberry	2	1
Tobacco, tree*	2 - 4	1 - 2
Toyon*	-	2
Trumpet creeper	2 - 3	1 - 1.5
Vine maple*	2 - 5	1 - 2
Virginia creeper	2 - 5	1 - 2
Waxmyrtle, southern*	2 - 5	1 - 2
Willow	3	1
Yerbasanta*	-	2

10.0 LIMIT OF WARRANTY AND LIABILITY

Repar-Glypho, LLC, warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the fullest extent permitted by law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, TO THE FULLEST EXTENT PERMITTED BY LAW. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

II. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Trademarks are the property of their respective owners. EPA Reg. No. 86004-

Repar-Glypho, LLC.

Silver Springs, MD 20910

III. SUPPLEMENTAL LABELING

Table of Contents: Supplemental labeling

	Name
A	FOR USE IN DORMANT ALFALFA
B	FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER FIRST CUTTING
C	FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS
D	FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARESTAIL, <i>Conyza canadensis</i>) IN COTTON, CORN, AND SOYBEANS
E	HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROL IN ROUNDUP READY CANOLA
F	FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRENDO COUNTY
G	FOR AERIAL APPLICATION IN ARKANSAS ONLY
H	FOR CONTROLLING BARNYARDGRASS (ECHINOCHLOA CRUS-GALLI) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY
I	FOR WEED CONTROL APPLICATION IN SEED PRODUCTION OF ALFALFA WITH THE ROUNDUP READY [®] GENE
J	FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY
K	BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS
L	FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD [®] TALL FESCUE AND AURORA GOLD [®] FINE FESCUE SELECTIONS
M	GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS
N	FOR GROUND AND AERIAL APPLICATION TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY
O	GLYPHOSATE 41% SL HERBICIDE FOR CONIFER RELEASE

III. SUPPLEMENTAL LABELING

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **GLYPHOSATE 41% SL** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **GLYPHOSATE 41% SL** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

GLYPHOSATE 41% SL Herbicide

EPA Reg. No. 86004-

Keep out of the reach of children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any other manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIREABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for **GLYPHOSATE 41% SL** for essential product performance information.

[INSERT SPECIFIC DIRECTIONS FOR USE TEXT FROM THE FOLLOWING SECTIONS]

Read the "Limit of Warranty and Liability" in the label booklet for **GLYPHOSATE 41% SL** before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

REPAR-GLYPHO, LLC. COMPANY
SILVER SPRING, MD 20910

III. SUPPLEMENTAL LABELING

A. FOR USE IN DORMANT ALFALFA

DIRECTIONS FOR USE

This product will control or suppress many weeds, including quackgrass, downy brome, and cheatgrass in dormant alfalfa.

Apply 8 to 12 fluid ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with **GLYPHOSATE 41% SL**.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product is limited to persons who have attended a Repar-Glypho, LLC.-approved training program. Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

B. FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER THE FIRST CUTTING

DIRECTIONS FOR USE

This product may be applied at 16 fluid ounces per acre to control the weeds listed below and most other winter annual grass and broadleaf weeds in established coastal Bermudagrass pastures.

Annual bluegrass	Oats
Cheat	Ryegrass, Italian
Crabgrass	Sandbur, field
Henbit	Sunflower
Johnsongrass, seedling	Wheat
Little barley	Wild mustard

TIMING OF APPLICATION

Applications prior to spring growth: Apply this product in

either late winter or early spring but before new coastal Bermudagrass growth begins in the spring. Applications to new growth can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 60 days after making this application before grazing or harvesting the treated area.

Application following the first cutting: Apply this product after the first Bermudagrass cutting when the Bermudagrass has not yet begun to regrow. Applications made after regrowth has begun can damage the Bermudagrass.

III. SUPPLEMENT LABELING

Remove domestic livestock from the pasture before making the application. Wait 28 days after making this application before grazing or harvesting the treated area.

NOTE: ONLY ONE APPLICATION PER YEAR MAY BE MADE TO ANY ONE FIELD. A SPRING APPLICATION PRIOR TO GROWTH AND AN APPLICATION FOLLOWING THE FIRST CUTTING MAY NOT BE MADE ON THE FIELD DURING THE SAME YEAR.

C. FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS

DIRECTIONS FOR USE

Refer to the **GLYPHOSATE 41% SL** label for rate recommendations and weeds controlled.

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 2 to 3 gallons of water per acre.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, draft, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray to drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as line spray droplets.

In aerial applications, do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Ensure uniform application. Use appropriate marking devices when applying herbicides by air.

Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR IS MOST SUSCEPTABLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

D. FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARETAIL, <i>Conyza Canadensis</i>) IN CROP AND NON-CROP AREAS.

III. SUPPLEMENTAL LABELING

DIRECTIONS FOR USE

For ground applications, use 10 to 20 gallons of water per acre. For aerial applications, use 3 to 15 gallons of water per acre. **For tank-mix recommendations, read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental or fact sheets published separately for all herbicides used.**

Cotton:

Preplant:

For the control of horseweed, apply this product (32 fluid ounces per acre) in a tank-mix with Clarity® (8 fluid ounces per acre). This application must be made 21 to 35 days before planting and before horseweed reaches 6 inches in height. In order to avoid crop injury, a minimum interval of 21 days during which there is at least 1 inch cumulative rainfall must be observed between Clarity application and planting of cotton.

2,4-D may be included in the tank-mixture with this product. Refer to the 2,4-D product label for the time intervals that are required between application and planting and other geographic use restrictions.

Post-directed (Roundup Ready® Cotton and Roundup Ready Flex cotton® varieties only):

Management of early season weed competition and the development of a crop height differential between cotton and horseweed is often achieved by a combination of preplant burndown and post emergence over-the-top and/or directed applications of this product. These measures enhance the development of a height differential that is necessary to successfully make post-directed treatments. In-crop post-directed applications of MSMA (2 pounds active ingredient per acre) tank-mixed with diuron (0.5 to 0.75 pounds active ingredient per acre) should be made when the temperature is 80°F or higher.

Soybeans:

Preplant:

It is strongly encouraged that horseweed be controlled prior to planting. Apply a tank mixture of this product (32 fluid ounces per acre) with 2,4-D (0.5 pounds a.i. per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

In-crop (Roundup Ready Soybean varieties only):

This treatment should be used as a salvage treatment only for a horseweed infestation that was not controlled prior to planting. At the time of treatment, horseweed should not exceed 6 inches in height. Apply a tank mixture of this product (32 fluid ounces per acre) with FirstRate® (0.3 ounces per acre). Application should be made between full emergence of the first trifoliate leaf and 50% flowering stage of soybeans.

Corn:

Preplant, At-Planting, Preemergence:

Apply a tank mixture of this product (32 fluid ounces per acre) plus 2,4-D (0.5 pounds i.e. or per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Atrazine (1 to 2 pounds active ingredients per acre) may be included in the tank-mixture to provide residual control. Refer to the atrazine product label for specific use instructions.

In-crop (Roundup Ready Corn hybrids only):

Apply a tank-mixture of this product (32 fluid ounces per acre) plus Clarity (8 to 16 ounces per acre) or 2,4-D (0.5 to 1.0 pounds of inactive ingredients per acre) between corn emergence and the 5-leaf stage of growth (approximately 8 inches tall).

III. SUPPLEMENTAL LABELING

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Tree and Vine Crops:

Orchards (Pome Fruit, Stone Fruit and Tree Nuts).

Apply two quarts of this product plus 2,4-D (18 fluid ounces per acre Dri-Clean® or 2 pints per acre Orchard Master® CA) before marestrail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is recommended. Follow labeling for 2,4-D product with regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply.

Vine Crops (grapes only):

Apply 2 quarts of this product per acre plus 2,4-D (18 fluid ounces per acre Dri-Clean®) before marestrail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is recommended. Follow Dri-Clean® labeling in regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply. Also, residual herbicides such as diuron may provide additional preemergence control.

General Non-crop Areas:

Apply a tank mixture of this product (1 quart per acre) before horseweed exceeds 6 inches in height. Control may be enhanced by making applications when horseweed is still in the rosette stage of growth. This product may be tank-mixed with the following products provided that the specific product is registered for use on the target site. Refer to these product labels for approved sites and application rates. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive precautionary statements for each product in the mixture.

- 2,4-D
- Crossbow™L
- Dicamba
- Gallery™ 75 DF
- Krova® 1 DF
- Landmark™ II MP
- Landmark™ MP
- Milestone™
- Overdrive®
- Telar™ DF
- Vanquish®
- Velpar® DF

E. HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROLS IN ROUNDUP READY CANOLA
--

DIRECTIONS FOR USE

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A ROUNDUP READY GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELEWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA.

III. SUPPLEMENTAL LABELING

Maximum Allowable Combined Application Quantities Per Season	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application for emergence to 6-leaf stage	2 quarts per acre

Preplant, At-planting Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Weeds Controlled: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" in this label

Single Application- Apply 16 to 32 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application- Apply 32 fluid ounces per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but not later than the 6-leaf stage. Sequential applications are recommended for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass or when controlling weeds with multiple application times.

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in the Roundup Ready crops. No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 64 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

F. FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRESNO COUNTY, CALIFORNIA

DIRECTIONS FOR USE

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this supplemental labeling and in the product label booklet. Refer to Aerial Equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the product label for additional information. Refer to the individual use site section of the product label, or to other supplemental labeling or technical fact sheets published separately for this product by Repar-Glypho, LLC., for specific use instructions.

AVOID DRIFT-- DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetations after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops:

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).

III. SUPPLEMENTAL LABELING

3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.
5. Apply by air only to nonresidential areas.

When applied as recommended under the conditions described, this product controls annual and perennial weeds as listed in the label booklet.

When tank mixing product with 2,4-D, only 2,4-D amine formulations may be used for aerial applications in California. Tank mixtures 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and alfalfa and pasture renovation applications only. This product, when tank mixed with dicamba, may not be applied by air in California.

ADDITIONAL INFORMATION FOR FRESNO COUNTY, CALIFORNIA

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California.

North:	Fresno County line
South:	Fresno County line
East:	State Highway 99
West:	Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Recommendations

A written recommendation MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation MUST state the proximity or surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to Fresno County Agricultural Commissioner.

Applications at Night- Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

III. SUPPLEMENTAL LABELING

Note: For aerial application from April 1 through February 14, refer to the other sections of this supplemental label.

G. FOR AERIAL APPLICATION IN ARKANSAS ONLY.

DIRECTIONS FOR USE

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Do not apply this product when winds are in excess of 10 mph.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 mph is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 mph toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

H. FOR CONTROLLING BARNYARDGRASS (*ECHINOCHLOA CRUS-GALLI*) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY

DIRECTIONS FOR USE

Renovation Treatment

USE INSTRUCTIONS: This product may be applied as a renovation treatment in rice crops to control barnyardgrass infestations using ground broadcast spray or hand-held equipment. Renovation is defined as

III. SUPPLEMENTAL LABELING

herbicide treatment that will produce crop and weed destruction in an entire field or contiguous area treated within a field. Follow the application methods and recommended treatment rates in the label booklet for **GLYPHOSATE 41 % SL** herbicide.

PRECAUTIONS, RESTRICTIONS: Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction. The rice straw and stubble from the treated area, including a 25-foot buffer zone on all sides, shall not be used for grazing, animal bedding or any feed purposed.

Aerial applications are not permitted for rice renovation using this supplemental label.

I. FOR WEED CONTROL APPLICATIONS IN SEED PRODUCTION OF ROUNDUP READY® ALFALFA
--

General Information

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FIRST OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for GLYPHOSATE 41% SL for essential product performance information.

The Roundup Ready designation indicates that the alfalfa contains a patented gene that provides tolerance to this product. Information on Roundup Ready Alfalfa varieties may be obtained from your seed supplier or Repar-Glypho, LLC, representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

See the "ROUNDUP READY CROPS" section of the GLYPHOSATE 41% SL label booklet for general precautionary instructions for use in Roundup Ready crops. Do NOT combine the instructions in this supplemental label with other recommendations listed in the "PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS" section of the **GLYPHOSATE 41 % SL** label booklet intended for alfalfa varieties that do not contain a glyphosate tolerance gene.

DIRECTIONS FOR USE

This product will control many troublesome emerged weeds with over-the-top applications in Roundup Ready alfalfa grown for seed. Over-the-top applications may be made from emergence through the late vegetative stage, and spot treatments may be made from early bud stage through seed harvest.

For ground applications with broadcast equipment, apply this product in 3 to 40 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial application: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre.

DO NOT EXCEED 2 QUARTS OF THIS PRODUCT PER ACRE WHEN MAKING APPLICATIONS BY AIR. FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATION IN THAT STATE. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

III. SUPPLEMENTAL LABELING

See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready alfalfa. Follow the cleaning procedures specified on the label of the product(s) used. Alfalfa can be very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence and Post-harvest of seed

Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Total Preplant, At-planting and Preemergence Applications	2 quarts per acre
Total in-crop application rate from emergence through the late vegetative stage*	6 quarts per acre
Spot-treatment during early bud stage through seed harvest (See the "Spot Treatment after late vegetative stage" section and the "PRECAUTIONS AND RESTRICTIONS" section of this label for complete instructions)	Apply spray-to-wet; do not apply to the point of runoff

There are no rotational crop restrictions following applications of this product. For any crop NOT listed in the label booklet, applications must be at least 30 days prior to planting.

Over-the-top applications: Broadcast applications of this product may be made using ground or aerial application equipment over the top of Roundup Ready alfalfa from emergence through the late vegetative stage. Do not make broadcast applications of this product between the initiation of alfalfa budding and the harvest of seed. Any single over-the-top broadcast application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart.

Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings are susceptible and will not survive or thrive after the first application of this product. A single application of at least 1 quart per acre of this product should be applied at or before the 3 to 4 trifoliate growth stage to eliminate the effects of stand gaps created by the loss of non-Roundup Ready plants.

Stop Treatment after late vegetative stage: For late emerging weeds, this product may be applied as a spot treatment in Roundup Ready alfalfa grown for seed, during the early bud stage through seed harvest. Applications made during this stage may result in reduced seed yield and quality and are the responsibility of the grower. Make applications on a spray-to-wet basis. Do not spray to the point of runoff. If a spot treatment is made after the late vegetative stage, harvested seed must not be used for alfalfa sprout production.

Post-harvest applications: Following harvest of Roundup Ready alfalfa seed, the stand may be managed for forage and hay production. Refer to the Roundup Ready alfalfa section of the label booklet **GLYPHOSATE 41% SL** for rates and timing of applications for forage and hay production.

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" in the **GLYPHOSATE 41% SL** label booklet. Some weeds with multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some re-growth of weeds has occurred.

In addition to those weeds listed in the **GLYPHOSATE 41% SL** label booklet, this product will suppress or control the parasitic weed, Dodder (*Cuscuta spp.*) in Roundup Ready alfalfa seed production. Repeat applications may be necessary for complete control.

III. SUPPLEMENT LABELING

Tank mixtures with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control and are NOT recommended for over-the-top applications of this product.

PRECAUTIONS AND RESTRICTIONS: Do not make over-the-top broadcast applications of this product between the initiation of alfalfa budding and the harvest of Roundup Ready alfalfa seed. The use of harvested Roundup Ready alfalfa seed is not suitable for, and is not recommended for, production of alfalfa sprouts.

Read the "LIMIT OF WARRANTY OR LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using. For (in-crop) over-the-top uses on Roundup Ready crop varieties, crop safety and weed control performance are not warranted by Repar-Glypho, LLC, when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

J. FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY

DIRECTIONS FOR USE

GENERAL INFORMATION

The use of the over-the-top applications described in this label on *other* than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (nodes) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this supplemental label are specific to, and should only be used with varieties designated as **Round Ready Flex cotton**. **DO NOT** combine the instructions in this supplemental label with those in the "Roundup Ready Cotton" or "Roundup Ready Flex Cotton" sections found in the label booklet for **GLYPHOSATE 41% SL**, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instruction on labeling for this or other glyphosate-containing product. See "Maximum Use Rate" in the "GENERAL INFORMATION" section of the label booklet for **GLYPHOSATE 41% SL** for additional information.

ROUNDUP READY FLEX COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATED THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop), Preharvest

Maximum Allowable Combined Application Quantities Per Season

Combined total per year for all application	8 quarts per acre
---	-------------------

Calculate the combined rate to be used for all preplant, in-crop and preharvest applications, to ensure that the total does not exceed the maximum allowed rate per acre year shown above.

Preplant, At-planting, Preemergence applications	5 quarts per acre
--	-------------------

Total in-crop applications from ground cracking to 60% percent open bolls	6 quarts per acre
---	-------------------

Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 fluid ounces per acre
---	-------------------------

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of the **GLYPHOSATE 41% SL** label booklet for general precautionary instructions for use in Roundup Ready crops.

Preplant, At-planting, Preemergence

III. SUPPLEMENTAL LABELING

USE INSTRUCTIONS: This product may be applied before, during or after planting Roundup Ready Flex cotton.

Postemergence (In-crop)

USE INSTRUCTIONS: When applied in accordance with this label, GLYPHOSATE 41% SL will control labeled annual grasses and broadleaf weeds in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds is recommended. This product may be applied by ground application equipment at rates up to 2 quarts per acre per application postemergence to Roundup Ready Flex cotton. In addition to broadcast applications, post-directed equipment may be used to achieve weed coverage.

NOTE: For specific rates of application and instructions, refer to the "ANNUAL" and "PERENNIAL WEEDS RATE TABLES" in the label booklet for **GLYPHOSATE 41% SL**.

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 2 quarts per acre made using ground application equipment. **In-crop application rates above 1 quart per acre made alone, or with the addition of other crop chemical products containing surfactant, may cause a crop response, including leaf speckling or leaf necrosis.** Do not exceed a maximum rate of 3 pints per acre of this product when making applications by air. Between layby and 60 percent open bolls, the maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre.

Preharvest

USE INSTRUCTIONS: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready Flex cotton after 60 percent boll crack. Up to 2 quarts of this product may be applied using either aerial or ground spray equipment. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of Roundup Ready Flex cotton. Do not apply this product over-the-top beyond first bloom to cotton grown for seed.

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY FLEX COTTON. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

K. BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS.
--

DIRECTIONS FOR USE

NOTE: IF IMPROPERLY APPLIED, THIS PRODUCT HAS THE POTENTIAL TO CAUSE SEVERE CHRISTMAS TREE INJURY. FOLLOW ALL LABEL DIRECTIONS.

This product may be applied as a broadcast spray over established Christmas trees. Ensure that adequate buffers are maintained to prevent drift onto nearby desirable crops or vegetation. Read the entire "APPLICATION"

III. SUPPLEMENTAL LABELING

"EQUIPMENT AND TECHNIQUES" section of the GLYPHOSATE 41% SL label booklet for additional application precautions

This application is approved for the following Christmas tree species:

Douglas fir (*Pseudotsuga menziesii*)

Fir species (*Abies* spp.)

Spruce species (*Picea* spp.)

Applications may be made only after trees have completed at least a full growing season since planting or transplanting. Applications should not be made within 1 full year prior to tree harvest.

Applications may only be made in the fall after the formation of final conifer resting buds. Final resting buds must be fully hardened and in the dormant stage. Applications made at any other time may result in unacceptable Christmas tree injury.

Avoid spray pattern overlap, as injury may occur.

Apply 1 quart of this product per acre in 5 to 30 galls of water per acre.

NOTE: DO NOT ADD SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO THIS PRODUCT AS SEVERE CHRISTMAS TREE INJURY MAY RESULT.

This product may be used at rates from 1 to 2 quarts per acre in some areas. Consult your local Repar-Glypho, LLC, representative or Roundup supplier for specific recommendations if you require rates greater than 1 quart per acre.

Drift control additives may increase Christmas tree injury and their use is not recommended.

The use of other herbicides tank mixed with this product is not recommended since severe Christmas tree injury may result.

L. FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

GLYPHOSATE 41% SL Tolerant Tall Fescue Selections For Seed Protection

Use this product on [INSERT BRAND NAME] tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 4 to 16 fluid ounces per acre as a postemergence spray on **GLYPHOSATE 41% SL** tolerant tall fescue selections. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass, mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the **GLYPHOSATE 41% SL** label booklet for a complete list of weeds controlled or suppressed.

NOTE: The recommended rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with **GLYPHOSATE 41% SL** tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

Timing of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy in the Spring after dormancy break until 60 days prior to harvest.

III. SUPPLEMENTAL LABELING

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

M. GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the GLYPHOSATE 41% SL label booklet for rate recommendations. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quart per acre of this product may be used for improved results.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any recommended rate of this product may be used in a tank mixture with these products.

<u>Tank-mix Product</u>	<u>Rate per Acre</u>
Arsenal TM *	0.5 to 4 pints
Barvel	1 to 4 pints
2, 4-D	0.5 to 1 pound
Garlon TM 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar TM X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar TM I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust TM	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike TM 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

*Arsenal is not approved for use in the state of California

III. SUPPLEMENTAL LABELING

Refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mixture products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

When used in combination as recommended by Repar-Glypho, LLC, Company, the liability of Repar-Glypho, LLC, shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Repar-Glypho, LLC, product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

TMArsenal is a trademark of American Cyanamid Company.

TMBanvel is a trademark of Sandoz Crop Protection Corporation.

TMGarlon and Spike are trademarks of Dow AgroSciences LLC.

TMHyvar, Krovar and Oust are trademarks of E. I DuPont de Nemours and Company.

N. FOR GROUND AERIAL APPLICATIONS TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY.
--

DIRECTIONS FOR USE

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Nonionic surfactants which are labeled for use with herbicides may be used to improve wetting of foliage. Do not reduce rates of GLYPHOSATE 41% SL when adding surfactant. Read and carefully observe surfactant rates, cautionary statements, and other information appearing on the surfactant label.

TIMING OF APPLICATION: Apply this product as a broadcast spray when plants are actively growing for partial control of undesirable vegetation listed on this label. Best results are obtained when application is made in the spring to early summer when brush species are at a high moisture content and flowering.

This product may be used as recommended for:

- Aid to burning treatment to establish and maintain fuel breaks
- Establishing fire perimeters and black lines
- Aid to prescribed burning
- Along fire roads and rights-of-way

APPLICATION RECOMMENDATION: Apply 2 quarts of this product per acre for partial control of the following emerged brush and chaparral species:

III. SUPPLEMENTAL LABELING

Ceanothus	Sage
Ceanothus spp.	Salvia spp.
Chamise	Scrub oak
Adenostroma fasciculatum	Quercus dumosa

Ground applications should be applied in 3 to 40 gallons of total spray solution per acre.

Aerial applications (helicopter only) should be applied in 3 to 15 gallons of total spray solution per acre.

Avoid direct application to any body of water.

AVOID DRIFT- DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

For aerial application of this product, please see the supplemental label directions for serial application in California.

O. GLYPHOSATE 41 % SLHERBICIDE FOR CONIFER RELEASE

DIRECTIONS FOR USE

AERIAL APPLICATION

This product may be applied using aerial spray equipment for conifer release treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of the label booklet for GLYPHOSATE 41% SLherbicide for information on how to properly spray this product by air.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

CONIFER RELEASE

For release, apply only where conifers have been established for more than one year. Vegetation should not be disturbed prior to treatment or until symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of the label booklet for GLYPHOSATE 41% SLherbicide.

For release of the following conifer species:

Douglas Fir	Pines*
Pseudotsuga menziesii	Pinus spp.
Fir	Spruce
Abies spp.	Picea spp.
Hemlock	
Tsuga spp.	

*Includes all species except eastern white pine, loblolly pine or slash pine.

III. SUPPLEMENTAL LABELING

Apply 1.5 to 2 quarts of this product per acre except in Washington and region, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of this product per acre before any major leaf drop of deciduous species.

For release of western hemlock apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly Pine

Pinus taeda

Eastern white pine

Pinus strobus

Slash Pine

Pinus elliotti

Late Season Application – Apply 1.5 to 2 quarts of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will result in potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later application. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label direction will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash

Fraxinus spp.

Cherry, black

Prunus serotina

Cherry, pin

Prunus pensylvanica

Elm

Ulmus spp.

Hawthorn

Crataegus spp.

Locust, black

Robinia pseudoacacia

Maple, red

Acer rubra

Oak, black

Quercus velutina

Oak, post

Quercus stellata

Oak, southern red

Quercus falcate

Oak, white

Quercus alba

Persimmon

Diospyros spp.

Poplar, yellow

Liriodendron tulipifera

Sassafras

Sassafras albidum

Sourwood

Oxydendrum arboreum

Sumac, poison

Rhus vernix

Sumac, smooth

Rhus glabra

Sumac, winged

Rhus copallina

Sweetgum

Liquidambar styraciflua

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

GLYPHOSATE 41% SL HERBICIDE PLUS OUST™ TANK MIXTURES FOR CONIFER RELEASE FROM HERBACEOUS WEEDS

To release loblolly pines from herbaceous weeds, tank mixtures of this product with Oust will provide control of annual weeds listed in the "WEEDS CONTROLLED" section of the label booklet for GLYPHOSATE 41% SL herbicide and the Oust label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of GLYPHOSATE 41% SL herbicide with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines.

This product plus Oust tank mixtures may not be applied by air in California.

III. SUPPLEMENTAL LABELING

This tank mixture may be applied using aerial equipment. When applying by air, use the recommended rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use the low rates of both products. Use the higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	Johnsongrass**
<i>Paspalum notatum</i>	<i>Sorghum halepense</i>
Broomsedge	Poorjoe**
<i>Andropogon virginicus</i>	<i>Diodia teres</i>
Dock, curly	Trumpet creeper*
<i>Rumex crispus</i>	<i>Campsis radicans</i>
Dogfennel	Vaseygrass
<i>Eupatorium capilliflorum</i>	<i>Paspalum urvillei</i>
Fescue, tall	Vervain, blue
<i>Festuca arundinacea</i>	<i>Verbena hastata</i>
*Suppression at the higher rates only	
**Control at the higher rates	

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water, insects or disease.

Read and observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Oust is a trademark of E.I. DuPont de Nemours and Company.

MASTER LABEL FOR EPA REG. NO. 86004-U

Registered Brand Names:
GLYPHOSATE 41% SL

Table of Contents for Master Label

	Main Label for Food Crop Uses	Page 2
	Main Label for Industrial, Turf, & Ornamental Uses	Page 70
	Supplemental Labeling	Page 100

See each label part for more detailed table of contents

I. MAIN LABEL FOR FOOD CROP USES

GLYPHOSATE 41% SL

Herbicide

Complete Directions for Use EPA Reg. No. 86004-U

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY[®] CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Herbicide for Roundup Ready[®] Crops.

Selective broad-spectrum weed control in Roundup Ready crops. Non-selective, broad-spectrum weed control for many agricultural systems and farmsteads.

Not all products on this label are registered for use in California. Check the registration status of each product in California before using.

Read the entire label before using this product.

Use only according to label instructions.

[Container Label Statement (5 gallons or smaller):]

Do not use this container for residential treatments.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC REPACKAGING OR TOLL REPACKAGING AGREEMENT.

Non-refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

CONTENTS

1.	1.0	INGREDIENTS
2.	2.0	IMPORTANT PHONE NUMBERS
3.	3.0	PRECAUTIONARY STATEMENTS
	3.1	Hazards to Humans and Domestic Animals
	3.2	Environmental Hazards
	3.3	Physical or Chemical Hazards
4	4.0	STORAGE AND DISPOSAL
5	5.0	GENERAL INFORMATION (How This Product Works)
6	6.0	Weed Resistance Management
	6.1	General Weed Management Recommendations
	6.2	Management Recommendations for Glyphosate Resistant Biotype
7	7.0	MIXING
	7.1	Mixing with Water
	7.2	Tank Mixtures
	7.3	Tank Mixing Procedures
	7.4	Mixing for Hand-Held Sprayers
	7.5	Surfactants
	7.6	Ammonium Sulfate
	7.7	Colorants or Dyes
	7.8	Drift Reduction Additives
8	8.0	APPLICATION EQUIPMENT AND TECHNIQUES
	8.1	Aerial Equipment
	8.2	Ground Broadcast Equipment
	8.3	Hand-Held or Backpack Equipment
	8.4	Selective Equipment
	8.5	Injection Systems 8.6 CDA Equipment
9	9.0	ANNUAL AND PERENNIAL CROPS (Alphabetical)
	9.1	Cereal and Grain Crops
	9.2	Corn
	9.3	Cotton
	9.4	Fallow Systems
	9.5	Grain Sorghum (Milo)
	9.6	Herbs and Spices
	9.7	Oil Seed Crops
	9.8	Soybeans
	9.9	Sugarcane
	9.10	Vegetable Crops 9.11 Miscellaneous Crops
10	10.0	TREE, VINE, AND SHRUB CROPS (Alphabetical)
	10.1	Berry Crops
	10.2	Citrus
	10.3	Miscellaneous Tree Food Crops
	10.4	Non-Food Tree Crops
	10.5	Pome Fruit

- 10.6 Stone Fruit
- 10.7 Tree Nuts
- 10.8 Tropical and Subtropical Trees and Fruits
- 10.9 Vine Crops
- 11 11.0 **PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS**
 - 11.1 Alfalfa, Clover, and Other Forage Legumes
 - 11.2 Conservation Reserve Programs (CRP)
 - 11.3 Grass Seed or Sod Production
 - 11.4 Pastures
 - 11.5 Rangelands
- 12 12.0 **ROUNDUP READY® CROPS**
 - 12.1 Roundup Ready® Alfalfa
 - 12.2 Roundup Ready® Canola (Spring Varieties)
 - 12.3 Roundup Ready® Canola (Winter Varieties)
 - 12.4 Roundup Ready® Corn 2 (Supplemental Labeling for Corn 2)
 - 12.5 Roundup Ready® Gene Cotton
 - 12.6 Roundup Ready® Flex Cotton
 - 12.7 Roundup Ready® Gene Soybeans
 - 12.8 Roundup Ready® Gene Sugar Beets
- 13 13.0 **NON-CROP USES AROUND THE FARMSTEAD**
 - 13.1 General Weed Control and Trim-And-Edge
 - 13.2 Greenhouse/Shade house
 - 13.3 Chemical Mowing
 - 13.4 Cut Stumps
 - 13.5 Habitat Management
- 14 14.0 **ANNUAL WEEDS RATE SECTION**
 - 14.1 Annual Weeds - Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K
 - 14.2 Annual Weeds - Hand-Held or Backpack Equipment
 - 14.3 Annual Weeds - Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems
- 15 15.0 **PERENNIAL WEEDS RATE SECTION**
- 16 16.0 **WOODY BRUSH AND TREES RATE SECTION**
- 17 17.0 **LIMIT OF WARRANTY AND LIABILITY**

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....41.0%

OTHER INGREDIENTS:.....59.0%

100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

2.0 IMPORTANT PHONE NUMBERS

(To be completed later)

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (c) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Precautions:

3.2 Environmental Hazards

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and transported in stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR TRANSPORT THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Supplemental labeling may be found on the cdms.net or greenbook.net websites or obtained by contacting your authorized Repar-Glypho, LLC, retailer or Repar-Glypho, LLC, Company representative.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/esppj>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

[Container Label Storage and Disposal Statements]:

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact Repar-Glypho, LLC. Company at 1-301-562-7330.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

Refiners: Do not reuse this container except for refill in accordance with a valid Repar-Glypho, LLC. Agreement for Repackaging or Toil Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Repar-Glypho, LLC. container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

Product Description: This product is a postemergence, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds effects may not be visible for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" for specific weed recommendations.

Always use the higher product application rate within the range when weed growth is heavy or dense or when weeds are growing in an undisturbed (non-cultivated) area.

Reduced weed control may result from treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by this herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates: The maximum application or use rates stated throughout this product's labeling are given in units of volume (fluid ounces, pints, or quarts) of this product per acre. However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as text mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of this product (6 pounds of glyphosate acid) per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product (8 pounds of glyphosate acid) per acre per year.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

6.0 WEED RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE
-------	---	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using cultural and mechanical practices.

6.1 General Weed Management Recommendations

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management recommendations:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready crops without rotation of crops is to rotate to other Roundup Ready crops to allow use of alternate tank mixtures.
- Utilize the label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Repair-Glypho, LLC. representative, local retailer, or county extension agent.

6.2 Management For Glyphosate Resistant Biotypes

Note: Contact your Repair-Glypho, LLC. representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

The following agronomic practices may be used to reduce the spread of confirmed or unconfirmed glyphosate resistant weed biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops that allows use of herbicides that controls the resistant biotypes.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes:

7.0 MIXING

Spray solutions of this product should be mixed, stored and applied using only clean stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: PRODUCT PERFORMANCE MAY BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

7.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Begin filling the mixing tank or spray tank with clean water. Add the amount of this product near the end of the filling process and mix gently. Use caution to avoid siphoning back into the carrier source. During mixing, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

7.2 Tank Mixtures This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and follow all label directions of all products in the tank mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not in this labeling, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC. for this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT for applications of this product unless otherwise noted in this product label, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC.

When a tank mixture with a generic active ingredient, such as diuron, atrazine, 2,4-D or dicamba, as in this label, the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank mixture.

Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture, and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Use according to the most restrictive label requirements.

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance. For best results, tank mixtures with this product be applied at a minimum spray volume rate of 10 gallons per acre.

7.2 Tank Mixing Procedure

Prepare tank mixtures of this product as follows:

1. Place a 20- to 35-mesh screen or wetting basket over the filling port of the tank.
2. Through the screen, fill the tank one-half full with water and start gentle agitation.
3. If ammonium sulfate is to be used, add it slowly through the screen into the tank, and continue adding water into the tank through the screen. If dry ammonium sulfate is being used, ensure that it is completely dissolved in the tank before adding other products.
4. If a wettable powder is used, make a slurry with the water, and add it SLOWLY through the screen into the tank while continuing gentle agitation.
5. If a flowable formulation is used, premix one part flowable with one part water, and add the diluted mixture SLOWLY through the screen into the tank while continuing gentle agitation.
6. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water, and add the diluted mixture slowly through the screen into the tank while continuing gentle agitation.
7. Continue filling the tank with water through the screen and add the required amount of this product near the end of the filling process.
8. If a nonionic surfactant is used, add it to the tank before completing the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquids (this product) followed by surfactant.

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying.

Keep by-pass and return lines on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

7.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product as indicated in the following table in water:

Spray Solution**Amount of GLYPHOSATE 41% SL**

Desired Volume	0.5%	1%	1.5%	2%	5%	10%
1 gal	0.7 oz.	1.3 oz.	2 oz.	2.7 oz.	6.5 oz.	13 oz.
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in backpack sprayers, the appropriate amount of this product be mixed with water in a larger container and then fill the sprayer with the mixed solution.

7.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

7.6 Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates in this label. Lower rates will result in reduced performance.

7.7 Colorants or Dyes

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce performance. Use colorants or dyes according to the manufacturer's recommendations.

7.8 Drift Reduction Additives

Drift reduction additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet Applicator (CDA) equipment. When a drift reduction additive is used, read and carefully observe the precautions, limitations, and all other information appearing on the additive label. Use of drift reduction additives can affect spray coverage, which can reduce product performance.

8.0 APPLICATION EQUIPMENT AND TECHNIQUES Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment: Aerial—Fixed wing and helicopter.

Ground Broadcast Spray—Boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held or Backpack Spray Equipment—Backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment—Shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems--Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA)—Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING THE DESIRED VOLUMES.

8.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED IN THIS LABEL.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling. Use the rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 2 quart per acre. Refer to the individual use area sections of this label for further instructions.

FOR AERIAL APPLICATION IN ARKANSAS OR CALIFORNIA, OR SPECIFIC COUNTIES THEREIN, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE OR COUNTY FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

This product, when tank mixed with dicamba, may not be applied by air in California.

Ensure uniform application - to avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target movement during aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Comply with all State regulations where applicable.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (read the following sections on "Wind", "Temperature and Humidity", and "Temperature Inversions").

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures rates for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing spray pressure.
- **Number of nozzles:** Use the minimum number of nozzles that will provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind speed, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke produced by a ground source or ^{an} aircraft smoke generator can also identify temperature inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water. Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may help prevent corrosion.

8.2 Ground Broadcast Equipment

Apply the product rates in this label at 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the range to ensure complete coverage. Carefully select proper nozzles to avoid generating a fine mist. For best results with ground application equipment, use flat spray nozzles. Check spray pattern for uniform distribution.

8.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled on a spray-to-wet basis; do not spray to the point of run-off. Spray coverage should be uniform and complete. Use coarse sprays only. For rates and timing, refer to the "ANNUAL WEEDS HAND-HELD OR BACKPACK EQUIPMENT" section of this product label.

8.4 Selective Equipment

This product may be diluted in water and applied through shielded sprayers, hooded sprayers, wiper applicators or sponge bars to weeds growing in any non-crop site specified on this label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants), and wipers may be used over-the-top of crops only when specifically in this product's labeling. Such equipment must be capable of preventing all crop contact with the herbicide solutions and operated without leakage of spray mists or dripping onto crop.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of this product with desirable vegetation may result in unwanted plant damage or destruction. Crop injury may occur when the foliage of treated weeds comes into contact with leaves of the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Sprayers

This product, when applied at rates under the conditions described in the following paragraphs for shielded and hooded sprayers, will control those weeds listed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Adjust the shields on these sprayers to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is capable of completely enclosing the spray pattern. If necessary, extend the front and rear flaps of the hooded applicator downward to reach the ground in deep furrows. EXTREME CARE MUST BE TAKEN TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

This equipment must be configured and operated in a manner that minimizes bouncing and avoids raising the hoods up off the ground at any time. If the hood is raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operating this equipment on rough or sloping terrain where the spray hoods might be raised up off the ground surface.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure low-drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is used. Spray volume should be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- Spray hoods must be operated on the ground or skimming across the ground surface.

- Leave at least an 8-inch untreated strip over the drill row. (For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.)
- Operate at ground speeds of no greater than 5 miles per hour to avoid bouncing of the spray hoods.
- Apply when wind speeds are 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Wiper Applicators

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators: Use solutions ranging from 33 to 75 percent of this product in water. **For**

Panel Applicators: Use solutions ranging from 33 to 100 percent of this product in water.

8.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products for use in injection systems.

8.6 CDA Equipment

The rate of this product applied per acre by vehicle-mounted controlled droplet applicator (CDA) equipment must not be less than the amount used in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply in 2 to 15 gallons of water per acre.

For control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre).

Controlled droplet applicators produce a spray pattern that is not easily visible. Extreme care must be taken to avoid spray or drift onto the foliage or any other green tissue of desirable vegetation, as damage or destruction of the plant may result.

9.0 ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the "ROUNDUP READY CROPS" section of this label or separately published Repar-Glypho, LLC. Supplemental Labeling for this product for instructions on treating Roundup Ready crops.

TYPES OF APPLICATIONS:

Chemical Fallow, Preplant Fallow Beds, Preplant, At-Planting, Preemergence, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest.

GENERAL USE INSTRUCTIONS:

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", AND WOODY BRUSH AND TREES RATE SECTIONS" in this label. Use rates of this product that are emphasized in this product's labeling to control tough weeds take precedence over the general rates in the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS". Repeat applications may be made up to a maximum of 8 quarts of this product per acre per year.

Post-directed hooded sprayers and wiper applicators capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted in the individual crop sections that follow, wipers may also be used above certain crops to control tall weeds. Refer to the "SELECTIVE EQUIPMENT" section of this label for essential precautions regarding crop injury. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified in this product's labeling. Refer to the "AERIAL EQUIPMENT" section of this label for additional information.

GENERAL PRECAUTIONS, RESTRICTIONS:

Do not allow this herbicide to contact foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of. When making at-planting and preemergence applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death of emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any crops not listed on this label. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction-

For broadcast postemergence treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

9.1 Cereal and Grain Crops

Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Rye, Quinoa, Teff, Teosinte, Triticale, Wheat (all types), Wild rice.

RESTRICTIONS: Do not treat rice fields or levees when flooded.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Red Rice Control Prior to Planting Rice, Spot Treatment (Except Rice), Over-The-Top Wiper Application (Feed Barley and Wheat only), Preharvest (Wheat And Feed Barley Only).

Preplant, Preenemergence, At-Planting

Apply before, during or after planting of cereal crops. Applications must be made prior to crop emergence.

TANK MIXTURES: In wheat, a tank mix with Aim may be used.

Red Rice Control Prior to Planting Rice

Apply 1.5 quarts of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are at the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

RESTRICTIONS: Do not spray during conditions of low humidity, as reduced control may result. Do not treat rice fields or levees when the fields contain floodwater. Do not flood treated fields for 8 days following application.

Spot Treatment (Except Rice)

Apply as a spot treatment in cereal crops. Apply this product before heading in small grains.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Over-the-Top Wiper Application (Feed Barley and Wheat only)

Use Wiper applications in wheat and feed barley. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth.

PRECAUTIONS, RESTRICTIONS: Allow at least 35 days between application and harvest. Do not use roller applicators. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation will result in discoloration, stunting or destruction.

Preharvest (Feed Barley And Wheat Only)

This product provides weed control when applied prior to harvest of feed barley or wheat. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard-dough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest.

For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1 quart of this product per acre. Allow 7 days between application and harvest or grazing. Preharvest application is not used for barley or wheat grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

Apply after harvest of cereal crops. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.2 Corn

TYPES OF CORN: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest.

Use directions for corn containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply alone or in a tank-mixture before, during or after planting corn. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with the following products to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Ensure that the specific product is registered for application prior to planting corn. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

2,4-D	Distinct™	Leadoff™
Aim [®]	Dual Magnum™	Alachlor
Atrazine	Dual II Magnum™	Linex™/Lorox™
Axiom™	Epic™	Keystone LA
Balance™ / Balance PRO	Frontier™/Outlook™	Marksman™
Banvel™/Clarity™	FulTime™	Micro-Tech [®]
Bicep Magnum™	Guardsman™/Guardsman MAX	Prowl™
Bicep II Magnum™	Harness [®]	Python™
Bullet [®]	Harness Xtra	Resource [®]
Degree [®]	Harness Xtra 5.6l	Simazine Surpass
Degree Xtra [®]	Hornet	Topnotch™
	Lariat [®]	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

RESTRICTIONS: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. The area covered by this restriction includes Illinois and Indiana south of Route 50, and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

Hooded Sprayers

Apply with hooded sprayers for weed control between rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instructions for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Corn must be at least 12 inches tall, measured without extending leaves. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Do not apply more than 1 quart of this product per acre for each application and no more than 3 quarts per acre per year for hooded sprayer applications.

Spot Treatment

This product may be applied as a spot treatment prior to silking of corn.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Take care not to spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

Apply at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the ear is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 2 quarts of this product per acre.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Preharvest application is not used for corn grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required to control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.3 Cotton

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Selective equipment, Spot treatment, Preharvest.

Use directions for cotton containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton.

TANK MIXTURES: This product may be tank-mixed with the following products provided that the specific product being used is registered for application prior to planting cotton. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Caparol®	Direx	Prowl
Clarity™	Dual Magnum™	Prowl H2O
Command	Dual II Magnum™	Staple®
Cotoran®	Karmex	Stalwart
Cotton PRO®	Meturon®	Zorial®

PRECAUTIONS, RESTRICTIONS: Applications must be made prior to crop emergence. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Selective Equipment

Apply through hooded sprayers, shielded sprayers or wiper applicators in cotton. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Allow at least 7 days between application and harvest. Do not treat more than 10 percent of the total field area to be harvested.

Spot Treatment

Apply as a spot treatment in cotton prior to boll opening.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in the treated area will be killed. Do not spray or to allow spray to drift outside target area to avoid unwanted crop destruction.

Preharvest

This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "ANNUAL WEEDS", and "PERENNIAL WEEDS RATE SECTIONS" of this label. For cotton regrowth inhibition, apply 1 pint to 2 quarts of this product per acre.

Up to 2 quarts of this product may be applied after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

TANK MIXTURES: Tank mix with DEF TM 6, Folex™, Ginstar, or Prep™ to provide additional enhancement of cotton leaf drop.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Do not apply preharvest for cotton grown for seed, as a reduction in germination or vigor may occur. THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF THIS PRODUCT TO COTTON IS PROHIBITED. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

9.4 Fallow Systems

Apply during the fallow period prior to planting or emergence of any crop listed on this label; for non-listed crops, applications must be made at least 30 days prior to planting.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Aid-to-Tillage, Chemical Fallow

Use as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Tank mixtures of this product with 2,4-D and dicamba may be used for a broader weed control spectrum. Applications up to 2 quarts per acre may be made by aerial application onto fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

RESTRICTIONS: Do not apply dicamba tank mixtures by air in California. Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

Preplant Fallow Beds

This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label prior to planting.

TANK MIXTURES: Use 12 fluid ounces of this product, plus 2 to 3 fluid ounces of Goai™ 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 3" -- common cheeseweed, chickweed, groundsel; 6" -- London rocket, shepherd's-purse.

Use 16 fluid ounces of this product, plus 2 to 3 fluid ounces of Goai™ 2XL, per acre to control the following weeds up to the maximum height, diameter or length indicated: 6" -- common cheeseweed, groundsel, marestail (*Coryza canadensis*), 12" -- chickweed, London rocket, shepherd's-purse.

Aid-to-Tillage

Apply in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 12 fluid ounces of this product in 3 to 10 gallons of water per acre before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

Tank mixtures with residual herbicides may result in reduced product performance.

9.5 Grain Sorghum (Milo)

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Over-the-Top Wiper Application, Preharvest.

Preplant, At-Planting, Preemergence

Apply alone or in tank-mixture before, during or after planting grain sorghum. Applications must be made prior to crop emergence.

TANK MIXTURES: Apply with one of the following pesticide products as a tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, provided that the specific product being used is labeled for application prior to planting grain sorghum.

Atrazine	IntRRo
Bicep II Magnum	Lariat
Bullet	Micro-Tech
Dual II Magnum	

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply a tank mixture of this product at 2 pints per acre plus the products listed above. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

For other labeled annual weeds listed on this label, apply 1.5 to 2 pints of this product per acre when weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased for acceptable weed control.

Spot Treatment, Over-the-Top Wiper Application

Apply as a spot treatment in grain sorghum before heading. Apply with wiper applicators to control or suppress tall weeds. For additional instructions, see "Wiper Applicators" in the "SELECTIVE EQUIPMENT" section of this label.

RESTRICTIONS: For spot treatment, do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Do not spray or to allow spray to drift outside target area to avoid unwanted crop destruction. Allow at least 40 days between wiper application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

Hooded Sprayers

Apply with hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Milo must be at least 12 inches tall, measured without extending leaves. Treat before mile, sends tillers between the drill rows. If tillers are sprayed, the main plant may be damaged or destroyed. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage, which shall be the sole responsibility of the applicator. Do not graze or feed milo forage or fodder following hooded sprayer applications. Do not apply more than 1 quart of this product per acre per application and no more than 3 quarts per acre per year for hooded sprayer applications.

Preharvest

Apply for weed control prior to harvest after sorghum grain has reached 30 percent grain moisture or less.

RESTRICTIONS: Do not apply more than 2 quarts of this product per acre. As with other herbicides that cause sudden plant death, avoid preharvest applications of this product to milo infected with charcoal rot as lodging can occur. Allow a minimum of 7 days between application and harvest of grain sorghum. Preharvest application is not used for sorghum grown for seed, as a reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California.

Post-Harvest

Apply after harvest of grain sorghum. Higher rates may be required to control large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product per acre for control, or 1.5 pints of this product per acre for suppression.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.6 Herbs and Spices

Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Cilantro (leaf), Cilantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Over-the-Top Wiper Application (Peppermint and Spearmint Only), Spot Treatment (Peppermint and Spearmint Only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Over-the-Top Wiper Application, Spot Treatment (Peppermint and Spearmint only)

Apply as a spot treatment or over the top of peppermint or spearmint with wiper applicators in spearmint and peppermint or apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution to a limited area.

RESTRICTIONS: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested should be treated at one time. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. For wiper applications, contact of the herbicide solution with the crop may result in discoloration, stunting, or destruction.

9.7 Oil Seed Crops

Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower.

Use directions for canola containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

Apply before, during or after planting oil seed crops listed in this section. Broadcast applications must be made prior to crop emergence. Wiper applicators or hooded sprayers may be used between the rows once the crop is established. See additional instructions of the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl may be applied before, during or after planting into conventionally tilled soil, a cover crop, established sod or previous crop residue.

RESTRICTIONS: For use with canola, do not apply more than 2 quarts of this product per acre. For use with sunflowers, do not apply more than 1 quart of this product per acre as a single preplant or preemergence application per year. Do not feed or graze sunflower forage following application of this product.

9.8 Soybeans

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: Spot Treatment, Preharvest, Selective Equipment.

Use directions for soybeans containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply alone or in a tank mixture before, during or after planting soybeans. Applications must be made prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with one or more of the following products to provide residual weed control, a broader spectrum, or an alternate mode of action. Ensure that the specific product being used in the tank mixture is registered for application prior to planting soybean, and follow all label directions of all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre.

Aim™	Domain™	Micro-Tech
Assure II™	Dual Magnum	Prowl
Authority™	Dual II Magnum	Pursuit™
Axiom	Firstate™	Pursuit Plus
Blanket	Flexstar™	Python
Boundary	Frontier™/Outlook™	Reflex™
Canopy™	Fusion™	Resource™
Canopy EXT™	Gauntlet™	Scepter™
Classic	Gangster	Sensor™/Lexone™
Command™	IntRRo	Squadron™
Command Xtra™	Linux™	Steel™
2,4-D	Lorox/Linaron	Valor™
	Lorox Plus™	

For tough-annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of this product per acre when

weeds are less than 6 inches tall, and 2 to 3 pints per acre when weeds are over 6 inches tall. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Spot Treatment

Apply as a spot treatment prior to initial pod set in soybeans.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction.

Preharvest

Apply for weed control prior to harvest of soybeans after pods have set and lost all green color. Apply at rates given in the "ANNUAL WEEDSRATE SECTION" and "PERENNIAL WEEDSRATE SECTION". Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Do not apply more than 5 quarts of this product per acre for preharvest applications. Do not apply more than 2 quarts of this product per acre by air. Allow a minimum of 7 days between application and harvest of soybeans. Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last preharvest application. (If the application rate is 1 quart per acre or lower, the grazing restriction is reduced to 14 days after last preharvest application.) Preharvest application is not used for soybeans grown for seed, as a reduction in germination or vigor may occur.

Selective Equipment

Apply through shielded or hooded sprayers, wiper applicators or sponge bars in soybeans. Additional instructions on the use of selective equipment are found in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow at least 7 days between application and harvest.

9.9 Sugarcane

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus Spot Treatment and Sugarcane Ripening.

Preplant, At-Planting, Preemergence

Apply in or around sugarcane fields, or in fields prior to the emergence of plant cane.

RESTRICTIONS: Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Spot Treatment

Apply as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using spray-to-wet or a spray-to-wet technique. Volunteer or diseased sugarcane should have at least 7 new leaves.

RESTRICTIONS: Do not allow spray contact with healthy sugarcane plants because severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

Fallow Treatments

Apply as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4 to 5 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Applications up to 3 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used.

Hooded Sprayers

Apply using hooded sprayers for weed control between the rows of sugarcane. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional use instructions.

RESTRICTIONS: Crop injury may occur when the foliage of treated weeds contact the crop. Do not apply this product when crop leaves are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction.

Sugarcane Ripening

Apply to foliage as a plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane. It is effective in both low and high-tonnage sugarcane.

When applied as directed under the conditions described, this product will hasten ripening and extend the period of high sucrose level in sugarcane.

As a result of leaf desiccation, improved trash burn can be expected.

Most of the sucrose increase is concentrated in the top nodes of the treated sugarcane stalk. In order to maximize sugar where topping is practiced during harvest, top at the base of the fourth leaf.

APPLICATION RATES:

NOTE: Use the higher rate within the used range when treating sugarcane under adverse ripening conditions or when less responsive varieties are to be treated.

Use the following application rates and timing instructions according to the State in which the sugarcane is grown.

FLORIDA—Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of LAST PLATOON CANE ONLY.

HAWAII—Apply 12 to 28 fluid ounces of this product per acre 4 to 10 weeks before harvest.

LOUISIANA—Apply 5 to 16 fluid ounces of this product per acre 3 to 7 weeks before harvest of PLATOON CANE ONLY.

PUERTO RICO—Apply 7 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

TEXAS—Apply 7 to 16 fluid ounces of this product per acre 3 to 5 weeks before harvest of PLATOON CANE ONLY.

PRECAUTIONS, RESTRICTIONS: Prior to application, consult your state sugarcane authority or local Repar-Glypho, LLC representative regarding the degree of sucrose response anticipated from the variety of sugarcane to be treated.

Application of this product may initiate development of shooting eyes. This product may not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2 to 3 weeks after application, this product may produce a slight yellowing to pronounce browning and drying of leaves, and a shortening of upper internodes. Spindle death may occur.

Rainfall within 6 hours after application may reduce effectiveness.

Application is not used for sugarcane grown for seed, as a reduction in germination or vigor may occur.

Do not feed or graze treated sugarcane forage following application. Do not plant subsequent crops in treated fields other than the following for 30 days after application: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybeans, squash (all types) or wheat.

9.10 Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL VEGETABLE CROPS LISTED ALPHABETICALLY IN THIS SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Chemical Fallow, Preplan Fallow Beds, Preplan, At-Planting, Preemergence, Prior to Transplanting Vegetables, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Application in Row-Middles, Post-Harvest, Selective Equipment Applications (Nonbearing Ginseng), Over-the-Top Wiper Application (Rutabagas only), Spot Treatment or Preharvest (Dry Beans, Peas, Lentils, and Chickpeas only).

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

9.10.1 Brassica Vegetables

Broccoli, Chinese broccoli (gai lan), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

9.10.2 Bulb Vegetables

Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

9.10.3 Cucurbit Vegetables and Fruits

Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Melons (all), *Momordica spp* (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon.

RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

9.10.4 Leafy Vegetables

Amaranth (Chinese spinach), Arugula (Rocket), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow koo, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

RESTRICTIONS: For Watercress, do not within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of crop injury.

9.10.5 Fruiting Vegetables

Eggplant, Groundcherry (*Physalis spp.*), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimiento, sweet pepper), Tomatillo, Tomato.

RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato and tomatillos grown on sandy soil, do not make hooded, shielded sprayer or wiper applications in row middles because of potential for crop injury.

9.10.6 Legume Vegetables (Succulent or Dried)

Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

Spot Treatment, or Preharvest (Dry Beans, Peas, Lentils and Chickpeas only)

Apply as an over-the-top broadcast spray or as a spot treatment to control labeled weeds in: dry beans, peas, lentils or chick peas. For spot treatment, to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed, apply in 10 to 20 gallons of water per acre through ground broadcast spray equipment or use a 2 percent solution in a hand-held sprayer. For preharvest treatments, apply in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less).

PRECAUTIONS, RESTRICTIONS: Follow the limitations listed in the table below. Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not specified for treatment in this label. Do not feed treated vines and hay from these crops to livestock. Do not treat cowpeas or field (feed) peas, since these crop are considered to be grown as livestock feed. Preharvest application is not used for dry legumes grown for seed, as a reduction in germination or vigor may occur.

Crop	Maximum Rate (fluid ounces per Acre)	Minimum Pre- Harvest Interval (days)	Allowed in
Dry Beans	32	7	All States
Dry Peas, Lentils, and Chickpeas	26	14	Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, and Washington

9.10.7 Root and Tuber Vegetables

Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

Selective Equipment Applications (Non-Bearing Ginseng only)

Apply for general weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and backpack wands, lances, and orchard guns or with wiper application equipment. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Conduct applications so that there is no contact of this product with the ginseng plant. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Applications must be made at least one year prior to harvest.

Over-the-Top Wiper Application (Rutabagas only)

Use Wiper applicators over the top of rutabagas for the control of taller weeds. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

9.11 Miscellaneous Crops

Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus the following: General Weed Control, Site Preparation, Spot Treatment (Asparagus).

Use directions for sugar beets containing a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

RESTRICTIONS: Avoid contact of this product with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe crop injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-Harvest or fallow applications must be made at least 30 days prior to planting any crop not listed on this label. See additional information in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

General Weed Control, Site Preparation

This product may be applied for general weed control or for site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS, RESTRICTIONS: This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Care must be taken to remove residues of this product from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or by irrigation. Care should be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Allow at least 21 days between residue removal and transplanting. Applications made at emergence will result in injury or death to emerged seedlings.

Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.

Spot Treatment (Asparagus)

Apply immediately after cutting, but prior to the emergence of new spears.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Post-Harvest (Asparagus)

Apply after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. See additional instructions under "Selective Equipment" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS, RESTRICTIONS: Direct contact of the spray with asparagus may result in serious crop injury.

10.0 TREE, VINE, AND SHRUB CROPS (Alphabetical)

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL TREE, VINE, AND SHRUB CROPS LISTED ALPHABETICALLY IN THE SECTIONS THAT FOLLOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Preplant (site preparation) Broadcast Sprays, General Weed Control, Middles (between rows of trees, vines or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (shielded sprayers, wiper treatments), Directed Spray, Spot Treatment, Perennial Grass Suppression, Cut Stump.

This product may be applied with boom equipment, CDA equipment, shielded sprayers, hand-held and backpack wands, lances, orchard guns or with wiper applicator equipment, unless prohibited in the specific Crop sections that follow.

Apply in middles (between rows of trees or vines), strips (within rows of trees or vines), and for general weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 1 pint to 5 quarts per acre according to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label. Use the higher rates in the range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts per acre per year. See the "GENERAL INFORMATION" section of this label for more information on Maximum Annual Use Rates.

GENERAL PRECAUTIONS, RESTRICTIONS: Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used in order to minimize the potential for overspray or drift of this product onto the crop. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with the crop may be used. See additional instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

Allow a minimum of 3 days between application and transplanting.

Middles (between rows)

This product will control or suppress annual and perennial weeds and ground covers growing between rows of tree and vine crops listed on this label. If weeds are under drought stress, irrigate prior to application. Reduced weed control may result if weeds have been recently mowed at the time of application.

TANK MIXTURES: A tank mixture of this product plus Goal 2XL may be applied for annual weed control between rows (middles) of citrus crops, tree fruits, tree nuts and vine crops. This mixture is used when weeds are stressed or growing in dense populations. Application of 16 to 32 fluid ounces of this product per acre plus 3 to 12 fluid ounces of Goal 2XL per acre will control annual weeds with a maximum height, length or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). This tank mixture will also control common cheeseweed (malva) or hairy fleabane (*Conyza bonariensis*) with a maximum height, length or diameter of 3 inches. Read and follow all labels and directions for use of all products being used in the tank mixture.

Strips (In Rows)

TANK MIXTURES: This product may be applied within rows of tree or vine crops in tank mixtures with the following products, provided that the label of the specific product used permits the desired use:

Devrinol TM 50 DF	Simazine 4L
Direx TM 4L	Simazine 80W
Goal 2XL	Sim-Trol TM 4L
Karmex DF	Solican TM DF
Krovarl	Surflan TM AS
Prowl	Surflan 75w
Princep Caliber TM 90	

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for information about specific crops, rates, geographic restrictions and precautionary statements. For more information see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Perennial Grass Suppression

This product will suppress perennial grasses such as bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

For burndown of Bermudagrass, apply 1 to 2 quarts of this product in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 6 to 16 fluid ounces of this product per acre east of the Rocky Mountains and 16 fluid ounces of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the Bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces of this product per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

Cut Stump (tree crops)

Make cut stump applications of this product during site preparation or site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

Citrus Trees: Calamondin, Chionja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo (Ugli), Tanger.

Fruit Trees: Apple, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince.

Nut Trees: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).

Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

PRECAUTIONS, RESTRICTIONS: DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

10.1 Berry Crops

Blackberry (including bingieberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucetlaberry, marionberry, nectarberry, ofallieberry, Oregon evergreen berry, phenomenalberry, rangberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Salal. [Optional Crop: Raspberry (black, red)].

TYPES OF APPLICATIONS: Those listed in Section 10.0, plus Spot Treatment in Cranberry Production and Post-Harvest Treatment in Cranberry Production.

RESTRICTIONS: To avoid damage, herbicide spray must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest of cranberries. Allow a minimum of 14 days between last application and harvest of other berry crops.

Spot Treatment in Cranberry Production

Use spot treatments to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in this label may be used. Drop water level to remove standing water in ditches prior to application. With hand-held sprayers, use 1- to 2-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off.

RESTRICTIONS: To achieve maximum weed control in treatments of dry ditches after water draw down, allow 2 or more days after treatment before reintroduction of water. Apply this product within 1 day after draw down to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not make applications by air. Do not apply directly to water. Use nozzles that emit medium- to large-sized droplets in order to minimize spray drift and avoid crop injury.

Post-Harvest Treatments in Cranberry Production

Apply this product may be made after the harvest of cranberries to control weeds growing in the field. Apply this product to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label may be used. If using hand-held sprayers, use a 0.5- to 1-percent solution of this product. Spray adequately to wet the vegetation only; do not spray to the point of run-off. If using hand-held boom sprayers, apply 2 to 4 quarts of this product per acre.

RESTRICTIONS: Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not make applications by air. Do not apply directly to water. Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

10.2 Citrus

Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugh), Tangor.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only.

The following use instructions pertain to applications in Florida and Texas only: For burndown or control of the weeds listed below, apply this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 2 to 3 quarts of this product in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall, and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, use of this product in a tank mixture with Krovar™ I or Kannex™ may improve weed control. Refer to the individual product labels for treatment information on specific crops, rates, geographic restrictions and precautionary statements.

Perennial weeds:

S = Suppression

B = Burndow

PC = Partial control

C = Control

WEED SPECIES	GLYPHOSATE 41% SL RATE PER ACRE			
	1 QT	2 QT	3 QT	5 QT
Bermudagrass	B	---	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	---	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	---	PC	C

10.3 Miscellaneous Tree Food Crops

Cactus (fruit and pads), Palm (heart, leaves), Palm (oil).

TYPES OF APPLICATIONS: Those listed in Section 10.0.

10.4 Non-Food Tree Crops

Pine, Poplar, Eucalyptus, Christmas trees.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

NOTE: unless otherwise directed, this product is not used for use as an over-the-top broadcast spray in plantations or other labeled tree crops.

Directed Spray, Spot treatment, Wiper Application

This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

PRECAUTIONS, RESTRICTIONS: Avoid contact of spray, drift or mist of this product with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Site preparation

Apply this product for weed control prior to planting non-food tree crops.

PRECAUTIONS: Precautions should be taken to protect non-target plants during site preparation applications.

10.5 Pome Fruit

Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince. TYPES OF

APPLICATIONS: Those listed in Section 10.0.

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in pome crops.

10.6 Stone Fruit

Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), Plumcot.

TYPES OF APPLICATIONS: Those listed in Section 10.0. For olive groves, apply only as a directed spray.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 17 days between last application and harvest in stone fruit crops.

Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. USE EXTREME CARE TO ENSURE THAT NO PART OF THE PEACH TREE IS CONTACTED WITH OVERSPRAY OR DRIFT OF THIS PRODUCT.

10.7 Tree Nuts

Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (black, English).

TYPES OF APPLICATIONS: Those listed in Section 10.0.

RESTRICTIONS: Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut.

10.8 Tropical and Subtropical Trees and Fruits

Ambarella, Atemoya, Avocado, Barana, Barbadoscherry (acerola), Biriba, Blinabe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates,

Durian, Feijoa, Figs, Governor's plum, Guava, Hama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genie), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots and leaves), Wax jambe.

TYPES OF APPLICATIONS: Those listed in Section 10.0, and as described below as a Bananacide (Bananas Only).

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit listed. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

Bananacide (Banana Only)

Apply to destroy banana plants infected with the Banana Bunchy Top Virus, as well as non-infected banana plants in order to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1/25 fluid ounce (1 mL) of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) within a 4-foot radius around a treated mat should be mechanically destroyed.

For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

PRECAUTIONS, RESTRICTIONS: Do not apply more than 1/2 fluid ounce (15 mL) of this product per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for general weed control.

10.9 Vine Crops

Grapes (raisin, table, wine) Hops, Kiwi, Passion fruit.

TYPES OF APPLICATIONS: Those listed in Section 10.0.

Apply this product for weed control only when green shoots, canes or foliage are not in the spray zone.

In the northeast and Great Lakes regions, apply this product to grape vineyards prior to the end of the bloom stage of grapes in order to avoid crop injury, or apply using shielded sprayers or wiper equipment.

RESTRICTIONS: Allow a minimum of 14 days between last application and harvest of vine crops. Do not use selective equipment in kiwi.

11.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELANDS

GENERAL USE INSTRUCTIONS: Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of this label for rates for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. Application rates of this product specified on this label for the control of tough weeds, or those specified on separate supplemental labeling for this product, supersede rates in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" of this label.

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling.

11.1 Alfalfa, Clover, and Other Forage Legumes

Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types).

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Preharvest (except Kenaf and Leucaena), Spot Treatment, Over-the-Top Wiper Application, Renovation, Stand Removal.

Use directions for alfalfa with a glyphosate tolerant gene are in the "ROUNDUP READY CROPS" section of this label.

Preplant, At-Planting, Preemergence

Apply before, during or after planting crops listed in this section. Applications must be made prior to crop emergence.

RESTRICTIONS: Remove domestic livestock before application. Do not feed or graze until crop has reached sufficient maturity.

Preharvest (Except Kenaf and Leucaena)

This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This product will control annual or perennial weeds, including quackgrass, when applied prior to crop harvest. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

PRECAUTIONS, RESTRICTIONS: Make only one application to an existing stand of crop per year. The treated crop and weeds can be harvested and fed to livestock according to the application rates and intervals defined below.

	Maximum single application rate (per acre)	Minimum interval between application and harvest or livestock grazing
Alfalfa	2 quarts	36 hours
All other legumes listed	3 pints	3 days

This application may destroy the alfalfa stand and may severely injure or destroy other labeled crops such as clover. Do not apply for preharvest application for alfalfa grown for seed, as a reduction in germination or vigor may result.

Spot Treatment, Over-the-Top Wiper Application

Apply as a spot treatment or over the top of crops listed in this section with wiper applicators. For wipers, see "Wiper Applicators" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field area should be treated at one time. Remove domestic livestock before application and wait 3 days after application before grazing livestock or harvesting.

Renovation, Stand Removal

Apply as a broadcast spray to remove established stands of alfalfa, clover, and other forage legumes listed on this label. If the crop is to be grazed or harvested for feed, use up to a maximum of 2 quarts per acre in alfalfa and up to 3 pints per acre in other labeled legumes. For complete removal of established stands of clover, it may be necessary to use treatment rates greater than this, as listed in the "PERENNIAL WEEDS RATE SECTION" of this label.

PRECAUTIONS, RESTRICTIONS: For applications up to 2 quarts per acre for alfalfa or 3 pints per acre for all other forage legumes, remove domestic livestock before application, follow the minimum harvest or grazing intervals in the table above under Preharvest in this section. For treatment rates above these levels, do not graze or harvest treated foliage for livestock feed or allow grazing. Crops listed in this label may be planted into the treated area at any time; for all other crops, wait 30 days or more between application and planting.

11.2 Conservation Reserve Program (CRP)

TYPES OF APPLICATIONS: Renovation (Rotating Out Of CRP), Site Preparation, Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application.

Renovation (Rotating Out of CRP), Site preparation

This product may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation recommendations.

RESTRICTIONS: Crops listed on this label may be planted into the treated area at any time; for any other crop, wait at least 30 days between application and planting.

Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Application

Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land. Applications may be made using wiper applicators to control tall weeds or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: *Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant.* No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

11.3 Grass Seed or Sod Production

Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation, Shielded Sprayers, Over-the-Top Wiper Application, Spot Treatment, Creating Rows in Annual Ryegrass.

Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stands, Site Preparation

This product controls most existing vegetation prior to renovating turf or forage grass seed areas or establishing turf grass grown for sod. It may be used to destroy undesirable grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting or for renovation. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turf management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates of up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed for treatment in this label, applications must be made at least 30 days prior to planting. Applications must be made prior to the crop emergence in order to avoid crop injury.

Shielded Sprayers

Apply 1 to 3 quarts of this product as a broadcast spray in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields. See additional instructions on the use of shielded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Over-the-Top Wiper Application

Apply this product for over the top of desirable grasses using wiper applicators for the control of tall weeds. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

PRECAUTIONS, RESTRICTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction.

Spot Treatment

Apply a 1- to 1.5-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may also be used to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS, RESTRICTIONS: Apply this product prior to heading of grasses. Do not treat more than 10 percent of the total field area. Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area to avoid unwanted crop destruction. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Creating Rows in Annual Ryegrass

Apply 16 to 32 fluid ounces of this product per acre. Best results are obtained when applications are made before ryegrass reaches 6 inches in height. Use the higher rate when the ryegrass is greater than 6 inches in height. Set nozzle heights to allow the establishment of the desired row spacing. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are used.

PRECAUTIONS: Take care not to spray or allow droplets, spray fines, or drift to settle outside the treatment area to avoid unwanted crop destruction.

Grower assumes all responsibility for crop losses from misapplication.

11.4 Pastures

Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label, including Bahiagrass, Bermudagrass, Bluegrass, *Bromegrass*, *Fescue*, *Guineagrass*, *Kikuyu grass*, *Orchardgrass*, *Pangola grass*, Ryegrass, Timothy, Wheatgrass.

TYPES OF APPLICATIONS: Preplant, Preemergence, Pasture Renovation, Spot Treatment, Over-the-Top Wiper Application, Postemergence Weed Control (Broadcast Treatments).

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be used to control perennial pasture species listed on this label prior to re-planting.

RESTRICTIONS: If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed on this label, applications must be made at least 30 days prior to planting.

Spot Treatment, Over-the-Top Wiper Application

This product may be applied in pastures as a spot treatment or over the top of desired grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section.

RESTRICTIONS: For spot treatment and wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acre, no more than 10 percent of the field may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing or harvesting for feed.

Postemergence Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after perennial grasses have reached dormancy.

PRECAUTIONS, RESTRICTIONS: Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto pasture grasses except for renovation use described above in this section. If replanting is needed due to severe stand reduction, wait at least 30 days after treatment prior to planting any crop not listed on this label.

11.5 Rangelands

TYPES OF APPLICATIONS: Postemergence.

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands. Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Apply 12 to 16 fluid ounces of this product to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are used where spring moisture is usually limited and fall germination allows for good weed growth.

For medusahead, apply 16 fluid ounces of this product at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

PRECAUTIONS, RESTRICTIONS: Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off. Do not use ammonium sulfate when spraying rangeland grasses with this product. No waiting period between treatment and feeding or livestock grazing is required. Do not apply more than 3 quarts of this product per acre per year.

12.0 ROUNDUP READY CROPS

The following instructions or those published separately on Repar-Glypho, LLC. supplemental labeling for this product include all applications of this product that can be made onto the specified Roundup Ready crops during the complete cropping season. DO NOT combine these treatment instructions with those for crops in the "ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)" section of this label that do not contain a glyphosate tolerant gene, unless otherwise directed in this product's labeling.

USE THIS PRODUCT FOR POST-EMERGENCE (IN-CROP) APPLICATION ONLY ON CROPS DESIGNATED AS CONTAINING A GLYPHOSATE TOLERANT GENE. Applying this product to crops that are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a glyphosate tolerant gene, since severe plant injury or destruction will result. The Roundup Ready designation indicates that the crop contains a patented gene that provides tolerance to this product. Information on Roundup Ready crops may be obtained from your seed supplier or Repar-Glypho, LLC. representative. Roundup Ready crops must be purchased from an authorized licensed seed supplier.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. **THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.**

For ground broadcast applications. Apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications. All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this product's labeling, in particular in the "Aerial Equipment" section. Apply this product in 3 to 15 gallons of water per acre. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

For proper stewardship of aerial applications over the top of Roundup Ready crops, read and follow all precautions and procedures contained in the use guide "A Guide to On-Target Aerial Application." Copies of this publication are available by calling 1-800-ROUNDUP (1-800-768-6387).

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE.

TANK MIXTURE: Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT for applications of this product over the top of Roundup Ready crops unless otherwise noted in this product label, or in separate supplemental labeling or fact sheets published by Repar-Glypho, LLC. for this product.

Unless otherwise directed, nonionic surfactant may be added to the spray solution for applications to Roundup Ready crops. The addition of certain surfactants to this product may result in some crop response including leaf necrosis, leaf chlorosis or leaf speckling due to the surfactant added to the spray mixture. Read

and carefully observe cautionary statements and other information appearing on the surfactant label.

Ammonium sulfate may be mixed with this product for applications to Roundup Ready crops. Refer to the "MIXING" section for instructions on the use of ammonium sulfate. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The following instructions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of this product is used to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed with multiple germination times, or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Rates of this product specified in the following sections, or in separate supplemental labeling published by Repar-Glypho, L.L.C. for this product on the control of tough weeds, supersede the general rate recommendations in the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" of this label.

12.1 Roundup Ready Alfalfa

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: Refer to the following table for the maximum application rates of this product.

Annual Maximum Application Rates	
Combined total per year for all applications, including preplant during year of establishment	8 quarts per acre
Combined total per year for in-crop applications for newly established and established stands	6 quarts per acre

GENERAL PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting alfalfa with a glyphosate tolerant gene, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

Apply this product may be made over the top of Roundup Ready alfalfa (in-crop) from emergence to 5 days prior to harvest. To maximize crop yield and quality potential, applications of this product should be made after weeds have emerged but before alfalfa growth or re-growth interferes with spray coverage of the target weeds.

Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" in this label for rates for specific weeds. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. In addition to those weeds listed in these sections, this product will suppress the parasitic weed Dodder (*Cuscuta spp*) in Roundup Ready alfalfa. Repeat applications may be necessary for complete control.

Stand Establishment—Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings may not contain a Roundup Ready gene and will not survive after the first application of this product. To eliminate the

undesirable effects of stand gaps created by this loss of plants, a single application of this product per acre should be applied at or before the 4-trifoliate growth stage. Refer to the following tables for maximum in-crop application rates during stand establishment (seeding year).

STAND ESTABLISHMENT (Seeding Year)	
Application Rates	
Prior to First Cutting	
From emergence up to 4 trifoliate leaves	1 to 2 quarts per acre
From 5 trifoliate leaves up to 5 days before first cutting	Up to 2 quarts per acre
After First Cutting	
In-crop application, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

Established Stands—Refer to the following table for directions and maximum application rates for in-crop applications to established stands of alfalfa (non-seeding year).

ESTABLISHED STANDS	
Application Rates	
In-crop applications, per cutting, up to 5 days before cutting	Up to 2 quarts per acre

PRECAUTIONS, RESTRICTIONS: Where Roundup Ready alfalfa is grown with a companion or cover crop, or is overseeded with a second species, in-crop (over-the-top) applications of this product will eliminate the non-glyphosate tolerant species. Any single in-crop application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart. The combined total per year for all in-crop applications in both newly established (seeding year) and established stands (non-seeding year) must not exceed 6 quarts per acre. Remove domestic livestock before application. Wait a minimum of 5 days after application before grazing, cutting and feeding of forage and hay.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application.

Newly Seeded Stands and Stand Establishment: For control of emerged annual grasses and broadleaf weeds, this product may be applied at up to 2 quarts per acre in a tank mixture with the following herbicides. Application should be made after weeds have emerged but before the alfalfa growth or regrowth would interfere with spray coverage of the target weeds

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, bromoxynil, clethodim, imazamox, imazethapyr, sethoxydim.

Arrow, Buctril, Butoxone, Butyrac, Poast, Pursuit, Raptor, Select.

Buctril can only be used in newly seeded stands]

Dormant Application and Winter Treatment: For control of emerged annual grasses and broadleaf weeds, up to 2 quarts per acre of this product may be applied in a tank mixture with the following herbicides. Tank mixtures should be applied when the temperature for the day remains above freezing.

Dormant Application

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: diuron, hexazinone, imazamox, metribuzin, pronamide, terbacil. Kerb 50-W, Lexone, Raptor, Sencor, Sinbar, Velpar AlfaMax.]

Winter Treatment

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa: 2,4-DB, diuron, hexazinone. Butoxone, Diuron, Velpar AlfaMax.]

PRECAUTIONS, RESTRICTIONS: Tank mixtures of this product with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control. Applications must not be made to frozen or snow covered ground.

12.2 Roundup Ready Canola (Spring)

Roundup Ready spring canola is defined as those Roundup Ready canola varieties that are seeded in the spring and harvested in the fall and do not enter a winter dormancy period.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

Annual Maximum Application Rates		GENERAL PRECAUTIONS
Preplant, At-planting, Preemergence applications	2 quarts per acre	
Total in-crop application from emergence to 6-leaf stage	1 quart per acre	

PRECAUTIONS AND RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates."

Preplant, At-planting, Preemergence,

This product may be applied before, during or after planting, up to a maximum of 2 quarts per acre. Postemergence (In-crop)

This product may be applied postemergence to Roundup Ready spring canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds. .

Single Application - Apply 16 to 24 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar crop injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application - Apply 16 fluid ounces per acre to 1 - to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Apply sequential applications for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass, or when multiple application times are needed for adequate weed control.

RESTRICTIONS: No more than two in-crop broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 32 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

12.3 Roundup Ready Canola (Winter)

Roundup Ready Winter Canola is defined as those Roundup Ready canola varieties that are seeded in early fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.

<u>Annual Maximum Application Rates</u>	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	1 quart per acre

DO NOT USE THIS PRODUCT ON WINTER CANOLA WITH A GLYPHOSATE TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA, EXCEPT FOR USES IN WILDLIFE FOOD PLOTS THAT WILL NOT BE HARVESTED FOR HUMAN OR LIVESTOCK FOOD.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL USE INSTRUCTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rate."

Preplant, At-Planting, Preemergence

Apply before, during or after planting, up to a maximum of 2 quarts per acre.

Postemergence (In-crop)

This product may be applied to Roundup Ready winter canola varieties from emergence to canopy closure in the fall and prior to bolting in the spring. Applications made during or after bolting may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Some weeds with multiple germination times, or suppressed (stunted) weeds, or weeds that have overwintered may require sequential applications of this product for control. The second application should be made after some regrowth has occurred and at least 60 days after a previous application of this product.

Single Application—Apply 24 to 32 fluid ounces of this product per acre in the fall. Applications in the fall should be made when weeds are small and actively growing. Use the higher rate in the range when weed densities are high, when weeds have overwintered or when weeds become large and well established. Applications of greater than 16 fluid ounces per acre prior to the 6-leaf stage may result in reduced crop growth in the fall. Avoid spray overlaps. Spray overlaps may result in temporary yellowing and/or growth reduction.

Sequential Applications - Apply 16 to 32 fluid ounces of this product per acre to 2-leaf or larger canola in the fall, followed by a sequential application at the same rate and at a minimum interval of 60 days, but before bolting in the spring. Sequential applications are for early emerging annual weeds and winter emerging weeds such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product will control or suppress most perennial weeds. For some perennial weeds, sequential applications may be required to reduce competition with the crop.

RESTRICTIONS: No more than two over-the-top broadcast applications may be made from crop emergence up to the onset of bolting, and the total in-crop application must not exceed 64 fluid ounces of this product per acre. Applications of greater than 24 fluid ounces per acre prior to the 6-leaf stage

may result in reduced crop growth in the fall. Allow a minimum of 60 days between last application and harvest of canola grain. No waiting period is required between application and open grazing of livestock.

12.4 Roundup Ready Corn 2

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Spot treatment, Preharvest, Post-harvest.

Annual Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from emergence through 48 inches	3 quarts per acre
Maximum preharvest application (See PRECAUTIONS, RESTRICTIONS section for Preharvest)	1 quart per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-planting, Preemergence

Apply alone or in a tank-mixture before, during or after planting.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for alfalfa application. Read and follow label directions for all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to corn: 2,4-D, acetochlor, alachlor, atrazine, carfentrazone-ethyl, clopyraïd, dicamba, diflufenzopyr, dimethenamid, dimethenamid-p, flufenacet, flumetsulam, flumiclorac pentyl ester, isoxaflutole, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, rimsulfuron.

Aim, Axiom, Balance, Balance PRO, Banvel, Bicep MAGNUM, Bicep II MAGNUM, Bicep Lite II MAGNUM, Bullet, Cinch, Cinch ATZ, Clarity, Define, Degree, Degree Xtra, Distinct, Dual II MAGNUM, Epic, Frontier, FulTime, Guardsman MAX, Harness, Harness Xtra, Harness Xtra 5.6L, Homer, Keystone, Keystone LA, Lariat, Linex, Lorox, Marksman, Me-Too-Lacitor II, Micro-Tech, Prowl, Python, Python II, Radius, Resolve, Resource, Stalwart, Stalwart Xtra, Surpass, TopNotch]

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

Apply alone or in tank mixtures postemergence (in-crop) to corn hybrids designated as Roundup Ready Corn 2. When applied as directed, this product controls annual grass and broadleaf *weeds* listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. A postemergence application of this product should be made before the weeds reach a height of 4 inches or before they become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre should be made before the weeds reach a height of 4 inches.

This product may be broadcast over-the-top or applied with drop nozzles to Roundup Ready Corn 2 from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing).

whichever comes first. Drop nozzles are for optimum spray coverage and weed control when corn height is 24 to 30 inches. For corn heights 30 to 48 inches (free standing), apply this product only by ground application using drop nozzles aligned to avoid spraying into the whorls of the corn plants.

TANK MIXTURES: This product may be tank-mixed with the following products. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to corn. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to corn: atrazine, acetochlor, alachlor, clopyralid, dicamba, diflufenzopyr, flumetsulam, flumiclorac pentyl ester, foramsulfuron, halosulfuron-methyl, iodosulfuron-methyl-sodium, mesotrione, nicosulfuron, rimsulfuron, thifensulfuron methyl.

Banvel, Basis, Basis Gold, Bullet, Callisto, Clarity, Degree, Degree Xtra, Distinct, Equip, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Marksman, Micro-Tech, Option, Resolve, Resource, YUKON[®]

Tank Mix Partner	Maximum Height Of Corn For Application
Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6L	11 inches
Bullet* Micro-Tech*	5 inches
Atrazine	12 inches

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet* or Micro-Tech* herbicides.

PRECAUTIONS, RESTRICTIONS: Single in-crop applications of this product should not exceed 3 pints per acre. Sequential in-crop applications of this product from emergence through 48 inches in height must not exceed a total of 3 quarts per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product in-crop and harvest of corn forage or grain. (For applications at preharvest timing, see **Preharvest** section of this label). The use of additional surfactants and other additives, including fertilizers and micro-nutrients, in the spray solution are not used with this product and may result in increased potential for crop injury or reduced yield, and is not used for over-the-top applications, unless otherwise directed in supplemental labeling of fact sheets published separately by Repar-Glypho, LLC..

Preharvest

A single preharvest application of up to 32 fluid ounces per acre of this product may be applied for annual and perennial weed control prior to crop harvest. Make application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

RESTRICTIONS: Do not make a preharvest application of this product if more than a combined total of 64 fluid ounces of this product has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

Postharvest

Apply for weed control after corn harvest. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

12.5 Roundup Ready Cotton

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Selective Equipment (In-Crop), Preharvest.

<u>Annual Maximum Application Rates</u>	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to layby	4 quarts per acre
Maximum preharvest application rate	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." **ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.**

Preplant, At-Planting, Preemergence

Apply before, during or after planting. Tank mixtures with other herbicides listed in Section 8.3 of this label maybe used.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to the emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyridiobac-sodium.

Caparol, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H2O, Stalwart, Staple]

RESTRICTIONS: The maximum quantity of this product that maybe applied for all preplant, at-planting and preemergence applications combined is 5 quarts per acre per season.

Postemergence (in-crop)

Apply over the top Roundup Ready cotton at rates up to 1 quart per acre per application from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). **Applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss.**

TANK MIXTURES: This product may be tank-mixed with the following products and applied over the top of Roundup Ready cotton. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clethodim, fluazifop-P-butyl, metolachlor, s-metolachlor, pyriithiobac-sodium, sethoxydim, quizalofop-P-ethyl. Assure II, Dual MAGNUM, Fusilade, Poast Plus, Select, Stalwart, Staple. Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop). Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting to exposed cotton leaves.]

RESTRICTIONS: The maximum quantity of this product that may be applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. No more than two over-the-top broadcast applications may be made from crop emergence through the 4-leaf (node) stage of development. No more than two applications should be made from the 5-leaf stage through layby. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications. Do not add additional surfactant or additives containing surfactant to this product (other than those contained in any tank-mix product) for over-the-top applications to roundup ready cotton.

Salvage Treatment. Apply after the 4-leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

Selective Equipment (In-crop)

Apply this product using precision post-directed or hooded sprayers at rates up to 1 quart per acre per application to Roundup Ready cotton through layby. At this stage, use post-directed equipment to direct the spray to the base of the cotton plants. Avoid contact of the herbicide spray with the cotton leaves to the maximum extent possible. To minimize spray contact maintain low spray pressure (less than 30 pounds per square inch) and place nozzles in a low position, directing a horizontal spray pattern under the cotton leaves and onto weeds in the row. For best results, make applications while weeds are small (less than 3 inches). See additional instructions on the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

TANK MIXTURES: This product can be tank-mixed with the following products for in-crop application using precision post-directed or hooded sprayers. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: carfentrazone-ethyl, diuron, flumioxazin, fluometuron, linuron, pendimethalin, prometryn, pyriithiobac-sodium. Akin, Caparol, Chateau, Cotoran, Direx, Layby-Pro, Prowl H20, Staple, Valor. Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop)]

RESTRICTIONS: The maximum quantity of this product that may be applied for all in-crop applications from ground-cracking to layby combined is 4 quarts per acre per season. Sequential over-the-top or post-directed applications of this product in-crop must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

Preharvest

Apply for annual and perennial weed control as a broadcast treatment prior to crop harvest after 20 percent boll crack. Apply up to 2 quarts of this product per acre. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Preharvest application is not used for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION: Use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.6 Roundup Ready Flex Cotton

The use of the over-the-top applications described in this section on **other** than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (node) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this section are specific to, and should only be used with, varieties designated as **Roundup Ready Flex cotton**. Do Not combine the instructions in this section, with those in the "Roundup Ready Cotton" section (12.5) of this label, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instructions on labeling for this or other glyphosate-containing product.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence(In-crop), Preharvest.

Annual Maximum Allowable Application Rates

Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to 60 percent open bolls	6 quarts per acre
Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 quarts per acre
Combined total in-crop application from emergence through harvest	6 quarts per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of cotton. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use prior to the emergence of cotton: diuron, fluometuron, metolachlor, s-metolachlor, pendimethalin, prometryn, pyriithiobac-sodium.]

Caparol, Cotoran, Direx, Dual MAGNUM, Prowl, Prowl H20, Stalwart, Staple

Postemergence (In-crop)

Apply to control annual grasses and broadleaf weeds listed on this label in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds is used. This product may be applied postemergence at rates up to 1.5 quarts per acre per application. In addition to broadcast applications, post-directed spray equipment may be used to achieve weed coverage.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready Flex cotton. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to cotton. Read and follow label directions of all products in the tank mixture.

Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to cotton: clothodim, fluazifop-Phutyl, metolachlor, s-metolachlor, pyriithiobac-sodium, quizalofop-p-ethyl, sethoxydim, trifloxysulfuron-sodium.

Assure II, Dual MAGNUM, Envoke, Fusilade, Poast Plus, Select, Stalwart, Staple

Staple may cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop).

Dual Magnum and Stalwart applied over the top of Roundup Ready cotton may cause leaf injury in the form of necrotic spotting in exposed cotton leaves.)

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 1.5 quarts per acre made using ground application equipment. In-crop application rates above 1 quart per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling or leaf necrosis. Do not exceed a maximum rate of 1 quart per acre of this product *when* making applications by air. Between layby and 60 percent open bolls, the maximum combined total rate of this product that may be applied is 2 quarts per acre. The maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre. Do not add additional surfactant or additives containing surfactant to this product for over-the-top applications to Roundup Ready Flex cotton.

Preharvest

Apply up to 2 quarts of this product may be applied for annual and perennial weed control as a broadcast treatment prior to harvest after 60 percent boll crack. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Do not apply preharvest for cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION: use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready Flex Cotton, however, due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors it is impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

12.7 Roundup Ready Soybeans

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Postemergence (In-crop), Preharvest, Post-Harvest.

Annual Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from cracking through flowering (R2 stage soybeans)	3 quarts per acre
Maximum preharvest application rate	1 quart per acre

GENERAL PRECAUTIONS/RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates."

Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

This product may be applied before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with 2,4-D, Banvel or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to soybeans: alachlor, atrazine, *n*-arfenazone-ethyl, chlorimuron ethyl, clethodim, clomazone, cloransulam-methyl, dimethenamid, dimethenamid-p, fenoxypyr, fluazifop-p-butyl, flufenacet, flumetsulam, flumiclorac pentyl ester, flumioxazin, fomesafen, imazaquin, imazethapyr, lactofen, linuron, metolachlor, smetolachlor, metribuzin, pendimethalin, sulfentrazone, tribenuron methyl, quizalofop P-ethyl.

Aim, Amplify, Assure II, Axiom, Authority, Blanket, Boundary, Canopy, Canopy EX, Classic, Cobra, Command, Command Xtra, Domain, Dual MAGNUM, Dual II MAGNUM, FirstRate, Flexstar, Frontier, Fusion, Gangster, Gauntlet, INTRRO, Lexone, Linex, Linuron, Lorox, Lorox Plus, Me-Toolachlor, Micro-Tech, Outlook, Pendimax, Prowl, Prowl H2O, Pursuit, Pursuit Plus, Python, Reflex, Resource, Scepter, Select, Select MAX, Sencor, Spartan, Squadron, Steel, Valor.]

Postemergence (In-crop)

For use of this product to control annual grasses and broadleaf weeds listed on this label in Roundup Ready soybeans. Applications of this product can be made in: from emergence (cracking) through flowering (R2 stage soybeans ends when a pod reaches 5 millimeters (0.2 inches) in length at one of the four uppermost nodes on the main stem with a fully developed leaf. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate recommendations for specific annual weeds. In general, an initial application of 1 quart per acre on 2- to 8-inch tall weeds is used. Weeds will generally be 2 to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 2 quarts per acre in any single in-crop application for control of annual weeds and where dense weed populations exist.

A 1- to 2-quarts per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marehail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds. **IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE ROUNDUP READY SOYBEAN CROP.** To control giant ragweed, 1 quart per acre of this product be applied when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready soybeans. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to soybeans. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to soybeans: acifluorfen, bentazonechlorimuron ethyl, clothodim, imazethapyr, clomansulam-methyl, fenoxypop, fluazifop-pbutyl, flumiclorac pertyl ester, fomesafen, imazamox, imazethapyr, lactofen, pendimethalin, quizalofop P-ethyl, sethoxydem, thifensulfuron-methyl.

Arrow, Assure II, Basagran, Classic, Cobra, Extreme, FirstRate, Flexstar, Fusilade DX, Fusion, Harmony GT XP, Poast, Poast Plus, Pursuit, Pursuit Plus, Raptor, Reflex, Select, Select MAX, Synchrony STS, Targa, Ultra Blazer.]

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in-crop application is 2 quarts per acre. The maximum combined total of this product that can be applied during flowering is 2 quarts per acre.

Preharvest

Apply for weed control prior to harvest of soybeans after pods have set and lost all green color. Up to 1 quart per acre of this product can be applied by aerial or ground application.

PRECAUTIONS, RESTRICTIONS: Care should be taken to avoid excessive seed shatter loss due to ground application equipment. Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

Post-Harvest

Apply this product after harvest of Roundup Ready soybeans. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

12.8 Roundup Ready Sugar Beet

Maximum Annual Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Emergence to 8-leaf stage	2.5 quarts per acre
Between 8-leaf stage and canopy closure	2 quarts per acre

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop).

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. See the "GENERAL INFORMATION" section of this label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-Planting, Preemergence

Apply before, during or after planting.

TANK MIXTURES: This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of sugar beet. Read and follow label directions of all products in the tank mixture.

Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to sugar beet: dimethenamid, S-metolachlor.

Dual MAGNUM, Frontier]

Postemergence (In-Crop)

Apply this over-the-top of Roundup Ready sugar beets for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate recommendations for specific annual weeds. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready sugar beet. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to sugar beet. Read and follow label directions of all products in the tank mixture.

Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to sugar beet: clethodim, clopyralid, desmedipham, ethofumesate, phenmedipham, quizalofop-p-ethyl, sethoxydim, trisulfuron-methyl.

Assure II, Batamix, Betanex, Norton SC, Poast, Progress, Select, Stinger, Upbeat]

RESTRICTIONS: The combined total application from crop emergence through harvest must not exceed 4.5 quarts per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 1.5 quarts per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 1 quart per acre. Allow a minimum of 30 days *between* last application and sugar beet harvest.

13.0 NON-CROP USES AROUND THE FARMSTEAD

TYPES OF APPLICATIONS: General Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stump, Habitat Management.

13.1 General Weed Control, Trim-And-Edge

Apply to control annual weeds, perennial weeds and woody brush found in any part of the farmstead, including around building foundations and equipment storage areas, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, rangeland, rights-of-way, shelterbelts, and prior to planting landscape ornamentals. Refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" in this label for treatment rates. For application through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "Annual Weeds -- Hand-Held or Backpack Equipment" section of this label for further instructions.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product used is registered for treatment of these non-crop sites. Refer to the individual product labels for approved farmstead sites and application rates. For more information, see the "TANK MIXTURES" and "TANK MIXING PROCEDURES" sections of this label.

Arsenal	Plateau
Barricade 65WG	Princep DF
Crossbow L	Princep 4L
Diuron	Ronstar 50 WP
Dicamba	Sahara
Endurance	Simazine

Escort	Surflan AS
Goal 2XL	Surflan WDG
Karmex DF	Telar
Krovar I DF	Transire
Outrider	Vanquish
Oust	Velpar DF
Pendimethalin	Velpar L
Pendulum 3.3 EC	2,4-D
Pendulum WDG	

This product plus dicamba tank mixtures may not be applied by air in California.

13.2 Greenhouse/Shadhouse

To control weeds in and around greenhouses and shade houses.

PRECAUTIONS, RESTRICTIONS: Desirable vegetation must not be present during application. Air circulation fans must be turned off. Do not use in residential greenhouses.

13.3 Chemical Mowing

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Use 16 fluid ounces of this product per acre when treating Bermudagrass. Use 64 fluid ounces of this product per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

PRECAUTIONS, RESTRICTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

13.4 Cut Stump

TYPES OF APPLICATIONS: Treating cut stumps in any non-crop site listed on this label.

Apply this product for control of growth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Red, giant
Eucalyptus	Salt-cedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow
Pine, Austrian	

PRECAUTIONS: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

13.5 Habitat Management

TYPES OF APPLICATIONS: Habitat Restoration and Maintenance, Wildlife Food Plots.

Habitat Restoration and Maintenance

Apply to control other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species, including Roundup ready canola, may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage. For specific product instructions for Roundup Ready canola wildlife food plots, see the "ROUNDUP READY CANOLA" section of this label.

RESTRICTIONS: Do not process treated Roundup ready canola seeds from Roundup ready canola wildlife food plots for food. Do not graze or feed treated Roundup Ready canola from wildlife food plots to livestock. There are no rotational restrictions for planting any wildlife food species or for allowing native species to repopulate the area following applications of this product.

14.0 ANNUAL WEEDS RATE SECTION

When water carrier volumes are between 16 and 40 gallons per acre for ground applications and between 6 and 15 gallons per acre for aerial applications, the following use rates will control the annual weeds listed in the table below:

- 1 quart per acre – grass and broadleaf annual weeds less than 6 inches in height or circumference and vines less than 3 inches in length.
- 3 pints per acre – grass and broadleaf annual weeds 6 to 12 inches in height or circumference and vines 3 to 6 inches in length.
- 2 quarts per acre – grass and broadleaf annual weeds greater than 12 inches in height or circumference and vines greater than 6 inches in length.

WHEN WATER CARRIER VOLUMES ARE BETWEEN 3 AND 15 GALLONS PER ACRE FOR GROUND APPLICATIONS AND BETWEEN 3 AND 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS, USE THE RATES SPECIFIED FOR INDIVIDUAL WEEDS AS FOLLOW IN THE "ANNUAL WEEDS RATE TABLE (Alphabetically by Species)".

Apply to actively growing annual weeds. Annual weeds are generally easiest to control when they are small.

Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Maximum size refers to the maximum plant height, length of runners for vines, or circumference of rosette plants in inches.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment. This product may be used up to 64 fluid ounces per acre where heavy weed densities exist.

ANNUAL WEEDS RATE TABLE
(Alphabetically by Species)

WEED SPECIES	RATE (Fluid Ounces Per Acre)				
	16	24	32	40	48
	(Maximum Size (in inches))				
Ammannia, purple	3	6	12	-	18
Anode, spurred	-	2	3	5	8
Barley	18	18+	-	-	-
Barnyardgrass	-	3	6	7	9
Basile, fivehook	-	-	6	-	-
Beggarweed, Florida	-	5	8	-	-
Bittercress	12	20	-	-	-
Bluegrass, annual	10	-	-	-	-

Bluegrass, bulbous	6	-	-	-	-
Brome, downy ^{1,2}	6	12	-	-	-
Brome, Japanese	6	12	24	-	-
Browntop panicum	6	8	12	-	24
Buckwheat, wild ³	-	1	2	-	-
Burcucumber	-	6	12	-	18
Buttercup	12	20	-	-	-
Carolina geranium	-	-	4	-	9
Carpetweed	-	6	12	-	-
Cheat ²	6	20	-	-	-
Chervil	20	-	-	-	-
Chickweed	-	12	18	-	-
Cocklebur	12	18	24	-	36
Copperleaf, hophornbeam	-	2	4	-	6
Copperleaf, Virginia	-	2	4	-	6
Coreopsis, plains	-	6	12	-	18
Corn, volunteer	6	12	20	-	-
Corn speedwell	12	-	-	-	-
Crabgrass	3	6	12	-	-
Crowfootgrass	-	-	6	-	12
Cutleaf evening primrose	-	-	3	-	6
Devilsclaw (unicorn plant)	-	3	6	-	-
Dwarf dandelion	12	-	-	-	-
Eastern manna grass	8	12	-	-	-
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
Falsedandelion	-	20	-	-	-
Falseflax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	6	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail, giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	6	12	-	-	-
Goosegrass	-	3	6	-	12
Grain sorghum (mild)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundsel, common	-	6	10	-	-
Hemp sesbania	-	2	4	6	8
Henbit	-	-	6	-	12
Horseweed/ Maretail (<i>Conyza canadensis</i>)*	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnsongrass, seedling	6	12	18	-	24
Junglerice	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia ⁴	-	3 to 6	12	-	-
Lambsquarters*	-	6	12	-	20
Little barley	6	12	-	-	-

London rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morningglory, annual (<i>Ipomoea spp</i>)	-	-	3	-	6
Mustard, blue	6	12	18	-	-
Mustard, tansy	6	12	18	-	-
Mustard, tumble	6	12	18	-	-
Mustard, wild	6	12	18	-	-
Nightshade, black	-	4	6	-	12
Nightshade, hairy	-	4	6	-	12
Oats	3	6	18	-	-
Pigweed, Palmer*	-	12	18	24	-
Pigweed species*	-	12	18	24	-
Prickly lettuce	-	6	12	-	-
Purslane	-	-	3	-	6
Ragweed, common*	-	6	12	-	18
Ragweed, giant*	-	6	12	-	18
Red rice	-	-	4	-	-
Rye, volunteer/cereal ²	6	18	18+	-	-
Ryegrass species*	-	-	6	-	12
Sandbur, field	6	12	-	-	-
Sandbur, longspine	6	12	-	-	-
Shattercane	6	12	20	-	-
Shepherd's-purse	6	12	-	-	-
Sicklepod	-	2	4	-	8
Signalgrass, broadleaf	-	3	6	7	9
Smartweed, ladythumb	-	-	6	-	9
Smartweed, Pennsylvania	-	-	6	-	9
Sowthistle, annual	-	-	6	-	12
Spanishneedles	-	-	6	-	12
Speedwell, purslane	12	-	-	-	-
Sprangletop	6	12	20	-	-
Spurge, prostrate	-	6	12	-	-
Spurge, spotted	-	6	12	-	-
Spurry, umbrella	6	-	-	-	-
Stinkgrass	-	12	-	-	-
Sunflower	12	18	-	-	-
Swinecress	-	5	12	-	-
Tarweed/ Prickly sida	-	2	4	-	6
Texas panicum	6	8	12	-	24
Thistle, Russian ⁵	-	6	12	-	-
Velvetleaf	-	-	6	-	12
Virginia pepperweed	-	18	-	-	-
Waterhemp*	-	-	6	-	12
Wheat ²	6	12	18	-	-
Wheat, (overwintered)	-	6	12	-	18
Wild oats	3	6	18	-	-
Wild proso millet	-	6	12	-	18
Witchgrass	-	12	-	-	-
Woolly cupgrass	-	6	12	-	-
Yellow rocket	-	12	20	-	-

¹ For control of downy brome in no-till systems, use 24 fluid ounces per acre.

² Performance is better if application is made before this weed reaches the boot stage of growth.

³ Use 24 fluid ounces per acre of this product to control wild buckwheat in the cotyledon to 2-leaf stage. Use 32 fluid ounces per acre to control 2- to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 32 fluid ounces followed by 32 fluid ounces of this product per acre.

⁴ Do not treat kochia in the button stage.

⁵ Control of Russian thistle may vary based on environmental conditions and spray coverage. Whenever possible, a tank mixture with 2,4-D as described below may improve control.

* A glyphosate resistant biotype has been confirmed. For additional information, refer to the "WEED RESISTANCE MANAGEMENT" section of this label. You may also visit on the Internet www.weedscience.org or www.weedresistancemanagement.com, or contact your Repar-Glypho, LLC, representative.

14.1 Annual Weeds — Tank Mixtures with 2,4-D or Dicamba or Tordon 22K

Better control of certain tough weeds can be achieved by tank mixing this product with 0.25 pound of dicamba, 0.5 pound of 2,4-D or 1 to 2 fluid ounces of Tordon 22K per acre. These other herbicides, combined with the rates of this product specified in the "ANNUAL WEEDS RATE TABLE" above, will control the following weeds up to the maximum height or length indicated: 6" -- prickly lettuce, maretail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Tordon 22K only); 12" -- cocklebur, lambsquarters, pigweed, Russian thistle (2,4-D only).

For better control of common ragweed, giant ragweed, Pennsylvania smartweed, or velvetleaf with a maximum height of 6-inches, tank mix this product with 0.5 pound of 2,4-D per acre.

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA.

14.2 Annual Weeds—Hand-Held or Backpack Equipment

For control of weeds listed in the "ANNUAL WEEDS RATE TABLE" above, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 -percent solution.

For best results, use a 2-percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 5-percent solution for annual and perennial weeds and a 5- to 10-percent solution for woody brush and trees.

14.3 Annual Weeds — Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

Applications of 24 to 28 fluid ounces of this product plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 28 ounces for control), Downy brome, Green foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass and Kochia (add 0.12 pound of dicamba for control).

Ensure that the specific product is registered for application at the desired site. Follow all precautions and limitations on the tank mix product's label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines.

15.0 PERENNIAL WEEDS RATE SECTION

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage. Best results are obtained when soil moisture is adequate for active weed growth.

PERENNIAL WEEDS RATE TABLE

(Alphabetically by Species)

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alfalfa Apply after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.	1 - 2	3 - 10	2%
Alligatorweed For partial control, apply when most of the plants are in bloom. Repeat applications will be required to achieve control.	4	3 - 20	1.5%
Anise (fennel) ¹ Apply as a spray-to-wet treatment.	-	-	1-2%
Bahiagrass²	3-5	3-20	2%
Bentgrass For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is used for best results.	3-5	10-20	2%
Bermudagrass For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to achieve control.	3 - 5	3 - 20	2%
Bermudagrass, water (knotgrass) Apply 3 pints of this product in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field. Fall applications only: Apply 1 quart this product in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length. This product is not registered in California for use on water Bermudagrass.	1 - 1.5	5 - 10	2%
Bindweed, field Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in	0.5-5	3 - 20	2%

late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 2 quarts of this product plus 0.5 pound of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. Irrigate at least once to promote active bindweed growth.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky	1 - 2	3 - 40	2%
----------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas	3 - 5	3 - 40	2%
------------------------	--------------	---------------	-----------

Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenfern	3 - 4	3 - 40	1 - 1.5%
--------------------	--------------	---------------	-----------------

Apply to fully expanded fronds that are at least 18 inches long.

Bromegrass, smooth	1 - 2	3 - 40	2%
---------------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf	---	3 - 20	2%
-----------------------------	------------	---------------	-----------

For control, apply 2 quarts of this product plus 0.5 pound of dicamba per acre. For partial control, apply 1 quart of this product plus 1/4 pound of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed²	2 - 3	3 - 40	2%
--------------------------------------	--------------	---------------	-----------

Cattail²	3 - 5	3 - 40	2%
----------------------------	--------------	---------------	-----------

Clover; red or white¹	3 - 5	3 - 20	2%
---	--------------	---------------	-----------

Also for control, apply 16 to 32 fluid ounces of this product plus 0.5 to 1 pound of 1,4-D, in 3 to 10 gallons of water per acre.

Cogongrass	3 - 5	10 - 40	2%
-------------------	--------------	----------------	-----------

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to achieve control.

Dallisgrass²	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Dandelion¹	3 – 5	3 – 40	2%
------------------------------	--------------	---------------	-----------

Also for control, apply 1 pint of this product plus 0.5 of 2,4-D, in 3 to 10 gallons of water per acre.

Dock, curly¹	3 – 5	3 – 40	2%
--------------------------------	--------------	---------------	-----------

Also for control, apply 1 to 2 pints of this product plus 0.5 to 1 pound of 2,4-D, in 3 to 10 gallons of water per acre.

Dogbane, hemp	4	3 – 40	2%
----------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.

For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre for ground applications, and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Fescue (except tall)¹	3 – 5	3 – 20	2%
---	--------------	---------------	-----------

Fescue, tall	1 – 3	3 – 40	2%
---------------------	--------------	---------------	-----------

Apply 3 quarts of this product per acre when most plants have reached boot-to-early seedhead stage of development.

Fall applications only: Apply 1 quart of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.

Guineagrass	2 - 3	3 - 40	1%
--------------------	--------------	---------------	-----------

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 2 quarts of this product per acre for control. In the flatwoods region of Florida, 3 quarts of this product per acre are required for control.

Horsenettle¹	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Horseradish	4	3 – 40	2%
--------------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Iceplant¹	---	---	1.5 – 2%
-----------------------------	------------	------------	-----------------

Thorough coverage is necessary for best control.

Jerusalem artichoke¹	3 – 5	3 – 20	2%
--	--------------	---------------	-----------

Johnsongrass	0.5 – 3	3 – 40	1%
---------------------	----------------	---------------	-----------

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 1 quart of this product per acre.

For burndown of Johnsongrass, apply 1 pint of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression)--Apply a 1-percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass	2 – 3	3 – 40	2%
--------------------	--------------	---------------	-----------

Knapweed	4	3 – 40	2%
-----------------	----------	---------------	-----------

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Lantana	--	--	1-1.25%
----------------	-----------	-----------	----------------

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Lespedeza	3 – 5	3-20	2%
------------------	--------------	-------------	-----------

Milkweed, common	3	3-40	2%
-------------------------	----------	-------------	-----------

Apply when most plants have reached the late bud to flower stage of growth.

Muhly, wirestem	1-2	3-40	2%
------------------------	------------	-------------	-----------

Use 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications, or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.

Mullein, common¹	3 – 5	3 – 20	2%
------------------------------------	--------------	---------------	-----------

Napiergrass²	3 – 5	3 – 20	2%
--------------------------------	--------------	---------------	-----------

Nutsedge, purple or yellow	0.5-3	3 - 40	1 - 2%
-----------------------------------	--------------	---------------	---------------

Apply 3 quarts of this product per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for longterm control of ungerminated tubers.

Sequential applications: 1 to 2 quarts of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.

For partial control of existing plants, apply 1 pint to 2 quarts of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

Orchardgrass	1 - 2	3 - 40	2%
---------------------	--------------	---------------	-----------

Apply 2 quarts of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of this product in 3 to 10 gallons

of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications, and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

Pampasgrass -- -- 1.5 - 2%

Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Paragrass² 3 - 5 3 - 20 2%

Phragmites 3 - 5 10 - 40 1 - 2%

For partial control and best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may result in reduced control. Due to the dense nature of this vegetation that may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to achieve control. Visual control symptoms will be slow to develop.

Poison hemlock -- -- 1 - 2%

For hand-held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Thorough coverage is necessary for best control.

Pokeweed, common 1 3-40 2%

Apply to actively growing plants up to 24 inches tall.

Quackgrass 1 - 3 3 - 40 2%

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of this product. Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.

In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

Redvine 0.75-2 5-10 2%

For suppression, apply 1.5 pints of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, giant -- -- 2%

Best results are obtained when applications are made in late summer to fall.

Ryegrass, perennial 1 - 3 3 - 40 1%

In annual cropping systems, apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In non-crop areas, or areas where annual tillage (no-till) is not practiced, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1 quart of this product per acre.

Smartweed, swamp¹	3-5	3-40	2%
Also for control, apply 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall.			
Sowthistle, perennial	2-3	3-40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
Spurge, leafy		3-10	2%
For suppression, apply 1 pint of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most plants are 12 inches tall.			
Starthistle, yellow	2	10-40	2%
Best results are obtained when applications are made during the rosette, bolting and early flower stages.			
Sweet potato, wild			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, artichoke			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, Canada	2-3	3-40	2%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
For suppression: in the spring, apply 1 quart of this product, or 1 pint of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.			
Timothy²	2-3	3-40	2%
Torpedograss	4-5	3-40	2%
For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will ^{not} be required to achieve control. Fall treatments must be applied before frost.			
Trumpet creeper	2	5-10	2%
For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.			
Vaseygrass²	3-5	3-20	2%
Velvetgrass²	3-5	3-20	2%

Wheatgrass, western² 2-3 3-40 2%

¹ Apply when most plants have reached the early bud stage of growth.

² Apply when most plants have reached the early heading stage of growth.

16.0 WOODY BRUSH AND TREES RATE SECTION

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Unless otherwise directed, apply broadcast treatments in 3 to 40 gallons of water per acre. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

WOODY BRUSH AND TREES RATE TABLE
(Alphabetically by species)

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alder	3-4	3-40	1-1.5%
Ash ¹	2-5	3-40	1-2%
Aspen, quaking	2-3	3-40	1-1.5%
Bearna ¹ (Bearclover)	2-5	3-40	1-2%
Beech ¹	2-5	3-40	1-2%
Birch	2-3	3-40	1-1.5%
Blackberry	3-4	10-40	1-1.5%

Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 0.75 percent solution of this product. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

Blackgum	2-5	3-40	1-2%
Bracken	2-5	3-40	1-2%
Broom; French, Scotch			1.5-2%
Buckwheat, California, 2			1-2%
Cascara ¹	2-5	3-40	1-2%
Catsclaw	---		1-1.5%
Ceanothus ¹	2-5	3-40	1-2%
Chamise ²			1%

Cherry; bitter, black, pin	2-3	3-40	1-5%
Coyote brush	--	--	1.5-2%
Apply when at least 50 percent of the new leaves are fully developed.			
Dogwood'	2-5	3-40	1-2%
Elderberry	2-3	3-40	1-1.5%
Elm'	2-5	3-40	1-2%
Eucalyptus	--	--	2%
For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.			
Florida holly (Brazilian Peppertree) 1	2-5	3-40	1-2%
Gorse'	2-5	3-40	1-2%
Hasardia*2	--	--	1-2%
Hawthorn	2-3	3-40	1-1.5%
Hazel	2-3	3-40	1-1.5%
Hickory'	2-5	3-40	1-2%
Honeysuckle	3-4	3-40	1-1.5%
Hornbeam, American'	2-5	3-40	1-2%
Kudzu	4-5	3-40	2%
Repeat applications may be required to achieve control.			
Locust, black'	2-4	3-40	1-2%
Madrone resprouts'	--	--	2%
Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.			
Manzanita'	2-5	3-40	1-2%
Maple, red	2-4	3-40	1-1.5%
Apply a 1- to 1.5-percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre.			
Maple, sugar	--	--	1-1.5%
Apply when at least 50 percent of the new leaves are fully developed			
Monkey flower ²	--	--	1-2%
Oak; black, white'	2-4	3-40	1-2%
Oak, post	3-4	3-40	1-1.5%
Oak; northern,	--	--	1-1.5%
Apply when at least 50 percent of the new pin leaves are fully developed			
Oak; southern red	2-3	3-40	1-1.5%
Persimmon'	2-5	3-40	1-2%
Pine	2-5	3-40	1-2%
Poison ivy/Poison oak	4-5	3-40	2%
Repeat applications may be required to achieve control. Fall treatments must be applied before leaves lose green color.			
Poplar, yellow'	2-5	3-40	1-2%
Redbud, eastern	2-5	3-40	1-2%

Rose, multiflora	2	3-40	1 %
Treatments should be made prior to leaf deterioration by leaf-eating insects.			
Russian olive ¹	2-5	3-40	1 -2%
Sage, black ¹			1%
Sage, white ¹	2-5	3-40	1 -2%
Sagebrush, California ²			1%
Salmonberry	2-3	3-40	1 -1.5%
Salt-cedar	2-5	3-40	1 -2%
Sassafras ¹	2-5	3-40	1 -2%
Sourwood ¹	2-5	3-40	1 -2%
Sumac; poison, smooth, winged ¹	2-4	3-40	1 -2%
Sweetgum	2-3	3-40	1 -1.5%
Swordfern ¹	2-5	3-40	1 -2%
Tallowtree, Chinese ¹			1%
Tan oak resprouts ¹	--	--	2%
Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.			
Thimbleberry	2-3	3-40	1 -1.5%
Tobacco, tree ¹			1 -2%
Trumpet creeper	2-3	3-40	1 -1.5%
Vine maple ¹	2-5	3-40	1 -2%
Virginia creeper	2-5	3-40	1 -2%
Waxmyrtle, Southern ¹	2-5	3-40	1 -2%
Willow	3-4	3-40	1 -1.5%

¹ Partial Control

² Thorough coverage of foliage is necessary for best results.

17.0 LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. To the extent consistent with applicable law, this warranty is also subject to the conditions and limitations stated on this labeling.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

For in-crop (over-the-top) uses on Roundup Ready crops, crop safety and weed control performance are not warranted by Repar-Glypho, LLC when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OF DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR , AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY TO THE FULLEST EXTENT PERMITTED BY LAW. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Amplify, Bullet, Degree, Farmsource, Harness, Lariat, , Micro-Tech, Monsanto and Vine symbol, and Permit are trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

II. MAIN LABEL FOR INDUSTRIAL, TURF, ORNAMENTAL USES

GLYPHOSATE 41% SL

The complete broad-spectrum postemergence herbicide for non-crop, industrial, turf and ornamental weed control.

Complete Directions for Use

EPA Reg. No. 86004-

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product. Use only according to label instructions.

Not all products on this label are registered for use in California. Check the registration status of each product in California before using.

[Container Label Statement (5 gallons or smaller):]

Do not use this container for residential use.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

[Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A REPAR-GLYPHO, LLC. REPACKAGING OR TOLL REPACKAGING AGREEMENT.

[Non-Refillable Container Label Statement]:

THIS IS AN END-USE PRODUCT. REPAR-GLYPHO, LLC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

1	1.0 INGREDIENTS
2	2.0 IMPORTANT PHONE NUMBERS
3	3.0 PRECAUTIONARY STATEMENTS 3.1 Hazards to Humans and Domestic Animals 3.2 Environmental Hazards 3.3 Physical or Chemical Hazards
4	4.0 STORAGE AND DISPOSAL
5	5.0 GENERAL INFORMATION (How This Product Works) 5.1 Weed Resistance Management 5.2 Management Recommendations for Glyphosate Resistant Weed Biotypes
6	6.0 MIXING 6.1 Mixing with Water 6.2 Tank Mixing 6.3 Tank Mixing Procedure 6.4 Mixing for Hand-Held Sprayers 6.5 Surfactants 6.6 Colorants or Dyes
7	7.0 APPLICATION EQUIPMENT AND TECHNIQUES 7.1 Aerial Equipment 7.2 Ground Broadcast Equipment 7.3 Hand-Held or Backpack Equipment 7.4 Selective Equipment 7.5 Injection Systems 7.6 CDA Equipment
8	8.0 SITE AND USE DIRECTIONS 8.1 Cut Stamp 8.2 Forestry Site Preparation 8.3 General Non-crop Areas and Industrial Sites 8.4 Habitat Management 8.5 Hollow Stem Injection 8.6 Injection and Frill (Woody Brush and Trees) 8.7 Non-food Tree, Shrub, or Vine Production Sites 8.8 Railroads 8.9 Roadsides 8.10 Rangelands 8.11 Utility Sites 8.12 Conservation Reserve program (CRP) 8.13 Grass Seed or Sod Production 8.14 Pastures
9	9.0 WEEDS CONTROLLED 9.1 Annual Weeds 9.2 Perennial Weeds 9.3 Woody Brush and Trees
10	10.0 LIMIT OF WARRANTY AND LIABILITY

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....	41.0%
OTHER INGREDIENTS (Including surfactant).....	59.0%
	100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

(To be added later)

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep Out of Reach of Children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED	Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical Emergency Assistance, call the National Pesticide Information Center 1-800-858-7378.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

3.2 Environmental Hazards

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wastewaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published REPAR-GLYPHO, LLC Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. PESTICIDE STORAGE: Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

See container label for additional CONTAINER DISPOSAL instructions.

[Container Label Storage and Disposal Statements]:

[FOR REFILLABLE PORTABLE MINI-BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If this container is damaged, leaking, or obsolete, or to obtain information about recycling portable refillable containers, contact: Repar-Glypho, LLC.

Users: When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling, if available.

Refiners: Do not reuse this container except for refill in accordance with a valid Monsanto agreement for Repackaging or Toll Repackaging. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. Do not reuse this container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, or worn-out threads / closure devices. Final disposal must be in compliance with State and local regulations. If not refilled, returned or recycled, triple rinse or pressure rinse, and offer for recycling or reconditioning, if possible. If burned, stay out of smoke.

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and place in a pesticide container recycling program, if available, or dispose of in a sanitary landfill, or by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke.

[FOR ONE-WAY PLASTIC DRUMS]

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse the container, then puncture and place into a container recycling program, if available, or dispose of in a sanitary landfill, or by incineration by burning, or, if allowed by state and local authorities. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

(How This Product Works)

Product Description: This product is a postemergent, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visual effects are gradual wilting and yellowing of the plant, which advances to complete browning of above ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at later growth stages approaching maturity.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Application Rates: The combined total of all treatments must not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate as the active ingredient, whether applied separately

or as mixtures. Calculate the application rates and ensure that the total use of this and other glyphosate containing products does not exceed stated maximum use rates. See the "INGREDIENTS" section of this label for necessary information.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

5.1 Weed Resistance Management

GROUP	9	HERBICIDE
-------	----------	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or by using other cultural or mechanical practices.

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management practices:

- Scout your fields before and after herbicide applications.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Utilize the label rate for the most difficult to control weed in your field. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture recommendations that encourage application rates of this product below the label recommendations.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
-
-

- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Reap-Glypho, LLC representative, local retailer, or county extension agent.

5.2 Management Practices For Glyphosate Resistant Biotypes

Note: Appropriate testing is critical in order to determine if a weed is resistant to glyphosate. Contact your Reap-Glypho, LLC representative to determine if resistance has been confirmed to any particular weed biotype in your area, or you can visit www.weedresistancemanagement.com or www.weedscience.org. For more information see the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" of this label.

Control recommendations for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling or fact sheets for this product and can be obtained from your local retailer or Reap-Glypho, LLC representative.

As the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Reap-Glypho, LLC Company is not responsible for any losses that may result from the failure of this product to control glyphosate resistant weeds biotypes.

The following good agronomic practices should be followed to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

6.0 MIXING

Spray solutions of this product should be mixed, stored, and applied using only clean stainless steel, aluminum, plastic, fiberglass, plastic or plastic-lined steel containers. Clean sprayer parts immediately after using this product by thoroughly flushing with water. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations.

6.1 Mixing with Water

NOTE: Product performance may be significantly reduced if water containing soil sediment is used as a carrier. Do not mix this product with water from ponds and ditches that is visibly muddy or murky.

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of clean water. Add the amount of this product near the end of the filling process and mix well. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

6.2 Tank Mixing

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

When this label gives directions for use of a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, or dicamba, the user is responsible for ensuring that the specific application is included on the label of the specific products used in the tank mixture. Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions and use according to the most restrictive label requirements. Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly used in this label. Mixing this product with herbicides or other materials on this label may result in reduced performance.

6.3 Tank Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the amount of this product. If needed, add nonionic surfactant before completing the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" and "GENERAL INFORMATION" sections for additional precautions.

6.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product indicated in the following table with water:

Spray Solution

Desired Volume	Amount of GLYPHOSATE 41% SL					
	0.5%	1 %	1.5%	2%	5%	10%
1 gal	0.7 oz	1.3 oz	2 oz	2.7 oz	6.5 oz	13 oz
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

For backpack, knapsack or pump-up sprayers, the appropriate amount of this product be mixed with water in a larger container, and then used to fill the sprayer.

6.5 Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution, unless otherwise directed. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

6.6 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to spray solutions of this product. However, they can reduce product performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system. Avoid direct application to any body of water.

APPLY SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial application.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversion" sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sunsets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use rates of this herbicide in 3 to 25 gallons of water per acre unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by Reap-Glypho, LLC, for this product.

Drift reduction additives may be used. When a drift reduction additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application --to avoid streaking and uneven or overlapped application, use appropriate marking devices.

Aircraft Maintenance; Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART.** LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may prevent corrosion.

7.2 Ground Broadcast Equipment

Apply at rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label for this product. As density of weeds increases, spray volume should be increased to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

7.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For low-volume directed spray applications, spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

7.4 Selective Equipment

This product may be diluted in water and applied through shielded applicators, hooded sprayers, wiper applicators or sponge bars to weeds listed on this label that are growing in any specified non-crop site.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators and Sponge Bars

Wiper applicators are devices that physically wipe the appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust the height of the applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper applicators when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from the use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators—Solutions ranging from 33 to 75 percent of this product in water may be used.

For Panel Applicators and pressure-feed systems—Solutions ranging from 33 to 100 percent of this product in water may be used.

When applied, this product **CONTROLS** the following weeds:

Corn, volunteer	Sicklepod
Panicum, Texas	Spanishneedles
Rye, common	Starbur, bristly Shattercane

When applied, this product **SUPPRESSES** the following weeds:

Beggarweed, Florida	Ragweed, common
Bermudagrass	Ragweed, giant
Dogbane, hemp	Smutgrass
Dogfennel	Sunflower
Guineagrass	Thistle, Canada
Johnsongrass	Thistle, musk
Milkweed	Vaseygrass
Nightshade, silverleaf	Velvetleaf
Pigweed, redroot	

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems.

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 INDUSTRIAL, TURF AND ORNAMENTAL SITES

Unless otherwise specified, applications may be made to control any weeds listed in the "WEEDS CONTROLLED" section 9.0 of this label. Refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" section for detailed instructions on different application methods.

8.1 Cut Stumps

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100-

percent solution of this product to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Salcedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow Pine, Austrian

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2 Forestry Site Preparation

This product is used for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product is also used in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is used in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

Use higher rates of this product for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

This product may be used in a tank mix with the following products for forestry site preparation.

Arsenal Applicators Concentrate
Chopper
Escort or Escort XP
Garlon 3A
Garlon 4A
Landmark XP
Oust or Oust XP
Westar

For control of herbaceous weeds, use the lower tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher tank mixture rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release, unless otherwise specified on this label for this product.

8.3 General Non-crop Areas, Industrial Sites

Use in areas such as airports, apartment complexes commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, rights-of-way, roadsides, shadehouses, sod or turf seed farms, storage areas, sports complexes, substations, turfgrass areas, utility sites, warehouse areas, and wildlife management areas

General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general non-crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product is registered for use on such non-crop sites. Refer to these products' labels for approved non-crop sites and application rates.

2,4-D	Goal 2XL	Princep TM 4L
Arsenal TM	Karmex TM DF	Princep TM DF
atrazine	Krovar TM 1 DF	Ronstar TM 50WP
Barricade TM 65WG	Landmark II MP	Sahara TM
Clarity	Milestone	simazine
Crossbow L	Oust	Surflan TM AS
dicamba	Oust XP	Surflan WDG
diuron	Outrider	Telar TM
Endurance TM	Pendulum WDG	Transline
Escort TM	Pendulum TM 3.3 EC	Vanquish TM
Gallery 75 DF	pendimethalin	Velpar DF
Garlon 4	Platacu TM	Velpar L
Garlon TM 3A	Poast	

This product plus dicamba tank mixtures may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 1 to 2 quarts of this product plus 2 to 4 ounces of Oust or Oust XP per acre.

Bahiagrass	Johnsongrass
Bermudagrass	Poorjoe
Broomsedge	Quackgrass
Dallisgrass	Vaseygrass
Dock, curly	Vervain, blue Dogfennel
Fescue, tall	

Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 8 to 64 fluid ounces of this product per acre. Apply 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust or Oust XP in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 16 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust or Oust XP in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate was greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

8.4 Habitat Management

Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.5 Hollow Stem Injection

This product may be applied through hand-held injection devices that deliver amounts of this product into targeted hollow-stem plants growing in any non-crop site specified on this label.

For control of the following hollow-stem plants, follow the use instructions below:

Japanese Knotweed, *Polygonum cuspidatum*

Inject 5 mL per stem of this product between second and third internode.

Bohemian Knotweed, *Polygonum bohemicum*

Inject 5 mL per stem of this product between the second and third internode.

Giant Hogweed, *Heracleum mantegazzianum*

Inject one leaf cane per plant 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Poison Hemlock, *Conium maculatum*

Inject one leaf cane per plant 10 to 12 inches above the root crown with 5 mL of a 5% v/v solution of this product.

Field Horsetail, *Equisetum arvense*

Inject one segment above the root crown with 0.5 mL per stem of this product. Use a small syringe that calibrates to this rate.

Canada Thistle, *Cirsium arvense*

Cut 8 to 9 of the tallest plants at bud stage in 1 clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.5 mL per stem of this product is injected into the stem.

Note: The combined total for all treatments must not exceed 10.6 quarts of this product per acre. At 5 mL per stem, 7 quarts should treat approximately 1300 stems per acre.

8.6 Injection and Frill (Woody Brush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 0.04 fluid ounce (1 mL) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50- to 100-percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100-percent concentration of this product. For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:

Control

Oak
Poplar
Sweetgum
Sycamore

Partial Control

Black gum
Dogwood
Hickory
Maple, red

8.7 Site Preparation for Planting of Non-Food Trees and Shrubs

Apply this product for general weed control prior to planting or around established ornamentals, or any woody tree, shrub, or vine species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce or yew, in any production site.

UNLESS OTHERWISE DIRECTED, THIS PRODUCT IS NOT FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be taken to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

TYPES OF APPLICATIONS: Site Preparation, Post-directed, Trim-and-edge, Wiper Application Site Preparation

This product may be used prior to planting any tree, shrub, or vine in an ornamental, nursery, or production setting, including Christmas tree species.

Post-directed, Trim-and-edge

This product may be used as a post-directed spray around established woody species, or to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a production setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Wiper Application

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs, or vines. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

8.8 Railroads

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along railroads.

Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

TANK MIXTURES: This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments, provided that the specific product is registered for use on such non-crop sites:

Arsenal	Oust XP
atrazine	Outrider
dicamba	Sahara DG
diuron	simazine
Escort	Spike™ 80 DF
Escort XP	Telar DF
Garlon 3A	Transline
Garlon 4	Vanquish
Hyvar™ X	Velpar DF
Hyvar X-L	Velpar L
Krovar I DF	2,4-D
Oust	

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.75- to 2-percent solution of this product when using high-volume spray-to-wet applications. Apply a 5- to 10-percent solution of this product when using low volume directed sprays for spot treatment.

TANK MIXTURES: This product may be mixed with the products listed above in this section for enhanced control of woody brush and trees along railroads, provided that the specific product is registered for use on such non-crop sites.

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestein, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestein, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpet creeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not done, since severe injury may occur.

8.9 Roadsides

All of the methods of application described in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section may be utilized along roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails, Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

TANK MIXTURES: This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments:

atrazine	Outrider
Clarity	Pendulum 3.3 EC
Crossbow I.	Pendulum WDG
dicamba	Plateau
diuron	Princep DF
Endurance	Princep 4L
Escort	Ronstar 50WP
Escort XP	Sahara
Gallery 75DF	simazine
Krovar I DF	Surflan

Landmark 11 MP	Telar
Landmark MP	Vanquish
Landmark XP	2,4-D
Oust	

See the "MIXING" section of this label for general instructions for tank mixing.

Release of Bermudagrass or Bahiagrass

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

TANK MIXTURES: This product may also be tank-mixed with Outrider, Oust or Oust XP for residual control. These tank mixtures may delay greenup.

Apply 8 to 64 fluid ounces of this product in a tank mix with 0.75 to 1.3 ounces of Outrider per acre. Read and follow all label directions for Outrider.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust or Oust XP in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Oust or Oust XP per acre on Bermudagrass and no more than 0.5 ounce of Oust or Oust XP per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in 10 to 40 gallons of water per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestein, silver	Trumpet creeper
Fescue, tall	Vaseygrass

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 9 to 32 fluid ounces of this product with 0.75 to 1.3 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank-mixed with Oust or Oust XP. If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Oust or Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust or Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Bluestein, silver	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Trumpet creeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not done, since severe injury may occur.

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product *per* acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

TANK MIXTURES: This product may be tank-mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 6 fluid ounces of this product with 0.75 to 2 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weed greater than 6 inches in height. Use only on well-established bahiagrass.

A tank mixture of this product plus Oust or Oust XP may be used. Apply 6 fluid ounces of this product plus 0.25 ounce of Oust or Oust XP per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.10 Rangelands

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands, pastures, and industrial sites. Preventing weed seed production is critical to the successful control of annual grassy weeds invading these perennial grass sites. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Bromus: This product may be used to control or suppress downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*), cheatgrass (*Bromus secalinus*), cereal rye and jointed goatgrass found in rangelands pastures and industrial sites. Apply 8 to 16 fluid ounces of this product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead: To control or suppress medusahead, apply 16 fluid ounces of this product per acre at the 3-leaf stage when plants are actively growing. Delaying applications beyond this stage will result in reduced or unacceptable control. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses. Applications may be made in the fall or spring.

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

Spot Treatment, Wiper Application

This product may be applied in rangeland, pastures or industrial sites as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire site or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 3 quarts of this product per acre, no more than 10 percent of the total site may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

8.11 Utility Sites

In utilities, this product is for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product is also for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

For control of herbaceous weeds, use the lower tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher tank mixture rates.

Bare Ground, Trim-and-Edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products provided that the specific product used is registered for application on these sites. Refer to the individual product's labels for approved sites and application rates.

2,4-D	Outrider
Arsenal	pendimethalin
atrazine	Plateau
Baricade 65WG	Princep
dicamba	Ronstar 50WP
diuron	Sahara
Endurance	simazine
Escort	Surflan AS
Escort XP	Surflan WDG
Garlon 3A	Telar DF
Garlon 4	Transline
Krenite	Vanquish
Krovar I DF	Veipar DF
Oust	Velpar L
Oust XP	

Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray incompatibility problems. For side trimming treatments, this product be used alone or in a tank mixture with Garlon 4.

8.12 Conservation Reserve Program (CRP)

This product may be used for renovation (rotating out of CRP), site preparation, postemergence weed control in dormant CRP grasses, or wiper application on CRP land.

Renovation (Rotating out of CRP), Site Preparation

This product may be used to prepare CRP land for crop production. Refer to Federal, State or local use guides for CRP renovation recommendations.

Postemergence Weed Control in Dormant CRP Grasses, Wiper Application

Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land.

Applications may be made using wiper applicators to control tall weeds, or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 8 to 12 fluid ounces of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts per acre per year onto CRP land.

8.13 Grass Seed or Sod Production

This product may be used in grass seed and sod production for preplant, at-planting, preemergence, removal of established stands, renovation, site preparation, shielded spraying, wiper application, spot treatment, and creating rows in annual ryegrass.

Preplant, Preemergence, At-Planting, Removal of Established Stands, Renovation, Site Preparation

This product controls most existing vegetation for purposes of renovating turf or forage grass seed areas or for establishing turfgrass grown for sod. It may also be used to destroy remaining undesired grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting, or for renovation purposes. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates up to 5 quarts per acre may be used to totally remove established stands of tough to kill grass species.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Applications must be made prior to crop emergence in order to avoid crop injury.

Shielded Sprayers

Apply 1 to 3 quarts of this product in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aids in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by the protective shields.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction.

Wiper Application

This product may be applied over the top of desirable grasses using wiper applicators for the control of tall weeds.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction.

Spot Treatment

Apply a 1.5-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

The crop sprayed in the treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

Creating Rows in Annual Ryegrass

Apply 1 to 2 pints of this product per acre. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height. Use the higher rate when ryegrass is greater than 6 inches in height.

Set nozzle heights to allow the establishment of the desired row spacing. Use low-pressure nozzles, or drop

nozzles designed to target the application over a narrow band.

Take care not to spray or allow spray to drift outside target area in order to avoid unwanted crop destruction.

8.14 Pastures

This product may be applied to any pasture grass (*Gramineae* family), including bahiagrass, Bermudagrass, bluegrass, bromie, fescue, guineagrass, kikuyugrass, orchardgrass, pangola grass, ryegrass, timothy, and wheatgrass. Application can be made as a spot treatment, wiper application, preplant, preemergence, pasture renovation, or postemergent broadcast.

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be applied to control perennial pasture species listed on this label prior to replanting.

If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment, Wiper Application

This product may be applied in pastures as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 3 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper application are made using rates above 3 quarts of this product per acre, no more than 10 percent of the total pasture may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

Postemergent Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3 quarts of this product per acre per year onto pasture grasses except for renovation uses as described previously in this section.

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for rates to the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results.

9.1 Annual Weeds

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions. Use the higher rate for tough-to-control species regardless of the weed size at application. Treat tough-to-control weeds early when they are relatively small. This product may be tank mixed provided the tank-mix product is registered for use on the target site. Refer to the individual product labels for approved sites and application rates. For spray-to-wet applications, apply a 0.5-percent solution of this product to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds.

For low volume directed spray applications, use a 4- to 7-percent solution of this product. Spray coverage should be

uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple spouts.

WEED SPECIES	
Anoda, spurred	Lamb's-quarters*
Barley*	Little barley*
Barnyardgrass*	London rocket*
Bittercress*	Mayweed
Bassia, fivehook	Medusahcad*
Black nightshade*	Morningglory (<i>Ipomoea spp</i>)
Bluegrass, annual*	Mustard, blue*
Bluegrass, bulbous*	Mustard, tansy*
Brome, downy*	Mustard, tumble*
Brome, Japanese*	Mustard, wild*
Brown-top panicum*	Oats
Buttercup*	Pigweed*
Carolina foxtail*	Plains/Tickseed coreopsis*
Carolina geranium Castor bean	Prickly lettuce*
Cheatgrass*	Puncturevine
Cheeseweed (<i>Malva parviflora</i>)	Purslane, common
Chervi*	Ragweed, common*
Chickweed*	Ragweed, giant
Cocklebur*	Red rice
Copperleaf, hophornbeam	Russian thistle
Corn*	Rye*
Corn speedwell*	Ryegrass*
Crabgrass*	Sandbar, field*
Dwarf dandelion*	Shepherd's-purse*
Eastern managrass*	Sicklepod
Eclipta*	Signalgrass, broadleaf*
Fall panicum*	Smartweed, ladythumb*
Falsedandelion*	Smartweed, Pennsylvania*
Falsesflax, smallseed*	Sowthistle, annual
Fiddleneck	Sowthistle, annual
Field pennycress*	Spanishneedles
Filaree	Speedwell, purslane*
Fleabane, annual*	Sprangletop*
Fleabane, hairy (<i>Conyza bonariensis</i>)*	Spurge, annual
Fleabane, rough*	Spurge, prostrate*
Florida pulley	Spurge, spotted*
Foxtail*	Spurry, umbrella*
Goatgrass, jointed*	Starthistle, yellow
Goosegrass	Stinkgrass*
Grain sorghum (milo)*	Sunflower*
Groundsel, common*	Teaweed/ Prickly sida
Hemp sesbania	Texas panicum*
Henbit	Velvetleaf
Horseweed/Marestail (<i>Conyza canadensis</i>)	Virginia copperleaf
Itchgrass*	Virginia pepperweed*
Johnsongrass, seedling	Wheat*
Jungle rice	Wild oats*
Knotweed	Witchgrass*
Kochia	Woolly cupgrass*
	Yellow rocket

*When using field broadcast equipment (aerial applications or boom sprayers using flat fan nozzles) these species will be controlled or partially controlled using 1 pint of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure through coverage of foliage and treat when weeds are in an early growth stage.

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate.

Ensure thorough coverage when using spray-to-wet treatments with hand-held equipment. For best results, use a 1.5-percent solution on harder-to-control perennials such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

For low volume directed spray applications, use a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Allow 7 or more days after application before tillage.

WEED SPECIES	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2 - 4	1 - 2
Bahiagrass	3 - 5	2
Beachgrass, European (<i>Ammophila arena</i>)	-	5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4 - 5	2
Bluegrass, Kentucky	2	2
Blueweed, Texas	4 - 5	2
Brackenfern	3 - 4	1-1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf	-	2
Canarygrass, reed	2 - 3	2
Cattail	3 - 5	2
Clover: red, white	3 - 5	2
Cogongrass	3 - 5	2
Dallisgrass	3 - 5	2
Dandelion	3 - 5	
Dock, curly	3 - 5	2
Dogbane, hemp	4	2
Fescue (except tall)	3 - 5	2
Fescue, tall	1 - 3	2
German ivy	2 - 4	1 - 2
Guineagrass	3	1
Horsenettle	3 - 5	2
Horseradish	4	2
Iceplant	2	1.5-2
Jerusalem artichoke	3 - 5	2
Johnsongrass	2 - 3	1
Kikuyugrass	2 - 3	2
Knapweed	4	2
Lantana	-	1 - 1.25
Lespedeza	3 - 5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3 - 5	2
Napiergrass	3 - 5	2
Nightshade, silverleaf	2	2

Nutsedge: purple, yellow	3	1 - 2
Orchardgrass	2	2
Pampasgrass	3 - 5	1.5 - 2
Paragrass	3 - 5	2
Pepperweed, perennial	4	2
Phragmites*	3 - 5	1 - 2
Poison hemlock	2 - 4	1 - 2
Quackgrass	2 - 3	2
Reed, giant	4 - 5	2
Ryegrass, perennial	2 - 3	1
Smartweed, swamp	3 - 5	2
Spurge, leafy*	-	2
Sweet potato, wild*	-	2
Thistle, artichoke	2 - 3	1 - 2
Thistle, Canada	2 - 3	2
Timothy	2 - 3	2
Torpedograss*	4 - 5	2
Trumpetcreeper*	2 - 3	2
Vaseygrass	3 - 5	2
Velvetgrass	3 - 5	2
Wheatgrass, western	2 - 3	2

* Partially Controlled

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

For best results when using hand-held equipment, use a 1.5-percent solution on harder-to-control woody brush and trees. For low volume directed-spray applications, apply a 5- to 10-percent solution of this product. Spray coverage should be uniform with at least 50-percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple spouts.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

WEED SPECIES	Rate (QT/A)	Hand-Held % Solution
Alder	3-4	1-1.5
Ash*	2-5	1-2
Aspen, quaking	2-3	1-1.5
Bearclover (Bearmat)*	2-5	1-2
Beech*	2-5	1-2
Birch	2	1
Blackberry	3-4	1-1.5
Blackgum	2-5	1-2
Bracken	2-5	1-2
Broom; French, Scotch	2-5	1.5-2
Buckwheat, California*	2-4	1-2
Cascara	2-5	1-1.5
Catalaw*	-	1-1.5
Ceanothus*	2-5	1-2
Chamise*	2-5	1
Cherry; bitter, black, pin	2-3	1-1.5
Coyote brush	3-4	1.5-2
Deerweed	2-5	1
Dogwood*	2-5	1
Elderberry	2	1
Elm*	2-5	1-2
Eucalyptus	--	2
Gorse*	2-5	1-2
Hasardia*	2-4	1-2
Hawthorn	2-3	1-1.5
Hazel	2	1
Hickory*	2-5	1-2
Honeysuckle	3-4	1-1.5
Hornbeam, American*	2-5	1-2
Kudzu	4	2
Locust, black*	2-4	1-2
Madrone resprouts*	--	2
Manzanita*	2-5	1-2
Maple, red	2-4	1-1.5
Maple, sugar	--	1-1.5
Monkey flower*	2-4	1-2
Oak; black, white*	2-4	1-2
Oak, post	3-4	1-1.5
Oak; northern, pin	2-4	1-1.5

Oak, Scrub*	2 - 4	1 - 1.5
Oak; southern red	2 - 3	1 - 1.5
Peppertree, Brazilian (Florida holly)*	2 - 5	1 - 2
Persimmon*	2 - 5	1 - 2
Pine	2 - 5	1 - 2
Poison ivy	4 - 5	2
Poison oak	4 - 5	2
Poplar, yellow*	2 - 5	1 - 2
Redbud, eastern	2 - 5	1 - 2
Rose, multiflora	2	1
Russian olive*	2 - 5	1 - 2
Sage, black	2 - 4	1
Sage, white*	2 - 4	1 - 2
Sage brush, California	2 - 4	1
Salmonberry	2	1
Saltcedar*	2 - 5	1 - 2
Sassafras*	2 - 5	1 - 2
Sourwood*	2 - 5	1 - 2
Sumac; laurel, poison, smooth, sugarbur	2 - 4	1 - 2
Sweetgum	2 - 3	1 - 1.5
Swordfern*	2 - 5	1 - 2
Tallowtree, Chinese	-	1
Tan oak resprouts*	-	2
Thimbleberry	2	1
Tobacco, tree*	2 - 4	1 - 2
Toyon*	-	2
Trumpet creeper	2 - 3	1 - 1.5
Vine maple*	2 - 5	1 - 2
Virginia creeper	2 - 5	1 - 2
Waxmyrtle, southern*	2 - 5	1 - 2
Willow	3	1
Yerbasanta*	-	2

* Partially Controlled

10.0 LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. To the extent consistent with applicable law, this warranty is also subject to the conditions and limitations stated on this labeling.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this

Company.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

EPA Reg. No. 86004-U
Repar-Glypho, LLC.
Silver Springs, MD 20910

III. SUPPLEMENTAL LABELING

Table of Contents: Supplemental labeling

	Name
A	FOR USE IN DORMANT ALFALFA
B	FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER FIRST CUTTING
C	FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS
D	FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARESTAIL, <i>Conyza canadensis</i>) IN COTTON, CORN, AND SOYBEANS
E	HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROL IN ROUNDUP READY CANOLA
F	FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRENCO COUNTY
G	FOR AERIAL APPLICATION IN ARKANSAS ONLY
H	FOR CONTROLLING BARNYARDGRASS (ECHINOCHLOA CRUS-GALLI) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY
I	FOR WEED CONTROL APPLICATION IN SEED PRODUCTION OF ALFALFA WITH THE ROUNDUP READY® GENE
J	FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY
K	BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS
L	FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS
M	GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS
N	FOR GROUND AND AERIAL APPLICATION TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY
O	GLYPHOSATE 41% SL HERBICIDE FOR CONIFER RELEASE

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **GLYPHOSATE 41% SL** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **GLYPHOSATE 41% SL** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

GLYPHOSATE 41% SL
Herbicide

EPA Reg. No. S6004-

Keep out of the reach of children.

CAUTION / CUIDADO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any other manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIREABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for **GLYPHOSATE 41% SL** for essential product performance information.

[INSERT SPECIFIC DIRECTIONS FOR USE TEXT FROM THE FOLLOWING SECTIONS]

Read the "Limit of Warranty and Liability" in the label booklet for **GLYPHOSATE 41% SL** before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

REPAR-GLYPHO, LLC. COMPANY
SILVER SPRING, MD 20910

A. FOR USE IN DORMANT ALFALFA

DIRECTIONS FOR USE

This product will control or suppress many weeds, including quackgrass, downy brome, and cheatgrass in dormant alfalfa.

Apply 8 to 12 fluid ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with **GLYPHOSATE 41% SL**.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

B. FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER THE FIRST CUTTING

DIRECTIONS FOR USE

This product may be applied at 16 fluid ounces per acre to control the weeds listed below and most other winter annual grass and broadleaf weeds in established coastal Bermudagrass pastures.

Annual bluegrass	Oats
Cheat	Ryegrass, Italian
Crabgrass	Sandbur, field
Henbit	Sunflower
Johnsongrass, seedling	Wheat
Little barley	Wild mustard

TIMING OF APPLICATION

Applications prior to spring growth: Apply this product in

either late winter or early spring but before new coastal Bermudagrass growth begins in the spring. Applications to new growth can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 60 days after making this application before grazing or harvesting the treated area.

Application following the first cutting: Apply this product after the first Bermudagrass cutting when the Bermudagrass has not yet begun to regrow. Applications made after regrowth has begun can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 28 days after making this application before grazing or harvesting the treated area.

NOTE: ONLY ONE APPLICATION PER YEAR MAY BE MADE TO ANY ONE FIELD. A SPRING APPLICATION PRIOR TO GROWTH AND AN APPLICATION FOLLOWING THE FIRST CUTTING MAY NOT BE MADE ON THE FIELD DURING THE SAME YEAR.

C. FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS (WHEAT AND BARLEY)

DIRECTIONS FOR USE

Refer to the GLYPHOSATE 41% SL label for rates and weeds controlled.

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 2 to 3 gallons of water per acre.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray to drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets.

In aerial applications, do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Ensure uniform application. Use appropriate marking devices when applying herbicides by air.

Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR IS MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

D. FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED (MARESTAIL, *Conyza Canadensis*) IN CROP AND NON-CROP AREAS.

DIRECTIONS FOR USE

For ground applications, use 10 to 20 gallons of water per acre. For aerial applications, use 3 to 15 gallons of water per acre. **For tank-mix recommendations, read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental or fact sheets published separately for all herbicides used.**

Cotton:

Preplant:

For the control of horseweed, apply this product (32 fluid ounces per acre) in a tank-mix with Clarity® (8 fluid ounces per acre). This application must be made 21 to 35 days before planting and before horseweed reaches 6 inches in height. In order to avoid crop injury, a minimum interval of 21 days during which there is at least 1 inch cumulative rainfall must be observed between Clarity application and planting of cotton.

2,4-D may be included in the tank-mixture with this product. Refer to the 2,4-D product label for the time intervals that are required between application and planting and other geographic use restrictions.

Post-directed (Roundup Ready® Cotton and Roundup Ready Flex cotton® varieties only):

Management of early season weed competition and the development of a crop height differential between cotton and horseweed is often achieved by a combination of preplant burn-down and post emergence over-the-top and/or directed applications of this product. These measures enhance the development of a height differential that is necessary to successfully make post-directed treatments. In-crop post-directed applications of MSMA (2 pounds active ingredient per acre) tank-mixed with diuron (0.5 to 0.75 pounds active ingredient per acre) should be made when the temperature is 80°F or higher.

Soybeans:

Preplant:

It is strongly encouraged that horseweed be controlled prior to planting. Apply a tank mixture of this product (32 fluid ounces per acre) with 2,4-D (0.5 pounds a.i. per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

In-crop (Roundup Ready Soybean varieties only):

This treatment should be used as a salvage treatment only for a horseweed infestation that was not controlled prior to planting. At the time of treatment, horseweed should not exceed 6 inches in height. Apply a tank mixture of this product (32 fluid ounces per acre) with FirstRate® (0.3 ounces per acre). Application should be made between full emergence of the first trifoliate leaf and 50% flowering stage of soybeans.

Corn:

Preplant, At-Planting, Preemergence:

Apply a tank mixture of this product (32 fluid ounces per acre) plus 2,4-D (0.5 pounds i.e. or per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Atrazine (1 to 2 pounds active ingredients per acre) may be included in the tank-mixture to provide residual control. Refer to the atrazine product label for specific use instructions.

In-crop (Roundup Ready Corn hybrids only):

Apply a tank-mixture of this product (32 fluid ounces per acre) plus Clarity (8 to 16 ounces per acre) or 2,4-D (0.5 to 1.0 pounds of inactive ingredients per acre) between corn emergence and the 5-leaf stage of growth (approximately 8 inches tall).

Dicamba may be included in the tank-mixture with this product. Refer to the dicamba product label for the time intervals that are required between application and planting and other geographic use restrictions.

Tree and Vine Crops:

Orchards (Pome Fruit, Stone Fruit and Tree Nuts).

Apply two quarts of this product plus 2,4-D (18 fluid ounces per acre Dri-Clean® or 2 pints per acre Orchard Master® CA) before marestail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is suggested. Follow labeling for 2,4-D product with regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply.

Vine Crops (grapes only):

Apply 2 quarts of this product per acre plus 2,4-D (18 fluid ounces per acre Dri-Clean®) before marestail exceeds 6 inches in height. A carrier volume of 15 gallons per acre is suggested. Follow Dri-Clean® labeling in regard to directions for use, precautionary statements and applicable schedules. Further local restrictions may apply. Also, residual herbicides such as diuron may provide additional preemergence control.

General Non-crop Areas:

Apply a tank mixture of this product (1 quart per acre) before horseweed exceeds 6 inches in height. Control may be enhanced by making applications when horseweed is still in the rosette stage of growth. This product may be tank-mixed with the following products provided that the specific product is registered for use on the target site. Refer to these product labels for approved sites and application rates. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive precautionary statements for each product in the mixture.

- 2,4-D
- Crossbow™
- Dicamba
- Gallery™ 75 DF
- Krovar® 1 DF
- Landmark™ II MP
- Landmark™ MP
- Milestone™
- Overdrive™
- Tejar™ DF
- Vanquish™
- Velpar® DF

HIGHER-RATE POSTEMERGENCE USE DIRECTIONS FOR WEED CONTROLS IN ROUNDUP READY CANOLA

DIRECTIONS FOR USE

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop).

DO NOT USE THIS PRODUCT ON CANOLA WITH A ROUNDUP READY GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA.

Maximum Allowable Combined Application Quantities Per Season	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application for emergence to 6-leaf stage	2 quarts per acre

Preplant, At-planting Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting

Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Weeds Controlled: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" in this label

Single Application- Apply 16 to 32 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

Sequential Application-Apply 32 fluid ounces per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but not later than the 6-leaf stage. Sequential applications are used for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass or when controlling weeds with multiple application times.

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in the Roundup Ready crops. No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 64 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

F. FOR LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRESNO COUNTY, CALIFORNIA

DIRECTIONS FOR USE

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this supplemental labeling and in the product label booklet. Refer to Aerial Equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the product label for additional information. Refer to the individual use site section of the product label, or to other supplemental labeling or technical fact sheets published separately for this product by Repar-Glypho, LLC., for specific use instructions.

AVOID DRIFT--DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetations after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops:

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).

3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.
5. Apply by air only to nonresidential areas.

When applied under the conditions described, this product controls annual and perennial weeds as listed in the label booklet

When tank mixing product with 2,4-D, only 2,4-D amine formulations may be used for aerial applications in California. Tank mixtures 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and alfalfa and pasture renovation applications only. This product, when tank mixed with dicamba, may not be applied by air in California.

ADDITIONAL INFORMATION FOR FRESNO COUNTY, CALIFORNIA

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California.

North:	Fresno County line
South:	Fresno County line
East:	State Highway 99
West:	Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Recommendations

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity or surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rate of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to Fresno County Agricultural Commissioner.

Applications at Night- Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

Note: For aerial application from April 1 through February 14, refer to the other sections of this supplemental label.

G. FOR AERIAL APPLICATION IN ARKANSAS ONLY

DIRECTIONS FOR USE

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Do not apply this product when winds are in excess of 10 mph.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 mph is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 mph toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

H. FOR CONTROLLING BARNYARDGRASS (*ECHINOCHLOA CRUS-GALLI*) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY

DIRECTIONS FOR USE

Renovation Treatment

USE INSTRUCTIONS: This product may be applied as a renovation treatment in rice crops to control barnyardgrass infestations using ground broadcast spray or hand-held equipment. Renovation is defined as

herbicide treatment that will produce crop and weed destruction in an entire field or contiguous area treated within a field. Follow the application methods and treatment rates in the label booklet for **GLYPHOSATE 41% SL** herbicide.

PRECAUTIONS, RESTRICTIONS: Crop sprayed in treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction. The rice straw and stubble from the treated area, including a 25-foot buffer zone on all sides, shall not be used for grazing, animal bedding or any feed purposed.

Aerial applications are not permitted for rice renovation using this supplemental label.

I. FOR WEED CONTROL APPLICATIONS IN SEED PRODUCTION OF ROUNDUP READY® ALFALFA
--

General Information

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FURSTI OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for GLYPHOSATE 41% SL for essential product performance information.

The Roundup Ready designation indicates that the alfalfa contains a patented gene that provides tolerance to this product. Information on Roundup Ready Alfalfa varieties may be obtained from your seed supplier or Repar-Glypho, LLC, representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

See the "ROUNDUP READY CROPS" section of the GLYPHOSATE 41% SL label booklet for general precautionary instructions for use in Roundup Ready crops. Do NOT combine the instructions in this supplemental label with other recommendations listed in the "PSTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS" section of the **GLYPHOSATE 41% SL** label booklet intended for alfalfa varieties that do not contain a glyphosate tolerance gene.

DIRECTIONS FOR USE

This product will control many troublesome emerged weeds with over-the-top applications in Roundup Ready alfalfa grown for seed. Over-the-top applications may be made from emergence through the late vegetative stage, and spot treatments may be made from early bud stage through seed harvest.

For ground applications with broadcast equipment, apply this product in 3 to 40 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial application: Use rates of this product in 3 to 15 gallons of spray solution per acre.

DO NOT EXCEED 2 QUARTS OF THIS PRODUCT PER ACRE WHEN MAKING APPLICATIONS BY AIR. FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATION IN THAT STATE. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready alfalfa. Follow the cleaning procedures specified on the label of the product(s) used. Alfalfa can be very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence and Post-harvest of seed

Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Total Preplant, At-planting and Preemergence Applications	2 quarts per acre
Total in-crop application rate from emergence through the late vegetative stage*	6 quarts per acre
Spot-treatment during early bud stage through seed harvest (See the "Spot Treatment after late vegetative stage" section and the "PRECAUTIONS AND RESTRICTIONS" section of this label for complete instructions)	Apply spray-to-wet; do not apply to the point of runoff

There are no rotational crop restrictions following applications of this product. For any crop NOT listed in the label booklet, applications must be at least 30 days prior to planting.

Over-the-top applications: Broadcast applications of this product may be made using ground or aerial application equipment over the top of Roundup Ready alfalfa from emergence through the late vegetative stage. Do not make broadcast applications of this product between the initiation of alfalfa budding and the harvest of seed. Any single over-the-top broadcast application of this product should not exceed 2 quarts per acre. Sequential applications of this product should be at least 7 days apart.

Due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings are susceptible and will not survive or thrive after the first application of this product. A single application of at least 1 quart per acre of this product should be applied at or before the 3 to 4 trifoliate growth stage to eliminate the effects of stand gaps created by the loss of non-Roundup Ready plants.

Spot Treatment after late vegetative stage: For late emerging weeds, this product may be applied as a spot treatment in Roundup Ready alfalfa grown for seed, during the early bud stage through seed harvest. Applications made during this stage may result in reduced seed yield and quality and are the responsibility of the grower. Make applications on a spray-to-wet basis. Do not spray to the point of runoff. If a spot treatment is made after the late vegetative stage, harvested seed must not be used for alfalfa sprout production.

Post-harvest applications: Following harvest of Roundup Ready alfalfa seed, the stand may be managed for forage and hay production. Refer to the Roundup Ready alfalfa section of the label booklet GLYPHOSATE 41% SL for rates and timing of applications for forage and hay production.

Weeds controlled: For specific rates of application and instructions for control of various annual and perennial weeds, refer to the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" in the GLYPHOSATE 41% SL label booklet. Some weeds with multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some re-growth of weeds has occurred.

In addition to those weeds listed in the GLYPHOSATE 41% SL label booklet, this product will suppress or control the parasitic weed, Dodder (*Cuscuta spp.*) in Roundup Ready alfalfa seed production. Repeat applications may be necessary for complete control.

Tank mixtures with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control and are NOT for over-the-top applications of this product.

RESTRICTIONS: Do not make over-the-top broadcast applications of this product between the initiation of alfalfa budding and the harvest of Roundup Ready alfalfa seed. The use of harvested Roundup Ready alfalfa seed is not suitable for, and is not for, production of alfalfa sprouts.

Read the "LIMIT OF WARRANTY OR LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using. For (in-crop) over-the-top uses on Roundup Ready crop varieties, crop safety and weed control performance are not warranted by Repar-Glypho, LLC, when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

J. FOR APPLICATIONS TO ROUNDUP READY FLEX COTTON IN THE STATE OF ARIZONA ONLY

DIRECTIONS FOR USE

GENERAL INFORMATION

The use of the over-the-top applications described in this label on *other* than Roundup Ready Flex cotton will cause crop injury and reduced yields. Drift of this product from applications made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (nodes) Roundup Ready cotton may cause extensive injury including boll loss, delayed maturity and/or yield loss.

Note: The instructions provided in this supplemental label are specific to, and should only be used with varieties designated as **Round Ready Flex cotton**. **DO NOT** combine the instructions in this supplemental label with those in the "Roundup Ready Cotton" or "Roundup Ready Flex Cotton" sections found in the label booklet for **GLYPHOSATE 41% SL**, or with any other Roundup Ready cotton or Roundup Ready Flex cotton instruction on labeling for this or other glyphosate-containing product. See "Maximum Use Rate" in the "GENERAL INFORMATION" section of the label booklet for **GLYPHOSATE 41% SL** for additional information.

ROUNDUP READY FLEX COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATED THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

TYPES OF APPLICATIONS: Preplant, At-planting, Preemergence, Postemergence (In-crop), Preharvest

Maximum Allowable Combined Application Quantities Per Season	
--	--

Combined total per year for all application	8 quarts per acre
---	-------------------

Calculate the combined rate to be used for all preplant, in-crop and preharvest applications, to ensure that the total does not exceed the maximum allowed rate per acre year shown above.	
--	--

Preplant, At-planting, Preemergence applications	5 quarts per acre
--	-------------------

Total in-crop applications from ground cracking to 60% percent open bolls	6 quarts per acre
---	-------------------

Maximum allowed from 60 percent bolls open to 7 days prior to harvest	2 fluid ounces per acre
---	-------------------------

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of the **GLYPHOSATE 41% SL** label booklet for general precautionary instructions for use in Roundup Ready crops.

Preplant, At-planting, Preemergence

This product may be applied before, during or after planting Roundup Ready Flex cotton.

Postemergence (In-crop)

When applied in accordance with this label, GLYPHOSATE 41% SL will control labeled annual grasses and broadleaf weeds in Roundup Ready Flex cotton. To maximize yield potential, spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product. In general, an initial application of 1 quart per acre on 1 to 3 inch tall annual grass and broadleaf weeds. This product may be applied by ground application equipment at rates up to 2 quarts per acre per application postemergence to Roundup Ready Flex cotton. In addition to broadcast applications, post-directed equipment may be used to achieve weed coverage.

NOTE: For specific rates of application and instructions, refer to the "ANNUAL" and "PERENNIAL WEEDS RATE TABLES" in the label booklet for **GLYPHOSATE 41% SL**.

PRECAUTIONS, RESTRICTIONS: The maximum rate for any single in-crop application of this product is 2 quarts per acre made using ground application equipment. **In-crop application rates above 1 quart per acre made alone, or with the addition of other crop chemical products containing surfactant, may cause a crop response, including leaf speckling or leaf necrosis.** Do not exceed a maximum rate of 3 pints per acre of this product when making applications by air. Between layby and 60 percent open bolls, the maximum combined total of all applications made from crop emergence to 60 percent open bolls must not exceed 6 quarts per acre.

Preharvest

Apply this product for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready Flex cotton after 60 percent boll crack. Up to 2 quarts of this product may be applied using either aerial or ground spray equipment. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of Roundup Ready Flex cotton. Do not apply this product over-the-top beyond first bloom to cotton grown for seed.

PRECAUTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY FLEX COTTON, HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

K. BROADCAST APPLICATIONS IN CHRISTMAS TREE PLANTATIONS.

DIRECTIONS FOR USE

NOTE: IF IMPROPERLY APPLIED, THIS PRODUCT HAS THE POTENTIAL TO CAUSE SEVERE CHRISTMAS TREE INJURY. FOLLOW ALL LABEL DIRECTIONS.

This product may be applied as a broadcast spray over established Christmas trees. Ensure that adequate buffers are maintained to prevent drift onto nearby desirable crops or vegetation. Read the entire "APPLICATION"

“EQUIPMENT AND TECHNIQUES” section of the GLYPHOSATE 41% SL label booklet for additional application precautions

This application is approved for the following Christmas tree species:

Douglas fir (*Pseudotsuga menziesii*)

Fir species (*Abies* spp.)

Spruce species (*Picea* spp.)

Applications may be made only after trees have completed at least a full growing season since planting or transplanting. Applications should not be made within 1 full year prior to tree harvest.

Applications may only be made in the fall after the formation of final conifer resting buds. Final resting buds must be fully hardened and in the dormant stage. Applications made at any other time may result in unacceptable Christmas tree injury.

Avoid spray pattern overlap, as injury may occur.

Apply 1 quart of this product per acre in 5 to 30 galls of water per acre.

NOTE: DO NOT ADD SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO THIS PRODUCT AS SEVERE CHRISTMAS TREE INJURY MAY RESULT.

This product may be used at rates from 1 to 2 quarts per acre in some areas. Consult your local Repar-Glypho, LLC representative or Roundup supplier for specific guidance if you require rates greater than 1 quart per acre.

Drift control additives may increase Christmas tree injury.

The use of other herbicides tank mixed with this product is not to be used because severe Christmas tree injury may result.

L. FOR USE FOR SELECTIVE WEED CONTROL ON GLYPHOSATE 41% SL TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

GLYPHOSATE 41% SL Tolerant Tall Fescue Selections For Seed Protection

Use this product on [INSERT BRAND NAME] tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 4 to 16 fluid ounces per acre as a postemergence spray on GLYPHOSATE 41% SL tolerant tall fescue selections. See the label booklet for application instructions, rates, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the GLYPHOSATE 41% SL label booklet for a complete list of weeds controlled or suppressed.

NOTE: The rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with GLYPHOSATE 41% SL tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

Timing of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

M. GLYPHOSATE 41% SL HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the GLYPHOSATE 41% SL label booklet for rates. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quart per acre of this product may be used for improved results. Apply 10 quarts when weeds are dense.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any rates of this product may be used in a tank mixture with these products.

<u>Tank-mix Product</u>	<u>Rate per Acre</u>
Arsenal TM *	0.5 to 4 pints
Banvel	1 to 4 pints
2, 4-D	0.5 to 1 pound
Garlon TM 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar TM X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar TM I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust TM	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike TM 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

*Arsenal is not approved for use in the state of California

Refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mixture products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Read "LIMITS OF WARRANTY AND LIABILITY" in the label booklet for GLYPHOSATE 41% SL before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

To the extent consistent with applicable law, when used in combination, the liability of Repar-Glypho, LLC shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Repar-Glypho, LLC product in such combination use.

TMArsenal is a trademark of American Cyanamid Company.

TMBanvel is a trademark of Sandoz Crop Protection Corporation.

TMGarlon and Spike are trademarks of Dow AgroSciences LLC.

TMHyvar, Krovar and Oust are trademarks of E. I. DuPont de Nemours and Company.

N. FOR GROUND AND AERIAL APPLICATIONS TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY.
--

DIRECTIONS FOR USE

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Nonionic surfactants which are labeled for use with herbicides may be used to improve wetting of foliage. Do not reduce rates of GLYPHOSATE 41% SL when adding surfactant. Read and carefully observe surfactant rates, cautionary statements, and other information appearing on the surfactant label.

TIMING OF APPLICATION: Apply this product as a broadcast spray when plants are actively growing for partial control of undesirable vegetation listed on this label. Best results are obtained when application is made in the spring to early summer when brush species are at a high moisture content and flowering.

This product may be used for:

- Aid to burning treatment to establish and maintain fuel breaks
- Establishing fire perimeters and black lines
- Aid to prescribed burning
- Along fire roads and rights-of-way

APPLICATION RECOMMENDATION: Apply 2 quarts of this product per acre for partial control of the following emerged brush and chaparral species:

Ceanothus	Sage
Ceanothus spp.	Salvia spp.
Chamise	Scrub oak
Adenostroma fasciculatum	Quercus dumosa

Ground applications should be applied in 3 to 40 gallons of total spray solution per acre.

Aerial applications (helicopter only) should be applied in 3 to 15 gallons of total spray solution per acre.

Avoid direct application to any body of water.

AVOID DRIFT- DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

For aerial application of this product, please see the supplemental label directions for serial application in California.

O. GLYPHOSATE 41% SL HERBICIDE FOR CONIFER RELEASE

DIRECTIONS FOR USE

AERIAL APPLICATION

This product may be applied using aerial spray equipment for conifer release treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of the label booklet for GLYPHOSATE 41% SL herbicide for information on how to properly spray this product by air.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

CONIFER RELEASE

For release, apply only where conifers have been established for more than one year. Vegetation should not be disturbed prior to treatment or until symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of the label booklet for GLYPHOSATE 41% SL herbicide.

For release of the following conifer species:

Douglas Fir	Pines*
Pseudotsuga menziesii	Pinus spp.
Fir	Spruce
Abies spp.	Picea spp.
Hemlock	
Tsuga spp.	

*Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 1.5 to 2 quarts of this product per acre except in Washington and region, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of this product per acre before any major leaf drop of deciduous species.

For release of western hemlock apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly Pine

Pinus taeda

Eastern white pine

Pinus strobus

Slash Pine

Pinus elliotti

Late Season Application – Apply 1.5 to 2 quarts of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will result in potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later application. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label direction will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash

Fraxinus spp.

Cherry, black

Prunus serotina

Cherry, pin

Prunus pensylvanica

Elm

Ulmus spp.

Hawthorn

Crataegus spp.

Locust, black

Robinia pseudoacacia

Maple, red

Acer rubra

Oak, black

Quercus velutina

Oak, post

Quercus stellata

Oak, southern red

Quercus falcata

Oak, white

Quercus alba

Persimmon

Diospyros spp.

Poplar, yellow

Liriodendron tulipifera

Sassafras

Sassafras albidum

Sourwood

Oxydendrum arboreum

Sumac, poison

Rhus vernix

Sumac, smooth

Rhus glabra

Sumac, winged

Rhus copallina

Sweetgum

Liquidambar styraciflua

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

GLYPHOSATE 41% SL HERBICIDE PLUS OUST™ TANK MIXTURES FOR CONIFER RELEASE FROM HERBACEOUS WEEDS

To release loblolly pines from herbaceous weeds, tank mixtures of this product with Oust will provide control of annual weeds listed in the "WEEDS CONTROLLED" section of the label booklet for GLYPHOSATE 41% SL herbicide and the Oust label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of GLYPHOSATE 41% SL herbicide with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines.

This product plus Oust tank mixtures may not be applied by air in California.

This tank mixture may be applied using aerial equipment. When applying by air, use the selected rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use low rates of both products. Use higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	Johnsongrass**
<i>Paspalum notatum</i>	<i>Sorghum halepense</i>
Broomsedge	Poorjoe**
<i>Andropogon virginicus</i>	<i>Diodia teres</i>
Dock, curly	Trumpet creeper*
<i>Rumex crispus</i>	<i>Campsis radicans</i>
Dogfennel	Vaseygrass
<i>Eupatorium capilliflorum</i>	<i>Paspalum urvillei</i>
Fescue, tall	Vervain, blue
<i>Festuca arundinacea</i>	<i>Verbena hastata</i>

*Suppression at the higher rates only

**Control at the higher rates

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water, insects or disease.

Read and observe the cautionary statements and all other information appearing on the labels of herbicides used.

Oust is a trademark of E.I. DuPont de Nemours and Company.

MANDAVA ASSOCIATES, LLC

CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS
1050 CONNECTICUT AVENUE, N.W., SUITE 1000, WASHINGTON, DC 20036
Telephone: (202)-223-1424/1747 · Fax: (202)-223-0141 · E-MAIL: Mandava@compuserve.com

Hand Delivered

February 17, 2009

Mr. James Tompkins
Product Manager (PM 25)
Herbicide Branch
Registration Division (7504C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Washington, D.C. 20460

SUBJECT: ME-TOO Application for Registration
Product Name: **GLYPHOSATE 41% SL**
Active Ingredient: Glyphosate
Company Name: Repar-Glypho, LLC
EPA Company Number: 86004

Dear Mr. Tompkins:

On behalf of Repar-Glypho, LLC, we are submitting a **ME-TOO** application for the registration of **Glyphosate 41% SL** which is an end-use product containing Glyphosate as an active ingredient (AI). The source for AI is Glyphosate Technical (EPA Reg. No: To Be Assigned).

In support of the application for the subject product registration, we are providing the following administrative materials and product chemistry data on **GLYPHOSATE 41% SL**:

- 1) Application for Pesticide Registration
- 2) Confidential Statement of Formula
- 3) Certification with Respect to Citation of Data
- 4) Data Matrix
- 5) Five Copies of Draft Labeling for the Subject Product

Mr. James Tompkins
February 17, 2009

Page 2

- 6) Volume 5: **GLYPHOSATE 41% SL**: Product Properties Data in Support of FIFRA Registration; Data Requirement: 40 CFR 158.300-355; OPPTS Series 830 (Project No.: REPAR-GLYPHO41SL-4); Repar-Glypho, LLC.

S

Please note that the physical and chemical properties data presented in **Volumes 4.1 to 4.5 for GLYCEL 41% SL** will also support the **GLYPHOSATE 41% SL** because both products have the same composition and are identical in all respects.

The product chemistry data for the subject product was developed by Repar-Glypho, LLC.


Please also note that Repar-Glypho, LLC is relying on the safety data submitted by other registrants on Glyphosate.

In addition to the above studies, we are also relying on other items of data on the subject product in the Agency's files.

We are requesting an "**Expedited Review**" in accordance with FIFRA Section 3(c)(3)(b)(i), because our product (Glyphosate 41% SL) is substantially similar or identical in composition and labeling to Roundup Original Herbicide (EPA Reg. No.: 524-445). Based on the Signal Word approved for Gly Star Original (EPA Reg. No.: 42750-60) which is also a ME-Too product, we have assigned **CAUTION** for the subject product.

Since we have fulfilled all the registration requirements, we request you grant the registration for the subject product, so that the applicant can market it for the upcoming growing season.

Sincerely,



N. Bhushan Mandava, Ph.D., RAC
Agent for Repar-Glypho, LLC

Enclosure

TRANSMITTAL DOCUMENT

NAME AND ADDRESS OF SUBMITTER:

Repar-Glypho, LLC
c/o Mandava Associates, LLC
1050 Connecticut Avenue, NW / Suite 1000
Washington, DC 20036

REGULATORY ACTION SUPPORTED BY THIS PACKAGE:

FIFRA Registration Application for:

Product Name:	Glyphosate 41% SL
Active Ingredient:	Glyphosate
EPA Reg. No.:	(Not assigned)


TRANSMITTAL DATE:

February 10, 2009

LIST OF SUBMITTED STUDIES:

47685101 Volume 5: **GLYPHOSATE 41% SL**: Product Properties Data in Support of
FIFRA Registration; Data Requirement: 40 CFR 158.300-355;
OPPTS Series 830 (Project No.: REPAR-GLYPHO41SL-~~4~~); Repar-
Glypho, LLC. 5

COMPANY OFFICIAL:



N. Bhushan Mandava
Registration Agent for Repar-Glypho, LLC

COMPANY CONTACT:

N. Bhushan Mandava
Mandava Associates, LLC
1050 Connecticut Avenue, NW/Suite 1000
Washington, D.C. 20036
(202) 223-1424/1747



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

March 4, 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

OPP Decision Number: D-406734
EPA File Symbol or Registration Number: 86004-U
Product Name: GLYPHOSATE 41% SL
EPA Receipt Date: 23-Feb-2009
EPA Company Number: 86004
Company Name: REPAR-GLYPHO, LLC

N. BHUSHAN MANDAVA, PH.D.
MANDAVA ASSOCIATES, LLC
REPAR-GLYPHO, LLC
1050 CONN. AVE., NW, STE 1000
WASHINGTON, DC 20036-

SUBJECT: Receipt of Registration Application Subject to Registration Service Fee

Dear Registrant:

The Office of Pesticide Programs has received your application and certification of payment. If you submitted data with this application, the results of the PRN-86-5 screen will be communicated separately. During the administrative screen, the Office of Pesticide Programs has determined that this Action is subject to a Pesticide Registration Service Fee as defined in the Pesticide Registration Improvement Act.

The Action has been identified as Action Code: R310.1

NEW PRODUCT;NON-FAST TRACK (INCLUDES REVIEWS OF PRODUCT
CHEMISTRY;ACUTE TOXICITY;PUBLIC HEALTH PEST EFFICACY);Reduced Fee:
Linked to PRIA Application;

No additional payment is due at this time.

If you have any questions, please contact the Pesticide Registration Service Fee
Ombudsman at (703) 305-6249.

Sincerely,

A handwritten signature in cursive script, appearing to read "Teresa Torres", is written over the typed name.

Front End Processing Staff
Information Technology & Resources Management Division



United States
Environmental Protection Agency
Washington, DC 20460

☒

Registration
Amendment
Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 86004- U	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Glyphosate 41% SL	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Repar-Glypho, LLC 8070 Georgia Avenue, Suite 209 Silver Spring, MD. 20910 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 524-445 Product Name Roundup Original Herbicide	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input checked="" type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

- 1) Application for FIFRA Section 3 Registration. Submission of Product Chemistry Studies in support of registration of Glyphosate 41% SL.
- 2) The action covered is New Product, Me-Too Product, Fast Track since the application is for formulation product registration that is substantially similar and/or identical to a currently registered product as cited in this application. The company Number is 86004. The file symbol is not assigned.
- 3) The applicant requested fee waiver.
- 4) Please direct all correspondence to: Mandava Associates, LLC; 1050 Connecticut Avenue, NW / Suite 1000; Washington, D.C. 20036
- 5) The contact Person fax number is (202) 223 - 0141 and e-mail address is: mandava@compuserve.com.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 20 Kg, 50 Kg		5. Location of Label Directions <input checked="" type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name N. Bhushan Mandava		Title Agent for Repar-Glypho, LLC		Telephone No. (Include Area Code) (202) 223-1424	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, or both under applicable law.					6. Date Application Received (Stamped)
2. Signature N. Bhushan Mandava		3. Title Agent for Repar-Glypho, LLC			
4. Typed Name N. Bhushan Mandava		5. Date 02/17/09			

MANDAVA ASSOCIATES, LLC

CONSULTANTS IN SCIENCE, TECHNOLOGY AND REGULATORY AFFAIRS

1050 CONNECTICUT AVENUE, N.W., SUITE 1000, WASHINGTON, DC 20036

Telephone: (202)-223-1424/1747 · Fax: (202)-223-0141 · E-MAIL: Mandava@compuserve.com

Hand Delivered

February 17, 2009

Mr. James Tompkins
Product Manager (PM 25)
Herbicide Branch
Registration Division (7504C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Washington, D.C. 20460

SUBJECT: ME-TOO Application for Registration
Product Name: **GLYPHOSATE 41% SL**
Active Ingredient: Glyphosate
Company Name: Repar-Glypho, LLC
EPA Company Number: 86004

Dear Mr. Tompkins:

On behalf of Repar-Glypho, LLC, we are submitting a **ME-TOO** application for the registration of **Glyphosate 41% SL** which is an end-use product containing Glyphosate as an active ingredient (AI). The source for AI is Glyphosate Technical (EPA Reg. No: To Be Assigned).

In support of the application for the subject product registration, we are providing the following administrative materials and product chemistry data on **GLYPHOSATE 41% SL**:

- 1) Application for Pesticide Registration
- 2) Confidential Statement of Formula
- 3) Certification with Respect to Citation of Data
- 4) Data Matrix
- 5) Five Copies of Draft Labeling for the Subject Product

- 6) Volume 5: **GLYPHOSATE 41% SL**: Product Properties Data in Support of FIFRA Registration; Data Requirement: 40 CFR 158.300-355; OPPTS Series 830 (Project No.: REPAR-GLYPHO41SL-4); Repar-Glypho, LLC.

Please note that the physical and chemical properties data presented in **Volumes 4.1 to 4.5 for GLYCEL 41% SL** will also support the **GLYPHOSATE 41% SL** because both products have the same composition and are identical in all respects.

The product chemistry data for the subject product was developed by Repar-Glypho, LLC.


Please also note that Repar-Glypho, LLC is relying on the safety data submitted by other registrants on Glyphosate.

In addition to the above studies, we are also relying on other items of data on the subject product in the Agency's files.

We are requesting an "**Expedited Review**" in accordance with FIFRA Section 3(c)(3)(b)(i), because our product (Glyphosate 41% SL) is substantially similar or identical in composition and labeling to Roundup Original Herbicide (EPA Reg. No.: 524-445). Based on the Signal Word approved for Gly Star Original (EPA Reg. No.: 42750-60) which is also a ME-Too product, we have assigned **CAUTION** for the subject product.

Since we have fulfilled all the registration requirements, we request you grant the registration for the subject product, so that the applicant can market it for the upcoming growing season.

Sincerely,



N. Bhushan Mandava, Ph.D., RAC
Agent for Repar-Glypho, LLC

Enclosure

TRANSMITTAL DOCUMENT

NAME AND ADDRESS OF SUBMITTER:

Repar-Glypho, LLC
c/o Mandava Associates, LLC
1050 Connecticut Avenue, NW / Suite 1000
Washington, DC 20036

REGULATORY ACTION SUPPORTED BY THIS PACKAGE:

FIFRA Registration Application for:

Product Name:	Glyphosate 41% SL
Active Ingredient:	Glyphosate
EPA Reg. No.:	(Not assigned)


TRANSMITTAL DATE:

February 10, 2009

LIST OF SUBMITTED STUDIES:

Volume 5: **GLYPHOSATE 41% SL**: Product Properties Data in Support of FIFRA Registration; Data Requirement: 40 CFR 158.300-355; OPPTS Series 830 (Project No.: REPAR-GLYPHO41SL-4); Repar-Glypho, LLC.

COMPANY OFFICIAL:



N. Bhushan Mandava
Registration Agent for Repar-Glypho, LLC

COMPANY CONTACT:

N. Bhushan Mandava
Mandava Associates, LLC
1050 Connecticut Avenue, NW/Suite 1000
Washington, D.C. 20036
(202) 223-1424/1747



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 1.25 hours per response for registration and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the completed form to this address.

Certification with Respect to Citation of Data

Applicant's/Registrant's Name, Address, and Telephone Number Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910; (202) 223-1424	EPA Registration Number/File Symbol 86004-U
Active Ingredient(s) and/or representative test compound(s) Glyphosate (CAS # 1071-83-6)	Date March 5, 2009
General Use Pattern(s) (list all those claimed for this product using 40 CFR Part 158) Herbicide, Terrestrial, Food Crops	Product Name Glyphosate 41% SL

NOTE: If your product is a 100% repackaging of another purchased EPA-registered product labeled for all the same uses on your label, you do not need to submit this form. You must submit the Formulator's Exemption Statement (EPA Form 8570-27).

☐ I am responding to a Data-Call-In Notice, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).

SECTION I: METHOD OF DATA SUPPORT (Check one method only)

☐ I am using the cite-all method of support, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).

☒ I am using the selective method of support (or cite-all option under the selective method), and have included with this form a completed list of data requirements (the Data Matrix form must be used).

SECTION II: GENERAL OFFER TO PAY

[Required if using the cite-all method or when using the cite-all option under the selective method to satisfy one or more data requirements]

☒ I hereby offer and agree to pay compensation, to other persons, with regard to the approval of this application, to the extent required by FIFRA.

SECTION III: CERTIFICATION

I certify that this application for registration, this form for reregistration, or this Data-Call-In response is supported by all data submitted or cited in the application for registration, the form for reregistration, or the Data-Call-In response. In addition, if the cite-all option or cite-all option under the selective method is indicated in Section I, this application is supported by all data in the Agency's files that (1) concern the properties or effects of this product or an identical or substantially similar product, or one or more of the ingredients in this product; and (2) is a type of data that would be required to be submitted under the data requirements in effect on the date of approval of this application if the application sought the initial registration of a product of identical or similar composition and uses.

I certify that for each exclusive use study cited in support of this registration or reregistration, that I am the original data submitter or that I have obtained the written permission of the original data submitter to cite that study.

I certify that for each study cited in support of this registration or reregistration that is not an exclusive use study, either: (a) I am the original data submitter; (b) I have obtained the permission of the original data submitter to use the study in support of this application; (c) all periods of eligibility for compensation have expired for the study; (d) the study is in the public literature; or (e) I have notified in writing the company that submitted the study and have offered (i) to pay compensation to the extent required by sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA; and (ii) to commence negotiations to determine the amount and terms of compensation, if any, to be paid for the use of the study.

I certify that in all instances where an offer of compensation is required, copies of all offers to pay compensation and evidence of their delivery in accordance with sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA are available and will be submitted to the Agency upon request. Should I fail to produce such evidence to the Agency upon request, I understand that the Agency may initiate action to deny, cancel or suspend the registration of my product in conformity with FIFRA.

I certify that the statements I have made on this form and all attachments to it are true, accurate, and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

Signature <i>N. Bhushan Mandava</i>	Date March 5, 2009	Typed or Printed Name and Title N. Bhushan Mandava, Agent for Repar-Glypho, LLC
--	-----------------------	--



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 1 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.1550	Product Identity and Composition		Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910	OWN	
830.1600	Description of Materials Used to Produce the Product		Repar-Glypho, LLC	OWN	
830.1620	Description of Production Process		Repar-Glypho, LLC	OWN	
830.1650	Description of Formulation Process		Repar-Glypho, LLC	OWN	
830.1670	Discussion of Formation of Impurities		Repar-Glypho, LLC	OWN	
830.1700	Preliminary Analysis		Repar-Glypho, LLC	OWN	
830.1750	Certified Limits		Repar-Glypho, LLC	OWN	
830.1800	Enforcement Analytical Method		Repar-Glypho, LLC	OWN	
830.6302	Color		Repar-Glypho, LLC	OWN	
830.6303	Physical State		Repar-Glypho, LLC	OWN	
830.6304	Odor		Repar-Glypho, LLC	OWN	
830.6313	Stability to Normal & Elevated Temperatures, Metals and Metal Ions		Repar-Glypho, LLC	OWN	
830.6314	Oxidation/Reduction: Chemical Incompatibility		Repar-Glypho, LLC	OWN	
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 2 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.6315	Flammability		Repar-Glypho, LLC		Not Required
830.6316	Explosibility		Repar-Glypho, LLC		Not Required
830.6317	Storage Stability		Repar-Glypho, LLC	OWN	
830.6319	Miscibility		Repar-Glypho, LLC	OWN	
830.6320	Corrosion Characteristics		Repar-Glypho, LLC	OWN	
830.6321	Dielectric Breakdown Voltage		Repar-Glypho, LLC	OWN	
830.7000	pH		Repar-Glypho, LLC	OWN	
830.7050	UV/Visible Absorption		Repar-Glypho, LLC	OWN	
830.7100	Viscosity		Repar-Glypho, LLC	OWN	
830.7200	Melting Point/Melting Range		Repar-Glypho, LLC	OWN	
830.7220	Boiling Point/Boiling Range		Repar-Glypho, LLC	OWN	
830.7300	Density/Relative Density/Bulk Density		Repar-Glypho, LLC	OWN	
830.7370	Dissociation Constants in Water		Repar-Glypho, LLC	OWN	
830.7550	Partition Coefficient (n-octanol/water), Shake Flask Method		Repar-Glypho, LLC	OWN	
Signature <i>N. B. Mandava</i>		Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009	



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 3 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
830.7560	Partition Coefficient (n-octanol/water), Generator Column Method		Repar-Glypho, LLC	OWN	
830.7570	Partition Coefficient (n-octanol/water), Estimation by Liquid Chromatography		Repar-Glypho, LLC	OWN	
830.7840	Water Solubility: Column Elution Method; Shake Flask Method		Repar-Glypho, LLC	OWN	
830.7860	Water Solubility, Generator Column Method		Repar-Glypho, LLC	OWN	
830.7950	Vapor Pressure		Repar-Glypho, LLC	OWN	
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 4 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Syngenta Crop Protection, Inc.	Pay	
			Attn: Regulatory Affairs		
			P.O. BOX 18300		
			Greensboro, NC 27419 (Company # 100)		
			Nufarm Americas, Inc.	Pay	
			150 Harvester Drive, Suite 200		
			Burr Ridge, IL 60527 (Company # 228)		
			The Scotts Company	Pay	
			D/B/A The Ortho Group		
			P.O. BOX 190		
			Marysville, OH 43040 (Company # 239)		
			FMC Corp, Agricultural Products Group	Pay	
			1735 Market Street		
			Philadelphia, PA 19103 (Company # 279)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 5 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			E.I. DU PONT DE NEMOURS AND CO., INC	Pay	
			Dupont Crop Protection (S300/427)		
			P.O. BOX 30		
			Newark, DE 19714 (Company # 352)		
			Monsanto Company	Pay	
			1300 I Street, NW, Suite 450 East		
			Washington, DC 20005 (Company # 524)		
			Cheminova A/S	Pay	
			1600 Wilson Blvd., Suite 700		
			Arlington, VA 22209 (Company # 4787)		
			Helena Chemical Co.	Pay	
			225 Schilling Boulevard, Suite 300		
			Collierville, TN 38017 (Company # 5905)		
			Ritter Chemical, Inc. 1150 18th St., N.W., Suite	Pay	
			1000 Washington, DC 20036 (Company # 9468)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 6 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Drexel Chemical Company	Pay	
			PO BOX 13327		
			Memphis, TN 38113 (Company # 19713)		
			Industria Prodotti Chimici S.P.A.	Pay	
			122 C St., NW, Ste 740		
			Washington, DC 20001 (Company # 33660)		
			Nufarm Limited	Pay	
			P.O. BOX 13439		
			RTP, NC 27709 (Company # 35935)		
			Albaugh, Inc.	Pay	
			P.O. BOX 2127		
			Valdosta, GA 31604 (Company # 42750)		
			BASF Sparks, LLC	Pay	
			P.O. BOX 13528		
			RTP, NC 27709 (Company # 51036)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009			EPA Reg No./File Symbol 86004-		Page 7 of 11
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910			Product Glyphosate 41% SL		
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Valent U.S.A. Corporation	Pay	
			1600 Riviera Avenue, Suite 200		
			Walnut Creek, CA 94596 (Company # 59639)		
			Dow AgroSciences, LLC	Pay	
			9330 Zionsville RD 308/2E		
			Indianapolis, IN 46268 (Company # 62719)		
			Makhteshim-Agan of North America, Inc.	Pay	
			4515 Falls of Neuse RD, Suite 300		
			Raleigh, NC 27609 (Company # 66222)		
			Spray Drift Task Force	Pay	
			McKenna, Long & Aldridge, LLP		
			1900 K Street, NW		
			Washington, DC 20006 (Company # 66607)		
			Cheminova, Inc. 1600 Wilson Boulevard, Suite	Pay	
			700 Arlington, VA 22209 (Company # 67760)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009			EPA Reg No./File Symbol 86004-		Page 8 of 11
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910			Product Glyphosate 41% SL		
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Chemical Products Technologies, LLC	Pay	
			PO BOX 2275		
			Cartersville, GA 30120 (Company # 70829)		
			Nufarm, Inc.	Pay	
			150 Harvester Drive Suite 200		
			Burr Ridge, IL 60527 (Company #71368)		
			Outdoor Residential Exposure Task Force	Pay	
			1350 I Street, N.W.		
			Washington, DC 20005 (Company # 71754)		
			Agricultural Reentry Task Force	Pay	
			1350 I Street, N.W.		
			Washington, DC 20005 (Company # 71755)		
			Nations AG II, LLC	Pay	
			4515 Falls of Neuse Road, Suite 300		
			Raleigh, NC 27609 (Company # 72167)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 9 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Zhejiang Xinan Chemical Industrial Group	Pay	
			115 Obtuse Hill		
			Brookefield, CT 06804 (Company # 73432)		
			FIFRA Endangered Species Task Force, LLC	Pay	
			c/o Harold Himmelman (Beveridge & Diamond		
			PC) 1350 I Street, NW		
			Washington, DC 20005 (Company # 73989)		
			Helm Agro US, Inc.	Pay	
			8295 Tournament Drive, Suite 310		
			Memphis, TN 38125 (Company # 74530)		
			Residential Exposure Joint Venture (REJF)	Pay	
			900 17th Street, NW, Suite 300		
			Washington, DC 20006 (Company # 74888)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb .17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reviewing the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPF Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 10 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Agricultural Handlers Exposure Task Force	Pay	
			PO BOX 509		
			Macon, MO 63552 (Company # 75234)		
			Mey Corporation	Pay	
			121 S. Estes Drive, Suite 101		
			Chapel Hill, NC 27514 (Company # 80967)		
			Libertas Now, Inc.	Pay	
			108 Severn Ave., Suite B		
			Annapolis, MD 21403 (Company # 81142)		
			AXSS USA, LLC	Pay	
			115 Obtuse Hill Road		
			Brookfield, CT 06804 (Company # 83520)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date February 17, 2009		EPA Reg No./File Symbol 86004-		Page 11 of 11	
Applicant's/Registrant's Name & Address Repar-Glypho, LLC, 8070 Georgia Avenue, Suite 209, Silver Spring, MD 20910		Product Glyphosate 41% SL			
Ingredient Glyphosate (CAS # 1071-83-6)					
Guideline Reference Number	Guideline Study Name	MRID Number	Submitter	Status	Note
CITE-ALL			Celsius Property B.V., Amsterdam (NL)	Pay	
			Neuhausen A. RHF Branch		
			4515 Falls of Neuse Rd., Suite 300		
			Raleigh, NC 27609 (Company # 83558)		
			Generic Endangered Species Task Force	Pay	
			(GESTF) 500 Eighth Street, N.W.		
			Washington, DC 20004 (Company # 84653)		
			Red Eagle Chemicals Corp	Pay	
			PO Box 640		
			Hockessin, DE 19707 (Company # 85595)		
			Redeagle International LLC	Pay	
			PO BOX 640		
			Hockessin, DE 19707 (Company # 85678)		
Signature <i>N. B. Mandava</i>			Name and Title N. Bhushan Mandava, Ph.D., Agent for Repar-Glypho, LLC		Date Feb. 17, 2009



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☒ Amendment
☐ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 86004-U	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Glyphosate 41% SL	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Repar-Glypho, LLC 8070 Georgia Avenue, Suite 209 Silver Spring, MD. 20910 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 524-445 Product Name Roundup Original Herbicide	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

- 1) Submission of revised draft label with corrections in support of registration for Glyphosate 41% SL (EPA Reg. No.: 86004-U)
2) Please direct all correspondence to: Mandava Associates, LLC; 1050 Connecticut Avenue, NW / Suite 1000; Washington, D.C. 20036
3) The contact Person fax number is (202) 223 - 0141 and e-mail address is: mandava@compuserve.com.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Nat Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 20 Kg, 50 Kg		5. Location of Label Directions <input checked="" type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name N. Bhushan Mandava		Title Agent for Repar-Glypho, LLC		Telephone No. (Include Area Code) (202) 223-424	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature N. Bhushan Mandava		3. Title Agent for Repar-Glypho, LLC			
4. Typed Name N. Bhushan Mandava		5. Date 09/14/09			

Received
NTS/Anderson
8/24/09



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

**TECHNICAL REVIEW BRANCH
SIMILARITY CLINIC DETERMINATION**

21/AUG/2009

MEMORANDUM

Subject: Name of Pesticide Product: Glyphosate 41% SL
EPA Reg. No. /File Symbol: 86004-U
DP Barcode: D363699
Decision No: 406734
Action Code: R310.1
PC Code: 103601 (glyphosate IPA salt)

From: Eugenia McAndrew, Biologist *E. McAndrew* *M. Harshbarger*
Technical Review Branch
Registration Division (7505P)

To: Eugene Wilson/James Tompkins, RM Team 25
Herbicide Branch
Registration Division (7505P)

Applicant: Repar-Glypho, LLC
8070 Georgia Avenue, Suite 209
Silver Spring, MD 20910

FORMULATION FROM LABEL:

<u>Active Ingredient(s):</u>	<u>% by wt.</u>
Glyphosate, in the form of its isopropylamine salt	41.0
<u>Inert Ingredient(s):</u>	<u>59.0</u>
Total:	100.0%

ACTION REQUESTED: The Risk Manager requests: Please review enclosed information to determine if proposed product is similar enough to 524-445 for cited and submitted data to support registration.

BACKGROUND: Repar-Glypho, LLC has applied for registration of Glyphosate 41% SL, EPA File Symbol 86004-U, an end-use product with 41.0% glyphosate as the active ingredient. In a letter dated February 17, 2009, the Agent for the Registrant states:

...our product (Glyphosate 41% SL) is substantially similar or identical in composition and labeling to Roundup Original Herbicide (EPA Reg. No. 524-445). Based on the signal word approved for Gly Star Original (EPA Reg. No. 42750-60) which is also a ME-Too product, we have assigned "CAUTION" for the subject product.

The signal word for 524-445 is Warning and the signal word for 42750-60 is Caution. The Registrant may not claim similarity to one product and use the signal word from a second product. After examining the CSFs for the proposed product, 86004-U, and the two cited products, 524-445 and 42750-60, TRB believes that 42750-60 is the more appropriate product to cite for a similarity determination.

The Registrant is using the cite-all method of data support to satisfy the acute toxicity data requirements. A search of the OPP electronic databases shows that acute toxicity data were submitted to support the registration of the cited product, 42750-60. These studies were reviewed and classified as acceptable in two prior TRB memos (McAndrew; D266112; 07/JUN/2000 and McAndrew; D266765; 19/JUL/2000).

RECOMMENDATIONS: TRB has evaluated the formulations of the proposed product, 86004-U, and the cited product, 42750-60, and has determined that the two products are substantially similar.

The acute toxicity profile for the proposed product, Glyphosate 41% SL File Symbol 86004-U, is as follows:

acute oral toxicity	IV	cited	EPA Reg. No. 42750-60
acute dermal toxicity	IV	cited	EPA Reg. No. 42750-60
acute inhalation toxicity	IV	cited	EPA Reg. No. 42750-60
primary eye irritation	IV	cited	EPA Reg. No. 42750-60
primary skin irritation	IV	cited	EPA Reg. No. 42750-60
dermal sensitization	negative	cited	EPA Reg. No. 42750-60

The proposed basic CSF dated February 9, 2009 should be approved by the TRB Product Chemistry Team.

This memorandum pertains only to the decision concerning whether the subject product is similar to the cited product from an acute toxicological view point. For the purposes of this action, TRB has made no further determination of the adequacy of the toxicological data base or the precautionary label of the cited product.

LABELING: Based on the toxicity profile above, the following are the precautionary and first aid statements for this product as obtained from the Label Review System:

PRODUCT ID #: 086004-00004

PRODUCT NAME: Glyphosate 41% SL

PRECAUTIONARY STATEMENTS

SIGNAL WORD: CAUTION (optional)

Hazards to Humans and Domestic Animals:

Wear: Long-sleeved shirt and long pants, socks and shoes.

First Aid: No statements are required. Category III statements may be used.

Received
N.B. Kordan
8/19/09

DATE OUT: 13 Aug 2009

SUBJECT: PRODUCT CHEMISTRY REVIEW MP [] EP [x]
DP BARCODE No.: D363697
Reg. File Symbol No.: 86004-U
PRODUCT NAME: Glyphosate 41% SL
COMPANY: Repar-Glypho, LLC
Decision No.: 406734 PC CODE: 417300
FOOD USE: [x] Integrated Formulation []

FROM: Bruce F. Kitchens, Chemist
Technical Review Branch
Registration Division (7505P)

Bruce F. Kitchens
13 Aug 2009
SBW 8113107

TO: RM #25, Jim Tompkins/Mickie Walters
Herbicide Branch (7505P)
Registration Division (7505P)

INTRODUCTION:

The registrant, Repar-Glypho, LLC, is submitting an application to register the proposed end-use product, Glyphosate 41% SL. The active ingredients in this product is Glyphosate isopropylamine salt (% a.i.) at a label nominal concentration of 41% a.i. In addition, the registrant states that this product is substantially similar to EPA Reg. No. 524-445 Roundup Original Herbicide. This product is intended for use as a food and nonfood use herbicide. In support of this request, the registrant has submitted a basic Confidential Statement of Formula (CSF) dated 09 Feb 2009, a draft label, and product chemistry data contained in MRID# 476851-01. During the course of this review, the registrant submitted a revised basic CSF dated 13 Aug 2009 to correct an error. The Technical Review Branch (TRB) has been asked to review this submission.

SUMMARY OF FINDINGS

TRB has reviewed this submission and reports the following findings:

1. This product is produced from an integrated formulation system. This means that the source of the active ingredient is the result of intended chemical reactions. The source of the active ingredient is currently undergoing registration.
2. All inert ingredients are approved for use in formulated pesticide products. In addition, all inert ingredients are exempt from the requirement of a food tolerance when applied to growing crops.
3. The nominal concentration of the active ingredient listed on the proposed basic CSF and the draft label are the same.
4. The draft label contains the appropriate storage and disposal statements. In addition, the draft label contains the appropriate physical and chemical hazard statement prohibiting mixing, storage, and application of this product in galvanized steel or unlined steel containers or spray tanks. This product reacts with these containers to produce hydrogen gas which may form a highly combustible mixture. This mixture could explode or flash if ignited by open flame, spark, cigarette or other ignition source.
5. The active ingredient's certified limits as proposed on the basic CSF are acceptable.

6. The comparison of the proposed and cited product reveals that both products have the same active ingredient, the same active ingredient nominal concentration, and similar inert ingredients.

CONCLUSIONS:

TRB has reviewed this submission and concludes the following:

1. The basic formula CSF for the proposed end-use product, Glyphosate 41% SL dated 13 Aug 2009 is acceptable. Acceptance of this product for registration is contingent upon registration of the source of the active ingredient.
2. This submission (MRID# 476851-01) satisfies the data requirements as specified in 40 CFR 158.320, 158.325, 158.335, 158.340, 158.350, and 158.355 with respect to product identity and composition, description of materials used to produce the product, description of formulation process, discussion of formation of impurities, certified limits, and enforcement analytical method.
3. Except for storage stability/corrosion characteristics studies, the remaining product chemistry Group B data (MRID# 476851-01) adequately fulfill the data requirements specified in 40 CFR 158.310 with respect to physical and chemical properties. Inform the registrant that 1-year studies are required to satisfy the storage stability and corrosion characteristics data requirements.
4. The proposed product has been determined to be substantially similar in composition to EPA Reg. No. 534-445 Roundup Original Herbicide from a product chemistry standpoint.

PRODUCT CHEMISTRY DATA (SERIES 830 Subgroup A)

Subgroup A – Product Identity and Composition	<u>Data Required</u> <u>Fulfilled</u>	<u>MRID No.</u>
830.1550. Chemical Identity	Y	476851-01
830.1600. Beginning Materials	Y	476851-01
830.1650. Formulation Process	Y	476851-01
830.1670. Discussion of Impurities	Y	476851-01
830.1700. Preliminary Analysis	NA	
830.1750. Certified Limits	Y	see csf 2/9/09
830.1800. Enforcement Analytical Method	Y	476851-01

PRODUCT CHEMISTRY DATA (SERIES 830 Subgroup B)

Subgroup B – Physical and Chemical Properties	<u>Data Required Fulfilled</u>	<u>Value or Qualitative Descrip.</u>	<u>MRID No.</u>
830.6302. Color	Y	Amber	476851-01
830.6303. Physical State	Y	Liquid	476851-01
830.6304. Odor	Y	Characteristic odor	476851-01
830.6314. Oxidation/Reduction Action	NA	Product does not contain oxidizing or reducing agents	
830.6315. Flammability	NA	Product does not contain flammable components	
830.6316. Explodability	NA	Product does not contain explosive components	
830.6317. Storage stability	G		
830.6319. Miscibility	NA	Product not an EC nor mixed with petroleum solvents	
830.6320. Corrosion Characteristics	G		
830.6321. Dielectric Breakdown Voltage	NA	Product not used around electrical equipment	
830.7000. pH	Y	4.7	476851-01
830.7100. Viscosity	Y	48.6cP @ 25°C 20.6cP @ 40°C	476851-01
830.7000. Density/Bulk Density	Y	1.16 @ 25°C (specific gravity)	476851-01

Explanations: A = The Requirements Were Fulfilled; N = The Requirements Were Not Fulfilled; NA = Not Applicable; G = Data Gap; U = Requires Upgrading; I = Incomplete or In Progress; W = Waived.

Analytical Method: (MRID No. 476851-01)

The active ingredient, Glyphosate isopropylamine salt, was determined by high performance liquid chromatography (HPLC) using ultraviolet detection (UV) AT 195 nm and an external standard. Results of method validation not reported in this submission.

Equipment and Parameters

Instrument: High performance liquid chromatograph (HPLC)
Detector: Ultraviolet detector (UV)
Wavelength: 195 nm
Column: Whatman Partisil 10 SAX 250 x 4.6 mm i.d. stainless steel packed with a strong ion exchange resin or equivalent
Column Temperature: Ambient
Sample Size: 50 µl
Mobile Phase: Water:Methanol 9.6/4 with 0.8437g of KH₂PO₄ adjust pH to 1.9 with 85% phosphoric acid, filter and degas before use
Flow Rate: 2.3 ml/min
Pressure: ≈ 8.0 MPa
Retention Time: 2.5-4.0 minutes



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

February 26, 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

MANDAVA ASSOCIATES, LLC
REPAR-GLYPHO, LLC
1050 CONN. AVE., NW, STE 1000
WASHINGTON, DC 20036-

Report of Analysis for Compliance with PR Notice 86-5

Thank you for your submittal of 23-FEB-09. Our staff has completed a preliminary analysis of the material. The results are provided as follows:

Your submittal was found to be in full compliance with the standards for submission of data contained in PR Notice 86-5. A copy of your bibliography is enclosed, annotated with Master Record ID's (MRIDs) assigned to each document submitted. Please use these numbers in all future references to these documents. Thank you for your cooperation. If you have any questions concerning this data submission, please raise them with the cognizant Product Manager, to whom the data have been released.